

Bis(μ -naphthalene-1,8-dicarboxylato- $\kappa^2 O^1:O^8$)bis[aquabis(N,N' -dimethylformamide- κO)copper(II)]

Jun-Dan Fu, Chun-Yan Zhang, Qing-Yu Shi and Yi-Hang Wen*

Zhejiang Key Laboratory for Reactive Chemistry on Solid Surfaces, Institute of Physical Chemistry, Zhejiang Normal University, Jinhua, Zhejiang 321004, People's Republic of China

Correspondence e-mail: wyh@zjnu.edu.cn


Received 17 June 2010; accepted 16 July 2010

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.027; wR factor = 0.076; data-to-parameter ratio = 17.9.

In the centrosymmetric dinuclear title complex, $[Cu_2(C_{12}H_6O_4)_2(C_3H_7NO)_4(H_2O)_2]$, the coordination environment of each Cu(II) atom displays a distorted CuO_5 square-pyramidal geometry, which is formed by two carboxylate O atoms of two μ -1,8-nap ligands (1,8-nap is naphthalene-1,8-dicarboxylate), two O atoms of two DMF (DMF is N,N' -dimethylformamide) and one coordinated water molecule. The Cu—O distances involving the four O atoms in the square plane are in the range 1.9501 (11)–1.9677 (11) Å, with the Cu atom lying nearly in the plane [deviation = 0.0726 (2) Å]. The axial O atom occupies the peak position with a Cu—O distance of 2.885 (12) Å, which is significantly longer than the rest of the Cu—O distances. Each 1,8-nap ligand acts as bridge, linking two Cu^{II} atoms into a dinuclear structure. Intermolecular O—H···O and C—H···O hydrogen-bonding interactions consolidate the structure.

Related literature

For the coordination modes of the 1,8-nap ligand, see: Wen *et al.* (2007, 2008). For related complexes, see: Abourahma *et al.* (2002); Bencini *et al.* (2003); Fokin *et al.* (2004); Sun *et al.* (2009).



Experimental

Crystal data

$[Cu_2(C_{12}H_6O_4)_2(C_3H_7NO)_4 \cdot (H_2O)_2]$	$\beta = 102.249$ (2)°
$M_r = 883.83$	$V = 3948.00$ (13) Å ³
Monoclinic, $C2/c$	$Z = 4$
$a = 17.7078$ (4) Å	Mo $K\alpha$ radiation
$b = 9.9025$ (1) Å	$\mu = 1.15$ mm ⁻¹
$c = 23.0393$ (5) Å	$T = 296$ K
	0.40 × 0.26 × 0.13 mm

Data collection

Bruker APEXII area-detector diffractometer	29673 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	4625 independent reflections
$T_{\min} = 0.71$, $T_{\max} = 0.86$	4076 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.076$	$\Delta\rho_{\max} = 0.28$ e Å ⁻³
$S = 1.04$	$\Delta\rho_{\min} = -0.27$ e Å ⁻³
4625 reflections	
259 parameters	
5 restraints	

Table 1
Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
O1W—H1WA···O5	0.82 (2)	1.82 (2)	2.642 (2)	178 (2)
O1W—H1WB···O4 ⁱ	0.80 (1)	1.82 (2)	2.623 (2)	175 (2)
C3—H3A···O1 ⁱⁱ	0.93	2.49	3.396 (3)	164
C13—H13A···O3	0.93	2.59	3.140 (2)	119
C17—H17A···O6 ⁱⁱⁱ	0.96	2.51	3.424 (3)	159

Symmetry codes: (i) $-x + \frac{1}{2}, -y + \frac{3}{2}, -z + 1$; (ii) $x + \frac{1}{2}, y + \frac{1}{2}, z$; (iii) $x - \frac{1}{2}, y - \frac{1}{2}, z$.

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2300).

References

- Abourahma, H., Moulton, B., Kravtsov, V. & Zaworotko, J. M. (2002). *J. Am. Chem. Soc.* **124**, 9990–9991.
- Bencini, A., Dei, A., Sangregorio, C., Totti, F. & Vaz, M. G. F. (2003). *Inorg. Chem.* **42**, 8065–8071.
- Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fokin, S., Ovcharenko, V., Romanenko, G. & Ikorskii, V. (2004). *Inorg. Chem.* **43**, 969–977.
- Sheldrick, G. M. (1996). *SADABS*. University of Gottingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sun, C. Y., Liu, S. X., Liang, D. D., Shao, K. Z., Ren, Y. H. & Su, Z. M. (2009). *J. Am. Chem. Soc.* **131**, 1883–1888.
- Wen, Y. H., Feng, X., Feng, Y. L. & Tang, Z. W. (2008). *Inorg. Chem. Commun.* **11**, 659–661.
- Wen, Y.-H., Feng, X., He, Y.-H., Lan, Y.-Z. & Sun, H. (2007). *Acta Cryst. C* **63**, m504–m506.

supporting information

Acta Cryst. (2010). E66, m1003–m1004 [https://doi.org/10.1107/S1600536810028497]

Bis(μ -naphthalene-1,8-dicarboxylato- $\kappa^2O^1:O^8$)bis[aquabis(N,N'-dimethyl-formamide- κO)copper(II)]

Jun-Dan Fu, Chun-Yan Zhang, Qing-Yu Shi and Yi-Hang Wen

S1. Comment

It is well-known that appropriate metal and ligand are the two keys for design and construction of metal-organic frameworks. Here we choose 1,8-nap ligand (1,8-nap = naphthalene-1,8-dicarboxylate), due to its unique ability to form stable chelates in diverse coordination modes such as bidentate, meridian and bridging; which have been demonstrated in our previous work (Wen *et al.*, 2008; Wen *et al.*, 2007). Moreover, we select the copper to provide a set of well defined coordination geometry. As a result, we have prepared the title complex, Cu₂(1,8-nap)₂(DMF)₄(H₂O)₂, (I), a new dinuclear Cu^{II} compound based on 1,8-nap ligand.


A perspective view of the molecular structure of (I) is presented in Fig. 1. The coordination environment of each Cu atom displays a distorted CuO₅ square pyramidal coordination geometry, which is formed from two carboxylate oxygen atoms of two μ_2 -1,8-nap ligands, two oxygen atoms of two DMF and one coordinated water molecule; similar to some complexes reported earlier (Abourahma *et al.*, 2002; Bencini *et al.*, 2003; Fokin *et al.*, 2004; Sun *et al.*, 2009). Four oxygen atoms O₂-O₃ⁱ-O₁W-O₆ form a square plane (Cu—O distances in ranging 1.9501 (11) - 1.9677 (11) Å), and the Cu1 atom lies in the plane (deviation 0.0726 (2) Å). The fifth oxygen atom O₁ is on the peak of square pyramid, and Cu—O distance is 2.885 (12) Å, which is significantly longer than the rest of the Cu—O distances. Both carboxylate groups of the 1,8-nap ligand are deprotonated, and adopt a monodentate coordination mode. As a result, the whole 1,8-nap ligand acts as μ_2 -bridge linking two Cu^{II} atoms to form a sixteen-atoms ring. There are intramolecular hydrogen bonds between uncoordinated O atoms of 1,8-nap ligands and water molecules, O₁W···O₅, O₁W···O₄ⁱ(details are given in Tab. 1). In addition, weak interactions of the type C—H···O are also present. Such hydrogen-bonding interactions consolidate the dinuclear structure, as depicted in Fig. 2.

S2. Experimental


A mixture of naphthalene-1,8-dicarboxylate anhydride (0.1981 g, 1 mmol), CuCl₂.2H₂O (0.085 g, 0.5 mmol) and Na₂CO₃(0.053 g, 0.5 mmol) was dissolved in a mixed solution of DMF-H₂O (1:2 v/v, 25 ml) and stirred at 343 K for 2 h. The filtrate was allowed to stand at ambient temperature. Well formed blue crystals suitable for X-ray analysis were obtained after two months (yield 45%, based on Cu).

S3. Refinement


H atoms bonded to C atoms were positioned geometrically and included in the refinement in the riding-model approximation [C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$] and methyl groups were allowed to rotate to fit the electron density [C—H = 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$]. Water H atoms were located and refined with distance restraints of O—H = 0.85 (2) Å and H···H = 1.35 (2) Å, with displacement parameters set at 1.5 $U_{\text{eq}}(\text{O})$.

**Figure 1**

Perspective view of the structure of (I), showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. [Symmetry codes: (i)- $x + 1/2, -y + 3/2, -z + 1$]

**Figure 2**

A packing diagram of (I) with the hydrogen-bonding interaction depicted by dash lines.

Bis(μ -naphthalene-1,8-dicarboxylato- $\kappa^2O^1:O^8$)bis[aquabis(N,N' - dimethylformamide- κO)copper(II)]*Crystal data*

$M_r = 883.83$

Monoclinic, $C2/c$

Hall symbol: -C 2yc

$a = 17.7078 (4)$ Å

$b = 9.9025 (1)$ Å

$c = 23.0393 (5)$ Å

$\beta = 102.249 (2)^\circ$

$V = 3948.00 (13)$ Å³

$Z = 4$

$F(000) = 1832$

$D_x = 1.487 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9928 reflections

$\theta = 2.4\text{--}27.7^\circ$

$\mu = 1.15 \text{ mm}^{-1}$

$T = 296$ K

Block, blue

$0.40 \times 0.26 \times 0.13$ mm

Data collection

Bruker APEXII area-detector

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.71$, $T_{\max} = 0.86$

29673 measured reflections

4625 independent reflections

4076 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\max} = 27.7^\circ$, $\theta_{\min} = 2.4^\circ$

$h = -23 \rightarrow 23$

$k = -12 \rightarrow 12$

$l = -28 \rightarrow 30$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.027$

$wR(F^2) = 0.076$

$S = 1.04$

4625 reflections

259 parameters

5 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0415P)^2 + 2.4079P]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.002$

$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.124473 (10)	0.670966 (18)	0.464441 (8)	0.03052 (7)
C1	0.33281 (9)	0.85492 (16)	0.40975 (7)	0.0341 (3)
C2	0.31963 (10)	0.94374 (17)	0.35537 (7)	0.0395 (4)

C3	0.38265 (12)	1.0098 (2)	0.34354 (9)	0.0543 (5)
H3A	0.4291	1.0076	0.3714	0.065*
C4	0.37868 (14)	1.0808 (2)	0.29013 (11)	0.0684 (6)
H4A	0.4219	1.1266	0.2834	0.082*
C5	0.31253 (14)	1.0826 (2)	0.24877 (10)	0.0627 (6)
H5A	0.3110	1.1271	0.2130	0.075*
C6	0.24541 (13)	1.01821 (18)	0.25878 (8)	0.0485 (4)
C7	0.24722 (10)	0.95047 (16)	0.31396 (7)	0.0381 (3)
C8	0.17617 (14)	1.0204 (2)	0.21466 (8)	0.0568 (5)
H8A	0.1752	1.0654	0.1791	0.068*
C9	0.11186 (13)	0.9584 (2)	0.22331 (8)	0.0571 (5)
H9A	0.0680	0.9554	0.1929	0.069*
C10	0.11117 (11)	0.89818 (19)	0.27848 (8)	0.0460 (4)
H10A	0.0660	0.8583	0.2846	0.055*
C11	0.17582 (10)	0.89708 (16)	0.32346 (7)	0.0367 (3)
C12	0.16459 (9)	0.85435 (16)	0.38382 (7)	0.0334 (3)
C13	0.09853 (12)	0.4309 (2)	0.38365 (9)	0.0543 (5)
H13A	0.1348	0.4846	0.3707	0.065*
C14	0.1278 (3)	0.2623 (3)	0.31590 (17)	0.1263 (15)
H14A	0.1622	0.3321	0.3085	0.189*
H14B	0.0923	0.2401	0.2796	0.189*
H14C	0.1572	0.1836	0.3310	0.189*
C15	0.02931 (19)	0.2191 (3)	0.37687 (13)	0.0902 (9)
H15A	0.0055	0.2632	0.4055	0.135*
H15B	0.0551	0.1388	0.3940	0.135*
H15C	-0.0096	0.1955	0.3426	0.135*
C16	-0.03349 (10)	0.72571 (17)	0.46612 (8)	0.0418 (4)
H16A	-0.0370	0.6338	0.4735	0.050*
C17	-0.16911 (14)	0.7426 (3)	0.46719 (17)	0.0956 (10)
H17A	-0.1636	0.6470	0.4735	0.143*
H17B	-0.2080	0.7597	0.4320	0.143*
H17C	-0.1841	0.7838	0.5007	0.143*
C18	-0.09461 (13)	0.9433 (2)	0.44989 (12)	0.0661 (6)
H18A	-0.0439	0.9698	0.4457	0.099*
H18B	-0.1075	0.9902	0.4829	0.099*
H18C	-0.1315	0.9656	0.4143	0.099*
O1W	0.22801 (6)	0.59540 (12)	0.46744 (5)	0.0358 (2)
H1WA	0.2520 (12)	0.641 (2)	0.4477 (8)	0.054*
H1WB	0.2520 (12)	0.589 (2)	0.5009 (7)	0.054*
O1	0.06739 (7)	0.47781 (13)	0.42143 (6)	0.0511 (3)
O2	0.02961 (7)	0.77247 (12)	0.46251 (6)	0.0434 (3)
O3	0.13036 (6)	0.74302 (12)	0.38655 (5)	0.0389 (2)
O4	0.18619 (7)	0.93436 (12)	0.42557 (5)	0.0407 (3)
O5	0.30792 (7)	0.73769 (11)	0.40428 (5)	0.0389 (3)
O6	0.37233 (6)	0.90670 (12)	0.45727 (5)	0.0395 (3)
N1	0.08503 (14)	0.30962 (18)	0.35940 (9)	0.0687 (5)
N2	-0.09595 (8)	0.79924 (16)	0.46006 (7)	0.0458 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.02809 (11)	0.02933 (11)	0.03456 (11)	0.00029 (7)	0.00758 (7)	-0.00087 (7)
C1	0.0308 (7)	0.0369 (8)	0.0381 (8)	0.0040 (6)	0.0154 (6)	0.0037 (6)
C2	0.0456 (9)	0.0357 (8)	0.0405 (8)	0.0005 (7)	0.0167 (7)	0.0047 (7)
C3	0.0534 (11)	0.0562 (12)	0.0563 (11)	-0.0080 (9)	0.0181 (9)	0.0110 (9)
C4	0.0724 (15)	0.0669 (14)	0.0723 (15)	-0.0164 (12)	0.0299 (12)	0.0220 (12)
C5	0.0888 (16)	0.0545 (12)	0.0502 (11)	-0.0057 (11)	0.0271 (11)	0.0172 (9)
C6	0.0709 (12)	0.0388 (9)	0.0386 (9)	0.0042 (8)	0.0179 (8)	0.0068 (7)
C7	0.0525 (10)	0.0301 (7)	0.0338 (8)	0.0041 (7)	0.0137 (7)	0.0039 (6)
C8	0.0837 (15)	0.0503 (11)	0.0345 (9)	0.0080 (10)	0.0086 (9)	0.0115 (8)
C9	0.0696 (13)	0.0567 (11)	0.0391 (10)	0.0086 (10)	-0.0020 (9)	0.0083 (9)
C10	0.0512 (10)	0.0445 (10)	0.0388 (9)	0.0060 (8)	0.0018 (7)	0.0035 (7)
C11	0.0460 (9)	0.0296 (8)	0.0339 (8)	0.0074 (6)	0.0075 (7)	0.0016 (6)
C12	0.0294 (7)	0.0356 (7)	0.0348 (8)	0.0099 (5)	0.0057 (6)	0.0046 (6)
C13	0.0593 (12)	0.0406 (10)	0.0641 (12)	-0.0089 (9)	0.0157 (10)	-0.0125 (9)
C14	0.204 (4)	0.0723 (19)	0.129 (3)	-0.004 (2)	0.093 (3)	-0.0396 (19)
C15	0.120 (2)	0.0554 (14)	0.094 (2)	-0.0337 (16)	0.0198 (17)	-0.0182 (14)
C16	0.0412 (9)	0.0345 (8)	0.0528 (10)	0.0041 (7)	0.0171 (8)	0.0013 (7)
C17	0.0436 (12)	0.0725 (17)	0.180 (3)	0.0018 (12)	0.0436 (16)	0.0084 (19)
C18	0.0562 (12)	0.0503 (12)	0.0948 (17)	0.0194 (10)	0.0228 (12)	0.0182 (11)
O1W	0.0315 (5)	0.0401 (6)	0.0366 (6)	0.0036 (4)	0.0088 (4)	0.0051 (5)
O1	0.0484 (7)	0.0433 (7)	0.0632 (8)	-0.0096 (6)	0.0151 (6)	-0.0163 (6)
O2	0.0338 (6)	0.0384 (6)	0.0601 (8)	0.0058 (5)	0.0147 (5)	0.0016 (6)
O3	0.0368 (6)	0.0414 (6)	0.0378 (6)	0.0003 (5)	0.0067 (5)	0.0057 (5)
O4	0.0430 (6)	0.0438 (6)	0.0348 (6)	0.0076 (5)	0.0069 (5)	-0.0017 (5)
O5	0.0431 (6)	0.0337 (6)	0.0427 (6)	0.0024 (5)	0.0156 (5)	0.0040 (5)
O6	0.0365 (6)	0.0442 (6)	0.0388 (6)	-0.0036 (5)	0.0103 (5)	0.0041 (5)
N1	0.0966 (15)	0.0409 (9)	0.0714 (12)	-0.0079 (9)	0.0241 (11)	-0.0189 (8)
N2	0.0344 (7)	0.0431 (8)	0.0626 (10)	0.0063 (6)	0.0166 (7)	0.0011 (7)

Geometric parameters (\AA , $^\circ$)

Cu1—O2	1.9500 (11)	C12—O3	1.266 (2)
Cu1—O6 ⁱ	1.9505 (11)	C13—O1	1.217 (2)
Cu1—O3	1.9544 (11)	C13—N1	1.324 (2)
Cu1—O1W	1.9678 (11)	C13—H13A	0.9300
Cu1—O1	2.2885 (12)	C14—N1	1.456 (3)
C1—O5	1.2386 (19)	C14—H14A	0.9600
C1—O6	1.275 (2)	C14—H14B	0.9600
C1—C2	1.508 (2)	C14—H14C	0.9600
C2—C3	1.370 (2)	C15—N1	1.452 (3)
C2—C7	1.428 (2)	C15—H15A	0.9600
C3—C4	1.406 (3)	C15—H15B	0.9600
C3—H3A	0.9300	C15—H15C	0.9600
C4—C5	1.344 (3)	C16—O2	1.228 (2)
C4—H4A	0.9300	C16—N2	1.307 (2)

C5—C6	1.410 (3)	C16—H16A	0.9300
C5—H5A	0.9300	C17—N2	1.453 (3)
C6—C8	1.417 (3)	C17—H17A	0.9600
C6—C7	1.432 (2)	C17—H17B	0.9600
C7—C11	1.430 (2)	C17—H17C	0.9600
C8—C9	1.345 (3)	C18—N2	1.447 (3)
C8—H8A	0.9300	C18—H18A	0.9600
C9—C10	1.406 (3)	C18—H18B	0.9600
C9—H9A	0.9300	C18—H18C	0.9600
C10—C11	1.371 (2)	O1W—H1WA	0.82 (2)
C10—H10A	0.9300	O1W—H1WB	0.80 (1)
C11—C12	1.507 (2)	O6—Cu1 ⁱ	1.9505 (11)
C12—O4	1.242 (2)		
O2—Cu1—O6 ⁱ	94.60 (5)	O3—C12—C11	116.67 (14)
O2—Cu1—O3	90.25 (5)	O1—C13—N1	125.4 (2)
O6 ⁱ —Cu1—O3	175.05 (5)	O1—C13—H13A	117.3
O2—Cu1—O1W	171.31 (5)	N1—C13—H13A	117.3
O6 ⁱ —Cu1—O1W	88.48 (5)	N1—C14—H14A	109.5
O3—Cu1—O1W	86.58 (5)	N1—C14—H14B	109.5
O2—Cu1—O1	96.99 (5)	H14A—C14—H14B	109.5
O6 ⁱ —Cu1—O1	89.67 (5)	N1—C14—H14C	109.5
O3—Cu1—O1	90.71 (5)	H14A—C14—H14C	109.5
O1W—Cu1—O1	91.15 (5)	H14B—C14—H14C	109.5
O5—C1—O6	125.65 (15)	N1—C15—H15A	109.5
O5—C1—C2	118.24 (15)	N1—C15—H15B	109.5
O6—C1—C2	116.03 (14)	H15A—C15—H15B	109.5
C3—C2—C7	119.91 (16)	N1—C15—H15C	109.5
C3—C2—C1	117.06 (16)	H15A—C15—H15C	109.5
C7—C2—C1	122.75 (14)	H15B—C15—H15C	109.5
C2—C3—C4	121.4 (2)	O2—C16—N2	122.97 (16)
C2—C3—H3A	119.3	O2—C16—H16A	118.5
C4—C3—H3A	119.3	N2—C16—H16A	118.5
C5—C4—C3	120.1 (2)	N2—C17—H17A	109.5
C5—C4—H4A	120.0	N2—C17—H17B	109.5
C3—C4—H4A	120.0	H17A—C17—H17B	109.5
C4—C5—C6	121.06 (18)	N2—C17—H17C	109.5
C4—C5—H5A	119.5	H17A—C17—H17C	109.5
C6—C5—H5A	119.5	H17B—C17—H17C	109.5
C5—C6—C8	120.45 (17)	N2—C18—H18A	109.5
C5—C6—C7	119.75 (19)	N2—C18—H18B	109.5
C8—C6—C7	119.79 (18)	H18A—C18—H18B	109.5
C2—C7—C11	125.34 (14)	N2—C18—H18C	109.5
C2—C7—C6	117.57 (16)	H18A—C18—H18C	109.5
C11—C7—C6	117.08 (16)	H18B—C18—H18C	109.5
C9—C8—C6	121.18 (17)	Cu1—O1W—H1WA	110.9 (15)
C9—C8—H8A	119.4	Cu1—O1W—H1WB	111.5 (16)
C6—C8—H8A	119.4	H1WA—O1W—H1WB	110.4 (19)

C8—C9—C10	119.82 (18)	C13—O1—Cu1	113.79 (12)
C8—C9—H9A	120.1	C16—O2—Cu1	126.56 (11)
C10—C9—H9A	120.1	C12—O3—Cu1	118.88 (10)
C11—C10—C9	121.36 (19)	C1—O6—Cu1 ⁱ	122.57 (10)
C11—C10—H10A	119.3	C13—N1—C15	121.0 (2)
C9—C10—H10A	119.3	C13—N1—C14	120.5 (2)
C10—C11—C7	120.33 (15)	C15—N1—C14	118.4 (2)
C10—C11—C12	116.57 (15)	C16—N2—C18	121.61 (16)
C7—C11—C12	122.64 (14)	C16—N2—C17	121.86 (18)
O4—C12—O3	126.08 (15)	C18—N2—C17	116.40 (17)
O4—C12—C11	117.15 (14)		
O5—C1—C2—C3	-129.54 (18)	C2—C7—C11—C12	14.1 (2)
O6—C1—C2—C3	47.3 (2)	C6—C7—C11—C12	-164.70 (15)
O5—C1—C2—C7	44.3 (2)	C10—C11—C12—O4	-124.12 (16)
O6—C1—C2—C7	-138.81 (16)	C7—C11—C12—O4	48.1 (2)
C7—C2—C3—C4	-2.4 (3)	C10—C11—C12—O3	52.4 (2)
C1—C2—C3—C4	171.7 (2)	C7—C11—C12—O3	-135.41 (15)
C2—C3—C4—C5	-1.4 (4)	N1—C13—O1—Cu1	168.26 (19)
C3—C4—C5—C6	2.4 (4)	O2—Cu1—O1—C13	135.72 (15)
C4—C5—C6—C8	-179.7 (2)	O6 ⁱ —Cu1—O1—C13	-129.68 (15)
C4—C5—C6—C7	0.5 (3)	O3—Cu1—O1—C13	45.38 (15)
C3—C2—C7—C11	-173.81 (17)	O1W—Cu1—O1—C13	-41.21 (15)
C1—C2—C7—C11	12.5 (3)	N2—C16—O2—Cu1	-174.76 (13)
C3—C2—C7—C6	5.0 (2)	O6 ⁱ —Cu1—O2—C16	-59.16 (15)
C1—C2—C7—C6	-168.68 (15)	O3—Cu1—O2—C16	121.82 (15)
C5—C6—C7—C2	-4.1 (3)	O1—Cu1—O2—C16	31.08 (15)
C8—C6—C7—C2	176.05 (17)	O4—C12—O3—Cu1	-10.9 (2)
C5—C6—C7—C11	174.82 (17)	C11—C12—O3—Cu1	172.91 (10)
C8—C6—C7—C11	-5.0 (2)	O2—Cu1—O3—C12	87.85 (11)
C5—C6—C8—C9	179.4 (2)	O1W—Cu1—O3—C12	-84.05 (11)
C7—C6—C8—C9	-0.8 (3)	O1—Cu1—O3—C12	-175.16 (11)
C6—C8—C9—C10	4.5 (3)	O5—C1—O6—Cu1 ⁱ	-18.5 (2)
C8—C9—C10—C11	-2.2 (3)	C2—C1—O6—Cu1 ⁱ	164.90 (10)
C9—C10—C11—C7	-3.8 (3)	O1—C13—N1—C15	0.0 (4)
C9—C10—C11—C12	168.60 (17)	O1—C13—N1—C14	-177.9 (3)
C2—C7—C11—C10	-173.94 (16)	O2—C16—N2—C18	-1.4 (3)
C6—C7—C11—C10	7.2 (2)	O2—C16—N2—C17	-177.2 (2)

Symmetry code: (i) $-x+1/2, -y+3/2, -z+1$.

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1W—H1WA ⁱⁱ —O5	0.82 (2)	1.82 (2)	2.642 (2)	178 (2)
O1W—H1WB ⁱⁱ —O4 ⁱ	0.80 (1)	1.82 (2)	2.623 (2)	175 (2)
C3—H3A ⁱⁱ —O1 ⁱⁱ	0.93	2.49	3.396 (3)	164

C13—H13A···O3	0.93	2.59	3.140 (2)	119
C17—H17A···O6 ⁱⁱⁱ	0.96	2.51	3.424 (3)	159

Symmetry codes: (i) $-x+1/2, -y+3/2, -z+1$; (ii) $x+1/2, y+1/2, z$; (iii) $x-1/2, y-1/2, z$.