

(5-Bromo-2-methylphenyl)(4-ethoxyphenyl)methanone

Yong-Heng Shi,^a Gui-Long Zhao,^a Hua Shao,^a Wei Liu^a
and Wei-Ren Xu^{a,b*}

^aTianjin Key Laboratory of Molecular Design and Drug Discovery, Tianjin Institute of Pharmaceutical Research, Tianjin 300193, People's Republic of China, and ^bSchool of Pharmacy, Tianjin Medical University, Tianjin 300070, People's Republic of China

Correspondence e-mail: weirun_xu@yahoo.cn


Received 7 July 2010; accepted 9 July 2010

Key indicators: single-crystal X-ray study; $T = 113\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.041; wR factor = 0.107; data-to-parameter ratio = 14.2.

In the title compound, $\text{C}_{16}\text{H}_{15}\text{BrO}_2$, the dihedral angle between the benzene rings is $68.5(2)^\circ$. In the crystal structure, molecules are linked by weak C–H···O hydrogen bonds into chains parallel to the b axis.

Related literature

For details of the biological activity of SGLT2 inhibitors, see: Meng *et al.* (2008). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{15}\text{BrO}_2$

$M_r = 319.19$

Orthorhombic, $Pbca$	$Z = 8$
$a = 9.5730(19)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 13.188(3)\text{ \AA}$	$\mu = 2.93\text{ mm}^{-1}$
$c = 22.205(4)\text{ \AA}$	$T = 113\text{ K}$
$V = 2803.4(10)\text{ \AA}^3$	$0.30 \times 0.20 \times 0.16\text{ mm}$

Data collection

Rigaku Saturn CCD area-detector diffractometer	17506 measured reflections
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	2479 independent reflections
$R_{\text{int}} = 0.044$	2133 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.474$, $T_{\max} = 0.652$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$	175 parameters
$wR(F^2) = 0.107$	H-atom parameters constrained
$S = 1.10$	$\Delta\rho_{\max} = 0.84\text{ e \AA}^{-3}$
2479 reflections	$\Delta\rho_{\min} = -0.57\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C11–H11···O1 ⁱ	0.95	2.42	3.313 (3)	156
Symmetry code: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, z$.				

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2475).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Meng, M., Ellsworth, B. A., Nirschl, A. A., McCann, P. J., Patel, M., Girotra, R. N., Wu, G., Sher, P. M., Morrison, E. P., Biller, S. A., Zahler, R., Deshpande, P. P., Pullockaran, A., Hagan, D. L., Morgan, N., Taylor, J. R., Obermeier, M. T., Humphreys, W. G., Khanna, A., Discenza, L., Robertson, J. M., Wang, A., Han, S., Wetterau, J. R., Janovitz, E. B., Flint, O. P., Whaley, J. M. & Washburn, W. N. (2008). *J. Med. Chem.* **51**, 1145–1149.
- Rigaku. (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supporting information

Acta Cryst. (2010). E66, o2038 [https://doi.org/10.1107/S1600536810027327]

(5-Bromo-2-methylphenyl)(4-ethoxyphenyl)methanone

Yong-Heng Shi, Gui-Long Zhao, Hua Shao, Wei Liu and Wei-Ren Xu

S1. Comment

Dapagliflozin is an anti-diabetic agent through the inhibition of renal SGLT2, which was developed by Bristol-Myers Squibb Company and is now in the phase III clinical trial (Meng *et al.*, 2008). During the discovery of our own SGLT2 inhibitors as anti-diabetic agents, we prepared the derivatives of dapagliflozin (Meng *et al.*, 2008) for biological evaluation, and the title compound, (5-bromo-2-methylphenyl)(4-ethoxyphenyl)methanone, was prepared as an important intermediate.


In title compound, $C_{16}H_{15}BrO_2$, bond lengths are normal (Allen *et al.*, 1987)). The dihedral angle between the benzene rings ($C2—C7$ and $C9—C14$) is $68.5(2)^\circ$. In the crystal structure, molecules interact through weak $C—H\cdots O$ hydrogen bonds to form chains parallel to the b axis.

S2. Experimental

A round-bottomed flask was charged with 2.15 g (10 mmol) of 5-bromo-2-methylbenoic acid, 1 drop of DMF, 1.27 g (10 mmol) of oxalyl chloride and 3 ml of dried dichloromethane, and the mixture was stirred at room temperature over night until a clear solution formed. The reaction mixture was evaporated on a rotary evaporator to give crude 5-bromo-2-chlorobenzoyl chloride, which was dissolved in 15 ml of dried dichloromethane. The solution thus obtained was stirred while being cooled with an ice-salt bath, and 1.22 g (10 mmol) of phenetole was added followed by the addition of 1.60 g (12 mmol) of anhydrous aluminium chloride in a portionwise manner. The resulting mixture was stirred at this temperature for 1 h and poured into 150 ml of ice-water. The mixture formed was extracted with three 50 ml portions of dichloromethane, and the combined extracts were washed with saturated brine, dried over sodium sulfate and evaporated on a rotary evaporator to afford the crude title compound. Pure title compound was obtained by column chromatography. Crystals suitable for X-ray diffraction were obtained through slow evaporation of a solution of the pure title compound in ethyl acetate/petroleum ether (1/5 v/v).

S3. Refinement

All H atoms were found on difference Fourier maps, and included in the final cycles of refinement using a riding model, with $C—H = 0.95–0.99 \text{ \AA}$ and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aryl and methylene H atoms and $1.5U_{\text{eq}}(\text{C})$ for the methyl H atoms.

**Figure 1**

The molecular structure of the title compound, with displacement ellipsoids drawn at the 40% probability level.

(5-Bromo-2-methylphenyl)(4-ethoxyphenyl)methanone

Crystal data

$C_{16}H_{15}BrO_2$
 $M_r = 319.19$
Orthorhombic, $Pbca$
Hall symbol: -P 2ac 2ab
 $a = 9.5730 (19)$ Å
 $b = 13.188 (3)$ Å
 $c = 22.205 (4)$ Å
 $V = 2803.4 (10)$ Å³
 $Z = 8$

$F(000) = 1296$
 $D_x = 1.513 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 6268 reflections
 $\theta = 2.1\text{--}27.9^\circ$
 $\mu = 2.93 \text{ mm}^{-1}$
 $T = 113 \text{ K}$
Block, colorless
 $0.30 \times 0.20 \times 0.16$ mm

Data collection

Rigaku Saturn CCD area-detector
diffractometer
Radiation source: rotating anode
Confocal monochromator
Detector resolution: 7.31 pixels mm⁻¹
 ω and φ scans
Absorption correction: multi-scan
(*CrystalClear*; Rigaku, 2005)
 $T_{\min} = 0.474$, $T_{\max} = 0.652$

17506 measured reflections
2479 independent reflections
2133 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.8^\circ$
 $h = -11 \rightarrow 10$
 $k = -15 \rightarrow 15$
 $l = -26 \rightarrow 19$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.107$
 $S = 1.10$
2479 reflections
175 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0598P)^2 + 1.3319P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.004$
 $\Delta\rho_{\max} = 0.84 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.57 \text{ e } \text{\AA}^{-3}$

Extinction correction: *SHELXTL* (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0120 (8)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	-0.00393 (3)	0.34535 (2)	0.307507 (13)	0.03274 (18)
O1	0.0807 (2)	0.03567 (15)	0.10091 (9)	0.0353 (5)
O2	0.1222 (2)	0.43270 (15)	-0.05895 (8)	0.0322 (5)
C1	0.3438 (3)	0.0117 (2)	0.18087 (13)	0.0354 (7)
H1A	0.4351	0.0104	0.2009	0.053*
H1B	0.3562	0.0299	0.1384	0.053*
H1C	0.3005	-0.0555	0.1836	0.053*
C2	0.2512 (3)	0.08893 (19)	0.21113 (13)	0.0259 (6)
C3	0.2670 (3)	0.1078 (2)	0.27228 (13)	0.0286 (6)
H3	0.3328	0.0691	0.2946	0.034*
C4	0.1895 (3)	0.1816 (2)	0.30181 (12)	0.0303 (7)
H4	0.1999	0.1921	0.3439	0.036*
C5	0.0968 (3)	0.2395 (2)	0.26867 (11)	0.0269 (6)
C6	0.0771 (3)	0.2229 (2)	0.20780 (12)	0.0264 (6)
H6	0.0129	0.2633	0.1857	0.032*
C7	0.1528 (3)	0.1459 (2)	0.17905 (12)	0.0255 (6)
C8	0.1169 (3)	0.1222 (2)	0.11452 (12)	0.0267 (6)
C9	0.1212 (3)	0.2048 (2)	0.06972 (11)	0.0246 (6)
C10	0.2006 (3)	0.2915 (2)	0.07938 (11)	0.0256 (6)
H10	0.2518	0.2979	0.1158	0.031*
C11	0.2069 (3)	0.3687 (2)	0.03726 (11)	0.0250 (6)
H11	0.2634	0.4268	0.0442	0.030*
C12	0.1289 (3)	0.3597 (2)	-0.01543 (12)	0.0267 (6)
C13	0.0496 (3)	0.2727 (2)	-0.02606 (12)	0.0308 (7)
H13	-0.0032	0.2668	-0.0621	0.037*
C14	0.0480 (3)	0.1956 (2)	0.01559 (12)	0.0294 (6)
H14	-0.0034	0.1356	0.0075	0.035*
C15	0.2006 (3)	0.5248 (2)	-0.04965 (13)	0.0334 (7)
H15A	0.3020	0.5102	-0.0497	0.040*
H15B	0.1756	0.5557	-0.0105	0.040*
C16	0.1645 (3)	0.5955 (2)	-0.10032 (13)	0.0364 (7)
H16A	0.1906	0.5644	-0.1388	0.055*
H16B	0.2154	0.6594	-0.0952	0.055*

H16C	0.0638	0.6090	-0.1000	0.055*
------	--------	--------	---------	--------

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0346 (3)	0.0327 (3)	0.0309 (3)	0.00282 (11)	0.01170 (11)	-0.00058 (11)
O1	0.0412 (13)	0.0268 (11)	0.0379 (11)	-0.0026 (9)	-0.0050 (9)	-0.0040 (8)
O2	0.0373 (12)	0.0317 (12)	0.0276 (10)	-0.0050 (9)	-0.0020 (8)	0.0032 (8)
C1	0.0350 (17)	0.0322 (18)	0.0390 (16)	0.0046 (14)	-0.0043 (13)	-0.0038 (13)
C2	0.0237 (15)	0.0226 (15)	0.0314 (15)	-0.0026 (12)	-0.0001 (11)	0.0005 (11)
C3	0.0281 (16)	0.0264 (16)	0.0314 (15)	0.0010 (12)	-0.0050 (11)	0.0036 (11)
C4	0.0317 (16)	0.0343 (16)	0.0248 (14)	-0.0063 (13)	0.0020 (11)	0.0019 (11)
C5	0.0265 (15)	0.0269 (15)	0.0273 (14)	-0.0050 (12)	0.0061 (11)	0.0013 (11)
C6	0.0209 (14)	0.0275 (15)	0.0307 (14)	0.0008 (11)	0.0022 (11)	0.0026 (11)
C7	0.0225 (15)	0.0246 (16)	0.0293 (14)	-0.0032 (11)	-0.0001 (11)	-0.0010 (10)
C8	0.0180 (14)	0.0304 (16)	0.0318 (15)	0.0022 (11)	0.0000 (10)	-0.0044 (13)
C9	0.0207 (14)	0.0278 (15)	0.0254 (14)	0.0031 (11)	-0.0018 (10)	-0.0037 (11)
C10	0.0197 (14)	0.0326 (16)	0.0245 (13)	0.0032 (12)	-0.0018 (10)	-0.0054 (11)
C11	0.0199 (14)	0.0273 (14)	0.0278 (14)	-0.0022 (11)	0.0023 (10)	-0.0060 (11)
C12	0.0245 (15)	0.0302 (16)	0.0254 (14)	0.0011 (12)	0.0029 (11)	0.0004 (11)
C13	0.0298 (16)	0.0361 (18)	0.0266 (15)	-0.0041 (13)	-0.0052 (12)	-0.0055 (12)
C14	0.0253 (15)	0.0309 (16)	0.0321 (16)	-0.0037 (13)	-0.0017 (12)	-0.0033 (12)
C15	0.0311 (16)	0.0324 (17)	0.0367 (16)	-0.0045 (13)	0.0001 (12)	0.0007 (12)
C16	0.0335 (17)	0.0361 (18)	0.0397 (16)	-0.0011 (13)	0.0051 (13)	0.0063 (13)

Geometric parameters (\AA , ^\circ)

Br1—C5	1.903 (3)	C8—C9	1.475 (4)
O1—C8	1.230 (3)	C9—C10	1.389 (4)
O2—C12	1.365 (3)	C9—C14	1.397 (4)
O2—C15	1.442 (3)	C10—C11	1.385 (4)
C1—C2	1.509 (4)	C10—H10	0.9500
C1—H1A	0.9800	C11—C12	1.393 (4)
C1—H1B	0.9800	C11—H11	0.9500
C1—H1C	0.9800	C12—C13	1.396 (4)
C2—C3	1.389 (4)	C13—C14	1.374 (4)
C2—C7	1.400 (4)	C13—H13	0.9500
C3—C4	1.388 (4)	C14—H14	0.9500
C3—H3	0.9500	C15—C16	1.502 (4)
C4—C5	1.382 (4)	C15—H15A	0.9900
C4—H4	0.9500	C15—H15B	0.9900
C5—C6	1.382 (4)	C16—H16A	0.9800
C6—C7	1.401 (4)	C16—H16B	0.9800
C6—H6	0.9500	C16—H16C	0.9800
C7—C8	1.506 (4)		
C12—O2—C15	117.9 (2)	C10—C9—C8	121.2 (2)
C2—C1—H1A	109.5	C14—C9—C8	120.2 (2)

C2—C1—H1B	109.5	C11—C10—C9	121.7 (2)
H1A—C1—H1B	109.5	C11—C10—H10	119.2
C2—C1—H1C	109.5	C9—C10—H10	119.2
H1A—C1—H1C	109.5	C10—C11—C12	118.8 (3)
H1B—C1—H1C	109.5	C10—C11—H11	120.6
C3—C2—C7	118.3 (2)	C12—C11—H11	120.6
C3—C2—C1	119.5 (2)	O2—C12—C11	124.0 (2)
C7—C2—C1	122.1 (3)	O2—C12—C13	115.7 (2)
C2—C3—C4	122.0 (3)	C11—C12—C13	120.2 (2)
C2—C3—H3	119.0	C14—C13—C12	120.0 (3)
C4—C3—H3	119.0	C14—C13—H13	120.0
C5—C4—C3	118.6 (2)	C12—C13—H13	120.0
C5—C4—H4	120.7	C13—C14—C9	120.6 (3)
C3—C4—H4	120.7	C13—C14—H14	119.7
C6—C5—C4	121.4 (3)	C9—C14—H14	119.7
C6—C5—Br1	119.3 (2)	O2—C15—C16	107.2 (2)
C4—C5—Br1	119.3 (2)	O2—C15—H15A	110.3
C5—C6—C7	119.3 (3)	C16—C15—H15A	110.3
C5—C6—H6	120.4	O2—C15—H15B	110.3
C7—C6—H6	120.4	C16—C15—H15B	110.3
C2—C7—C6	120.4 (3)	H15A—C15—H15B	108.5
C2—C7—C8	121.7 (2)	C15—C16—H16A	109.5
C6—C7—C8	117.8 (2)	C15—C16—H16B	109.5
O1—C8—C9	121.8 (2)	H16A—C16—H16B	109.5
O1—C8—C7	119.4 (2)	C15—C16—H16C	109.5
C9—C8—C7	118.8 (2)	H16A—C16—H16C	109.5
C10—C9—C14	118.6 (3)	H16B—C16—H16C	109.5
C7—C2—C3—C4	-0.7 (4)	O1—C8—C9—C10	-159.8 (3)
C1—C2—C3—C4	176.7 (3)	C7—C8—C9—C10	22.6 (4)
C2—C3—C4—C5	-1.8 (4)	O1—C8—C9—C14	18.3 (4)
C3—C4—C5—C6	2.2 (4)	C7—C8—C9—C14	-159.3 (3)
C3—C4—C5—Br1	-177.0 (2)	C14—C9—C10—C11	0.9 (4)
C4—C5—C6—C7	-0.1 (4)	C8—C9—C10—C11	179.0 (2)
Br1—C5—C6—C7	179.1 (2)	C9—C10—C11—C12	1.5 (4)
C3—C2—C7—C6	2.8 (4)	C15—O2—C12—C11	-0.1 (4)
C1—C2—C7—C6	-174.6 (3)	C15—O2—C12—C13	179.1 (2)
C3—C2—C7—C8	-172.7 (2)	C10—C11—C12—O2	177.1 (2)
C1—C2—C7—C8	10.0 (4)	C10—C11—C12—C13	-2.0 (4)
C5—C6—C7—C2	-2.4 (4)	O2—C12—C13—C14	-179.0 (3)
C5—C6—C7—C8	173.2 (2)	C11—C12—C13—C14	0.2 (4)
C2—C7—C8—O1	53.5 (4)	C12—C13—C14—C9	2.2 (4)
C6—C7—C8—O1	-122.1 (3)	C10—C9—C14—C13	-2.8 (4)
C2—C7—C8—C9	-128.8 (3)	C8—C9—C14—C13	179.1 (3)
C6—C7—C8—C9	55.6 (3)	C12—O2—C15—C16	-174.5 (2)

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
C11—H11···O1 ⁱ	0.95	2.42	3.313 (3)	156

Symmetry code: (i) $-x+1/2, y+1/2, z$.