

## 5,8-Dibromo-15,18-dimethoxy-2,11-dithia[3.3]paracyclophane

Guojun Jin\* and Yinghui Lu

Key Laboratory of Pesticides and Chemical Biology of the Ministry of Education, College of Chemistry, Central China Normal University, Wuhan 430079, People's Republic of China

Correspondence e-mail: jgj6100449@163.com


Received 1 July 2010; accepted 21 July 2010

Key indicators: single-crystal X-ray study;  $T = 298\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$ ;  $R$  factor = 0.059;  $wR$  factor = 0.151; data-to-parameter ratio = 20.4.

In the title compound [systematic name: 1<sup>2</sup>,1<sup>5</sup>-dibromo-5<sup>2</sup>,5<sup>5</sup>-dimethoxy-2,7-dithia-1,5(1,4)-dibenzeno-octaphane],  $\text{C}_{18}\text{H}_{18}\text{Br}_2\text{O}_2\text{S}_2$ , the dihedral angle between the aromatic rings is 0.6 (2)° and their centroid separation is 3.251 (2) Å, indicating that a trans-annular  $\pi-\pi$  interaction occurs. The dimethoxy and dibromo substituents are located at crossed positions because of the electronic and the steric nature of the substituents.

## Related literature

For the preparation of the title compound, see: Kay & Baek (1997); Xu *et al.* (2008). For paracyclophane and its derivatives, see: Clément *et al.* (2009); Wang *et al.* (2006); Yamamoto *et al.* (1997). For studies on the benzene dimer of [2.2]paracyclophane, see: Ball *et al.* (2004); Dahmen & Bräse (2002); Rowlands (2008); Valentini *et al.* (2008). For studies of [3.3]paracyclophane, see: Wang *et al.* (2004).



## Experimental

## Crystal data

$\text{C}_{18}\text{H}_{18}\text{Br}_2\text{O}_2\text{S}_2$	$V = 1843.2 (3)\text{ \AA}^3$
$M_r = 490.26$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 8.9576 (8)\text{ \AA}$	$\mu = 4.63\text{ mm}^{-1}$
$b = 16.2291 (14)\text{ \AA}$	$T = 298\text{ K}$
$c = 13.0251 (11)\text{ \AA}$	$0.16 \times 0.12 \times 0.10\text{ mm}$
$\beta = 103.240 (1)$	

## Data collection

Bruker SMART CCD area-detector diffractometer	4475 independent reflections
12800 measured reflections	2812 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.126$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$	219 parameters
$wR(F^2) = 0.151$	H-atom parameters constrained
$S = 0.95$	$\Delta\rho_{\text{max}} = 1.00\text{ e \AA}^{-3}$
4475 reflections	$\Delta\rho_{\text{min}} = -1.27\text{ e \AA}^{-3}$

Data collection: SMART (Bruker, 1997); cell refinement: SAINT (Bruker, 1999); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

The authors thank Professor Sheng-Hua Liu for technical assistance with the structure analysis and Dr Xiang-Gao Meng for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SI2274).

## References

- Ball, P. J., Shtoyko, T. R., Bauer, J. A. K., Oldham, W. J. & Connick, W. B. (2004). *Inorg. Chem.* **43**, 622–632.
- Bruker (1997). SMART. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (1999). SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Clément, S., Guyard, L., Knorr, M., Däschlein, C. & Strohmann, C. (2009). *Acta Cryst. E65*, o528.
- Dahmen, S. & Bräse, S. (2002). *J. Am. Chem. Soc.* **124**, 5940–5941.
- Kay, K. Y. & Baek, Y. G. (1997). *Chem. Ber. Rec.* **130**, 581–584.
- Rowlands, G. J. (2008). *Org. Biomol. Chem.* **6**, 1527–1534.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Valentini, L., Marrocchi, A., Seri, M., Mengoni, F., Meloni, F., Taticchi, A. & Kenny, J. M. (2008). *Thin Solid Films.* **516**, 7193–7198.
- Wang, W., Xu, J., Lai, Y. H. & Wang, F. (2004). *Macromolecules*, **37**, 3546–3553.
- Wang, W., Xu, J., Sun, Z., Zhang, X., Lu, Y. & Lai, Y. H. (2006). *Macromolecules*, **39**, 7277–7285.

# supporting information

*Acta Cryst.* (2010). E66, o2144 [https://doi.org/10.1107/S1600536810029053]

## 5,8-Dibromo-15,18-dimethoxy-2,11-dithia[3.3]paracyclophane

Guojun Jin and Yinghui Lu

### S1. Comment

Various studies on the benzene dimer of [2.2]paracyclophane have focused on the face-to-face stacking (Rowlands, 2008), it is known to play a significant role in chiral catalysis (Dahmen & Bräse, 2002), molecular electronics (Ball *et al.*, 2004), and organic solar cells (Valentini *et al.*, 2008). However, the [3.3]paracyclophane have received less attention (Wang *et al.* 2004). In our research, we have synthesized a series of novel dithia[3.3]paracyclophane. The inter plane distance of the two benzene rings of 3.251 Å is less than the normal packing distance of aromatic rings in organic aromatic molecules (3.4°), thus suggesting probable transannular  $\pi$ - $\pi$  interaction.


For the preparation of the title compound, see: Kay & Baek (1997); Xu *et al.* (2008); for the paracyclophanes and its derivatives, see: Clément *et al.* (2009); Wang *et al.* (2006); Yamamoto *et al.* (1997).

### S2. Experimental

A solution with equimolar amounts of 2,5-dibromo-1,4-bis(mercaptomethyl)benzene (3.26 g, 10 mmol) and 1,4-dibromo-methyl-2,5-dimethoxybenzene (3.22 g, 10 mmol) in degassed THF (500 mL) was added dropwise under N<sub>2</sub> over 12 h to a refluxing solution of potassium carbonate (6.9 g, 50 mmol) in EtOH (1.5 L). After an additional 2 h at the reflux temperature (353 K), the mixture was cooled and the solvent were removed. The resulting residue was treated with CH<sub>2</sub>Cl<sub>2</sub> (500 mL) and water (500 mL). The organic phase was separated, the aqueous extracted with CH<sub>2</sub>Cl<sub>2</sub> three times. The combined organic layers was dried over Na<sub>2</sub>SO<sub>4</sub>, then the solvent was removed, and the resulting solid was chromatographed on silica gel using CH<sub>2</sub>Cl<sub>2</sub>/petroleum ether (1:1, v/v) as eluent. Colourless single crystals of the title compound suitable for X-ray diffraction were obtained by slow evaporation of a dichloromethane/n-hexane (1:30) solution over a period of 5 days.

### S3. Refinement

Hydrogen atoms were placed in calculated positions and refined using a riding model with C—H = 0.93 - 0.97 Å and U<sub>iso</sub>(H) = 1.2U<sub>eq</sub>(C-H, CH<sub>2</sub>), 1.5U<sub>eq</sub>(CH<sub>3</sub>).

**Figure 1**

Molecular structure of the title compound with displacement ellipsoids drawn at the 30% probability level.

### $1^2,1^5$ -dibromo- $5^2,5^5$ -dimethoxy-2,7-dithia-1,5(1,4)-dibenzenaoctaphane

#### Crystal data



$M_r = 490.26$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 8.9576 (8)$  Å

$b = 16.2291 (14)$  Å

$c = 13.0251 (11)$  Å

$\beta = 103.240 (1)^\circ$

$V = 1843.2 (3)$  Å<sup>3</sup>

$Z = 4$

$F(000) = 976$

$D_x = 1.767 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 3885 reflections

$\theta = 2.5\text{--}26.9^\circ$

$\mu = 4.63 \text{ mm}^{-1}$

$T = 298$  K

Block, colorless

$0.16 \times 0.12 \times 0.10$  mm

#### Data collection

Bruker SMART CCD area detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and  $\omega$  scans

12800 measured reflections

4475 independent reflections

2812 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.126$

$\theta_{\text{max}} = 28.3^\circ, \theta_{\text{min}} = 2.0^\circ$

$h = -11 \rightarrow 11$

$k = -19 \rightarrow 21$

$l = -15 \rightarrow 17$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.059$$

$$wR(F^2) = 0.151$$

$$S = 0.95$$

4475 reflections

219 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0753P)^2]$$
$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 1.00 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -1.27 \text{ e \AA}^{-3}$$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.56512 (6)	0.00221 (3)	0.68306 (4)	0.0672 (2)
Br2	1.04343 (6)	0.17844 (3)	0.44267 (4)	0.05796 (19)
C1	0.9651 (5)	0.0721 (2)	0.6001 (3)	0.0383 (9)
C2	0.8556 (5)	0.0352 (2)	0.6428 (3)	0.0414 (9)
H2	0.8858	-0.0067	0.6919	0.050*
C3	0.7037 (5)	0.0569 (2)	0.6166 (3)	0.0421 (9)
C4	0.6510 (5)	0.1210 (2)	0.5448 (3)	0.0426 (9)
C5	0.7568 (5)	0.1535 (2)	0.4937 (3)	0.0419 (9)
H5	0.7248	0.1924	0.4408	0.050*
C6	0.9099 (5)	0.1296 (2)	0.5193 (3)	0.0386 (9)
C7	1.1351 (5)	0.0552 (3)	0.6408 (3)	0.0478 (10)
H7A	1.1921	0.1026	0.6250	0.057*
H7B	1.1630	0.0085	0.6027	0.057*
C8	1.2053 (5)	0.1368 (3)	0.8357 (3)	0.0603 (13)
H8A	1.2330	0.1323	0.9120	0.072*
H8B	1.2865	0.1669	0.8140	0.072*
C9	1.0584 (5)	0.1853 (3)	0.8039 (3)	0.0483 (11)
C10	0.9312 (5)	0.1636 (3)	0.8413 (3)	0.0466 (10)
C11	0.7901 (5)	0.1997 (3)	0.7979 (3)	0.0461 (10)
H11	0.7041	0.1842	0.8220	0.055*
C12	0.7759 (5)	0.2590 (3)	0.7186 (3)	0.0423 (9)
C13	0.9054 (5)	0.2850 (3)	0.6891 (3)	0.0447 (10)
C14	1.0460 (5)	0.2476 (3)	0.7292 (3)	0.0480 (10)
H14	1.1321	0.2642	0.7061	0.058*
C15	0.6193 (5)	0.2917 (3)	0.6623 (4)	0.0519 (11)

H15A	0.6279	0.3141	0.5948	0.062*
H15B	0.5928	0.3368	0.7035	0.062*
C16	0.4926 (5)	0.1575 (3)	0.5276 (4)	0.0531 (11)
H16A	0.4179	0.1132	0.5143	0.064*
H16B	0.4745	0.1922	0.4654	0.064*
C17	0.8336 (7)	0.0803 (4)	0.9594 (4)	0.0767 (16)
H17A	0.7924	0.1276	0.9874	0.115*
H17B	0.8688	0.0411	1.0149	0.115*
H17C	0.7553	0.0555	0.9054	0.115*
C18	1.0126 (6)	0.3817 (3)	0.5888 (4)	0.0703 (14)
H18A	1.0703	0.4095	0.6503	0.106*
H18B	0.9807	0.4208	0.5328	0.106*
H18C	1.0756	0.3402	0.5673	0.106*
O1	0.9536 (4)	0.1039 (2)	0.9173 (2)	0.0666 (9)
O2	0.8860 (4)	0.34533 (19)	0.6119 (2)	0.0584 (8)
S1	1.19297 (14)	0.03433 (8)	0.77929 (9)	0.0601 (3)
S2	0.46307 (13)	0.21831 (8)	0.63885 (10)	0.0559 (3)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.0566 (3)	0.0782 (4)	0.0670 (4)	-0.0235 (2)	0.0142 (3)	0.0087 (2)
Br2	0.0558 (3)	0.0679 (3)	0.0542 (3)	-0.0029 (2)	0.0209 (2)	0.0073 (2)
C1	0.038 (2)	0.041 (2)	0.0332 (19)	0.0023 (16)	0.0015 (15)	-0.0064 (16)
C2	0.046 (2)	0.039 (2)	0.036 (2)	-0.0006 (18)	0.0028 (17)	0.0016 (17)
C3	0.043 (2)	0.046 (2)	0.036 (2)	-0.0103 (18)	0.0058 (17)	-0.0035 (17)
C4	0.039 (2)	0.048 (2)	0.035 (2)	-0.0046 (18)	-0.0025 (16)	-0.0061 (17)
C5	0.045 (2)	0.050 (2)	0.0287 (19)	-0.0017 (18)	0.0036 (16)	0.0002 (17)
C6	0.041 (2)	0.045 (2)	0.0292 (18)	-0.0044 (17)	0.0055 (16)	-0.0023 (16)
C7	0.043 (2)	0.059 (3)	0.040 (2)	0.005 (2)	0.0048 (18)	-0.0066 (19)
C8	0.041 (3)	0.097 (4)	0.036 (2)	0.002 (2)	-0.0063 (18)	-0.010 (2)
C9	0.036 (2)	0.072 (3)	0.032 (2)	-0.005 (2)	-0.0033 (17)	-0.018 (2)
C10	0.049 (3)	0.064 (3)	0.0253 (19)	0.000 (2)	0.0041 (17)	-0.0072 (18)
C11	0.037 (2)	0.064 (3)	0.039 (2)	-0.0033 (19)	0.0114 (17)	-0.0076 (19)
C12	0.036 (2)	0.051 (2)	0.038 (2)	-0.0044 (18)	0.0059 (16)	-0.0099 (18)
C13	0.043 (2)	0.054 (2)	0.034 (2)	-0.0112 (19)	0.0024 (17)	-0.0136 (18)
C14	0.037 (2)	0.068 (3)	0.037 (2)	-0.011 (2)	0.0034 (16)	-0.018 (2)
C15	0.040 (2)	0.059 (3)	0.056 (3)	-0.003 (2)	0.010 (2)	0.002 (2)
C16	0.032 (2)	0.072 (3)	0.050 (2)	-0.003 (2)	-0.0010 (18)	-0.001 (2)
C17	0.077 (4)	0.100 (4)	0.054 (3)	0.002 (3)	0.017 (3)	0.027 (3)
C18	0.068 (3)	0.069 (3)	0.072 (3)	-0.023 (3)	0.014 (3)	-0.004 (3)
O1	0.055 (2)	0.102 (3)	0.0419 (17)	0.0117 (19)	0.0104 (15)	0.0129 (17)
O2	0.056 (2)	0.0589 (18)	0.058 (2)	-0.0108 (16)	0.0077 (15)	0.0047 (15)
S1	0.0459 (7)	0.0797 (8)	0.0499 (7)	0.0193 (6)	0.0010 (5)	0.0161 (6)
S2	0.0312 (6)	0.0714 (8)	0.0654 (8)	-0.0008 (5)	0.0113 (5)	-0.0028 (6)

Geometric parameters ( $\text{\AA}$ ,  $\text{^{\circ}}$ )

Br1—C3	1.889 (4)	C10—C11	1.390 (6)
Br2—C6	1.898 (4)	C11—C12	1.397 (6)
C1—C2	1.372 (6)	C11—H11	0.9300
C1—C6	1.408 (5)	C12—C13	1.369 (6)
C1—C7	1.518 (6)	C12—C15	1.521 (6)
C2—C3	1.371 (6)	C13—O2	1.386 (5)
C2—H2	0.9300	C13—C14	1.388 (6)
C3—C4	1.407 (5)	C14—H14	0.9300
C4—C5	1.382 (6)	C15—S2	1.809 (4)
C4—C16	1.506 (6)	C15—H15A	0.9700
C5—C6	1.390 (6)	C15—H15B	0.9700
C5—H5	0.9300	C16—S2	1.822 (5)
C7—S1	1.792 (4)	C16—H16A	0.9700
C7—H7A	0.9700	C16—H16B	0.9700
C7—H7B	0.9700	C17—O1	1.369 (6)
C8—C9	1.508 (6)	C17—H17A	0.9600
C8—S1	1.812 (5)	C17—H17B	0.9600
C8—H8A	0.9700	C17—H17C	0.9600
C8—H8B	0.9700	C18—O2	1.372 (6)
C9—C10	1.384 (6)	C18—H18A	0.9600
C9—C14	1.389 (6)	C18—H18B	0.9600
C10—O1	1.368 (5)	C18—H18C	0.9600
C2—C1—C6	115.5 (4)	C12—C11—H11	119.6
C2—C1—C7	122.2 (4)	C13—C12—C11	118.7 (4)
C6—C1—C7	122.2 (4)	C13—C12—C15	120.3 (4)
C3—C2—C1	123.3 (4)	C11—C12—C15	120.9 (4)
C3—C2—H2	118.4	C12—C13—O2	116.6 (4)
C1—C2—H2	118.4	C12—C13—C14	120.9 (4)
C2—C3—C4	121.2 (4)	O2—C13—C14	122.3 (4)
C2—C3—Br1	119.0 (3)	C13—C14—C9	120.1 (4)
C4—C3—Br1	119.8 (3)	C13—C14—H14	119.9
C5—C4—C3	116.1 (4)	C9—C14—H14	119.9
C5—C4—C16	120.4 (4)	C12—C15—S2	116.4 (3)
C3—C4—C16	123.4 (4)	C12—C15—H15A	108.2
C4—C5—C6	121.7 (4)	S2—C15—H15A	108.2
C4—C5—H5	119.1	C12—C15—H15B	108.2
C6—C5—H5	119.1	S2—C15—H15B	108.2
C5—C6—C1	121.5 (4)	H15A—C15—H15B	107.3
C5—C6—Br2	117.6 (3)	C4—C16—S2	113.5 (3)
C1—C6—Br2	120.9 (3)	C4—C16—H16A	108.9
C1—C7—S1	114.9 (3)	S2—C16—H16A	108.9
C1—C7—H7A	108.5	C4—C16—H16B	108.9
S1—C7—H7A	108.5	S2—C16—H16B	108.9
C1—C7—H7B	108.5	H16A—C16—H16B	107.7
S1—C7—H7B	108.5	O1—C17—H17A	109.5

H7A—C7—H7B	107.5	O1—C17—H17B	109.5
C9—C8—S1	113.5 (3)	H17A—C17—H17B	109.5
C9—C8—H8A	108.9	O1—C17—H17C	109.5
S1—C8—H8A	108.9	H17A—C17—H17C	109.5
C9—C8—H8B	108.9	H17B—C17—H17C	109.5
S1—C8—H8B	108.9	O2—C18—H18A	109.5
H8A—C8—H8B	107.7	O2—C18—H18B	109.5
C10—C9—C14	119.4 (4)	H18A—C18—H18B	109.5
C10—C9—C8	120.5 (4)	O2—C18—H18C	109.5
C14—C9—C8	119.8 (4)	H18A—C18—H18C	109.5
O1—C10—C9	116.0 (4)	H18B—C18—H18C	109.5
O1—C10—C11	124.2 (4)	C10—O1—C17	119.3 (4)
C9—C10—C11	119.7 (4)	C18—O2—C13	119.4 (4)
C10—C11—C12	120.7 (4)	C7—S1—C8	102.2 (2)
C10—C11—H11	119.6	C15—S2—C16	104.1 (2)
C6—C1—C2—C3	-5.6 (6)	O1—C10—C11—C12	-178.4 (4)
C7—C1—C2—C3	171.5 (4)	C9—C10—C11—C12	-1.5 (6)
C1—C2—C3—C4	-1.6 (6)	C10—C11—C12—C13	-4.2 (6)
C1—C2—C3—Br1	-179.0 (3)	C10—C11—C12—C15	173.1 (4)
C2—C3—C4—C5	7.4 (6)	C11—C12—C13—O2	-178.4 (4)
Br1—C3—C4—C5	-175.2 (3)	C15—C12—C13—O2	4.2 (5)
C2—C3—C4—C16	-168.8 (4)	C11—C12—C13—C14	6.4 (6)
Br1—C3—C4—C16	8.6 (5)	C15—C12—C13—C14	-171.0 (4)
C3—C4—C5—C6	-6.0 (6)	C12—C13—C14—C9	-2.8 (6)
C16—C4—C5—C6	170.4 (4)	O2—C13—C14—C9	-177.7 (3)
C4—C5—C6—C1	-1.2 (6)	C10—C9—C14—C13	-3.0 (6)
C4—C5—C6—Br2	179.7 (3)	C8—C9—C14—C13	171.4 (4)
C2—C1—C6—C5	7.0 (5)	C13—C12—C15—S2	141.2 (3)
C7—C1—C6—C5	-170.2 (4)	C11—C12—C15—S2	-36.0 (5)
C2—C1—C6—Br2	-174.0 (3)	C5—C4—C16—S2	-105.9 (4)
C7—C1—C6—Br2	8.8 (5)	C3—C4—C16—S2	70.2 (5)
C2—C1—C7—S1	-32.5 (5)	C9—C10—O1—C17	178.9 (4)
C6—C1—C7—S1	144.4 (3)	C11—C10—O1—C17	-4.1 (7)
S1—C8—C9—C10	69.9 (5)	C12—C13—O2—C18	171.4 (4)
S1—C8—C9—C14	-104.4 (4)	C14—C13—O2—C18	-13.5 (6)
C14—C9—C10—O1	-177.8 (4)	C1—C7—S1—C8	-80.3 (4)
C8—C9—C10—O1	7.9 (6)	C9—C8—S1—C7	56.6 (4)
C14—C9—C10—C11	5.1 (6)	C12—C15—S2—C16	-76.7 (4)
C8—C9—C10—C11	-169.2 (4)	C4—C16—S2—C15	53.5 (4)