

Poly[μ_{10} -[(phosphonomethyl)imino-dimethylene]diphosphonato]-dithallium(I)]

Khodayar Gholivand* and Ali Reza Farrokhi*

Department of Chemistry, Tarbiat Modares University, PO Box 14115-175 Tehran, Iran
Correspondence e-mail: gholi_kh@modares.ac.ir, arfmhf@yahoo.com


Received 9 July 2009; accepted 4 August 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{N}-\text{C}) = 0.008$ Å; R factor = 0.025; wR factor = 0.055; data-to-parameter ratio = 16.8.

The title compound, $[\text{Tl}_2(\text{C}_3\text{H}_{10}\text{NO}_9\text{P}_3)]_n$, a Tl^{I} organic-inorganic hybrid complex, was synthesized by the reaction of nitrilotris(methylenephosphonic acid) with thallium(I) nitrate. There are two types of Tl^{I} ions in the complex, with coordination numbers of eight and seven and with stereochemically active and inactive lone-pair electrons, respectively. In the crystal, the doubly deprotonated ligands form two-dimensional hydrogen-bonded layers through $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds. The NH group is involved in a trifurcated intramolecular hydrogen bond. Coordination of the phosphonate ligands to the Tl^{I} ions creates a three-dimensional structure.

Related literature

For related metal phosphonate complexes of the same ligand, see: Sharma *et al.* (2001).



Experimental

Crystal data

$[\text{Tl}_2(\text{C}_3\text{H}_{10}\text{NO}_9\text{P}_3)]$	$\gamma = 68.085 (1)^{\circ}$
$M_r = 705.77$	$V = 635.06 (8)$ Å 3
Triclinic, $\overline{P}\bar{1}$	$Z = 2$
$a = 7.9236 (6)$ Å	Mo $K\alpha$ radiation
$b = 8.0932 (6)$ Å	$\mu = 25.76$ mm $^{-1}$
$c = 10.9136 (8)$ Å	$T = 100$ K
$\alpha = 81.422 (1)^{\circ}$	$0.16 \times 0.14 \times 0.10$ mm
$\beta = 79.023 (1)^{\circ}$	

Data collection

Bruker APEXII CCD area-detector diffractometer	6627 measured reflections
Absorption correction: numerical (<i>XPREP</i> ; Bruker, 2007)	2744 independent reflections
$T_{\min} = 0.104$, $T_{\max} = 0.183$	2437 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.025$	12 restraints
$wR(F^2) = 0.055$	H-atom parameters constrained
$S = 1.05$	$\Delta\rho_{\text{max}} = 1.72$ e Å $^{-3}$
2744 reflections	$\Delta\rho_{\text{min}} = -1.83$ e Å $^{-3}$
163 parameters	

Table 1
Hydrogen-bond geometry (Å, °).

D-H···A	D-H	H···A	D···A	D-H···A
N1-H1N···O3	0.87	2.39	2.882 (5)	116
N1-H1N···O5	0.87	2.53	2.957 (6)	111
N1-H1N···O8	0.87	2.19	2.837 (7)	131
O1-H1O···O1 ⁱ	0.82	1.78	2.504 (8)	147
O2-H2O···O2 ⁱⁱ	0.82	1.68	2.497 (7)	171
O6-H6O···O9 ⁱⁱⁱ	0.82	1.67	2.484 (6)	171
O7-H7O···O4 ^{iv}	0.82	1.70	2.521 (6)	180

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 2, -y, -z + 1$; (iii) $x + 1, y - 1, z$; (iv) $-x + 1, -y, -z$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Support of this investigation by Tarbiat Modares University Research Council is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2128).

References

- Bruker (2007). *APEX2, SAINT* and *XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sharma, C. V. K., Clearfield, A., Cabeza, A., Aranda, M. A. G. & Bruque, S. (2001). *J. Am. Chem. Soc.* **123**, 2885–2886.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supporting information

Acta Cryst. (2010). E66, m873 [https://doi.org/10.1107/S1600536809031006]

Poly[$\{\mu_{10}$ -[(phosphonomethyl)iminodimethylene]diphosphonato}dithallium(I)]

Khodayar Gholivand and Ali Reza Farrokhi

S1. Comment

Metal phosphonate complexes of the ligand nitrilotris(methylenephosphonic acid (H_6L)), giving two different types of 1:1 (M/L) metal phosphonate complexes, $M[NH(CH_2PO_3H)_3(H_2O)_3](M = Mn, Co, Ni, Cu, Zn, Cd)$, with two-dimensional hydrogen-bonded layered structures, have been prepared previously by (Sharma *et al.*, 2001).

In the structure of the title compound, synthesized by the reaction of the same ligand with thallium(I) nitrate, the nitrilotris(methylenephosphonate) (H_4L^{2-}) group is doubly deprotonated and two different environments for the Thallium atoms are observed, as shown in Figs. 1 and 2. Eight O-atoms of five H_4L^{2-} phosphonate ligands are coordinated to the Tl_1 ion (Fig. 1), while seven oxygen atoms of five H_4L^{2-} phosphonate ligands are coordinated to the Tl_2 ion (Fig. 2).

The conformation of the doubly deprotonated nitrilotris(methylenephosphonate) ligand is illustrated in Fig. 3. The hydrogen atoms on atoms O1 and O2 are positionally disordered with relative occupancies of 0.5:0.5. Each H_4L^{2-} dianion links to five Tl_1 and five Tl_2 ions (Fig. 4).


The doubly deprotonated H_4L^{2-} ligand forms two-dimensional hydrogen bonded layers *via* O—H \cdots O hydrogen bonds involving hydroxyl groups O1, O2, O6 and O7 (Table 1 and Fig. 5). Atoms O3, O5 and O8 do not contribute in this type of hydrogen bond, rather they coordinate only to the Tl^+ ions. The NH group is involved in a 4-centre trifurcated intramolecular hydrogen bond with O-atoms O3, O5 and O8 (Table 1). Coordination of the ligand to the metal ions in the interlayer space creates a three-dimensional structure (Fig. 6).

S2. Experimental


Thallium(I)nitrate (0.133 g, 0.5 mmol) was added in several portions to a solution of nitrilotris(methylenephosphonicacid) (H_6L) (0.104 g, 0.35 mmol) in 12 ml of a deionized water-ethanol mixture (3:5). The solution was stirred for seven days and a white precipitate was obtained. This was filtered off and recrystallized from deionized water at rt. Colorless prism-like crystals of the title compound were obtained in 68% yield (based on the Tl atom). Elemental analysis for $Tl_2(H_4L)$, $C_3H_{10}NO_9P_3Tl_2:C$, 5.06; H, 1.36; N, 2.03%. Calc.: C, 5.10; H, 1.41; N, 1.98%.

S3. Refinement


The NH and OH H-atoms were located in a difference electron-density map and were refined with distance restraints: O—H = 0.82 (2) and N—H = 0.86 (2) Å, with $U_{iso}(H) = 1.2U_{eq}(N,O)$. The C-bound H atoms were positioned geometrically and refined using a riding model: C—H = 0.99 Å, with $U_{iso}(H) = 1.2$ times $U_{eq}(C)$.

**Figure 1**


Environment of the Tl1 atom.

**Figure 2**


Environment of the Tl2 atom.

**Figure 3**


The conformation of the doubly deprotonated nitrilotris(methylenephosphonate) ligand.

**Figure 4**


Each H_4L^{2-} dianion links to five Tl^{+1} and five Tl^{+2} ions.

**Figure 5**

Two-dimensional hydrogen-bonded layers formed *via* O—H \cdots O hydrogen bonds involving hydroxyl groups.

**Figure 6**

The three-dimensional structure resulting from the coordination of the ligand to the metal ions in the interlayer space.

**Figure 7**

Reaction scheme.

Poly[μ -[(phosphonomethyl)iminodimethylene]diphosphonato]dithallium(I)]*Crystal data*

$M_r = 705.77$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.9236(6)$ Å

$b = 8.0932(6)$ Å

$c = 10.9136(8)$ Å

$\alpha = 81.422(1)^\circ$

$\beta = 79.023(1)^\circ$

$\gamma = 68.085(1)^\circ$

$V = 635.06(8)$ Å³

$Z = 2$

$F(000) = 628$

$D_x = 3.691$ Mg m⁻³

Mo K α radiation, $\lambda = 0.71073$ Å

Cell parameters from 266 reflections

$\theta = 3\text{--}26^\circ$

$\mu = 25.76$ mm⁻¹

$T = 100$ K

Prism, colorless

$0.16 \times 0.14 \times 0.10$ mm

*Data collection*Bruker APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: numerical
(XPREP; Bruker, 2007)

$T_{\min} = 0.104$, $T_{\max} = 0.183$

6627 measured reflections

2744 independent reflections

2437 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\max} = 27.0^\circ$, $\theta_{\min} = 1.9^\circ$

$h = -10 \rightarrow 10$

$k = -10 \rightarrow 10$

$l = -13 \rightarrow 13$

*Refinement*Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.025$

$wR(F^2) = 0.055$

$S = 1.05$

2744 reflections

163 parameters

12 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0225P)^2 + 1.104P]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.003$

$\Delta\rho_{\max} = 1.72$ e Å⁻³

$\Delta\rho_{\min} = -1.83$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Tl1	0.29076 (3)	-0.05875 (3)	0.25484 (2)	0.01064 (7)	
Tl2	0.23248 (3)	0.24843 (3)	0.54672 (2)	0.01703 (8)	
P1	0.7521 (3)	0.2462 (3)	0.42352 (16)	0.0201 (4)	
P2	0.7554 (2)	-0.0669 (2)	0.10302 (14)	0.0090 (3)	
P3	0.2557 (2)	0.4329 (2)	0.20298 (14)	0.0091 (3)	
O1	0.6684 (8)	0.4387 (7)	0.4597 (5)	0.0331 (13)	
H1O	0.5753	0.4894	0.5072	0.040*	0.50
O2	0.9424 (7)	0.1622 (8)	0.4631 (5)	0.0311 (12)	
H2O	0.9828	0.0586	0.4936	0.037*	0.50
O3	0.6291 (6)	0.1391 (5)	0.4655 (4)	0.0119 (9)	
O4	0.7469 (6)	-0.1263 (6)	-0.0199 (4)	0.0148 (9)	
O5	0.6365 (6)	-0.1126 (6)	0.2173 (4)	0.0151 (9)	
O6	0.9564 (6)	-0.1194 (5)	0.1274 (4)	0.0129 (9)	
H6O	1.0047	-0.2157	0.1660	0.015*	
O7	0.2360 (6)	0.4303 (6)	0.0624 (4)	0.0145 (9)	
H7O	0.2418	0.3313	0.0485	0.017*	
O8	0.2641 (6)	0.2594 (5)	0.2794 (4)	0.0108 (8)	
O9	0.1203 (6)	0.6039 (6)	0.2530 (4)	0.0141 (9)	
N1	0.6274 (7)	0.2583 (6)	0.2032 (4)	0.0086 (10)	
H1N	0.5590	0.2094	0.2560	0.010*	
C1	0.7944 (8)	0.2553 (8)	0.2523 (6)	0.0116 (12)	
H1A	0.8225	0.3638	0.2177	0.014*	
H1B	0.9018	0.1497	0.2251	0.014*	
C2	0.6768 (8)	0.1763 (8)	0.0791 (5)	0.0089 (11)	
H2A	0.7749	0.2130	0.0257	0.011*	
H2B	0.5677	0.2209	0.0351	0.011*	
C3	0.4836 (8)	0.4433 (8)	0.1986 (6)	0.0111 (12)	
H3A	0.5140	0.5133	0.1209	0.013*	
H3B	0.4828	0.5047	0.2708	0.013*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Tl1	0.00946 (12)	0.01138 (12)	0.01117 (12)	-0.00463 (9)	-0.00049 (9)	-0.00006 (9)
Tl2	0.01643 (14)	0.01862 (14)	0.00967 (13)	-0.00101 (10)	0.00066 (10)	0.00088 (10)
P1	0.0291 (10)	0.0340 (10)	0.0090 (8)	-0.0260 (9)	-0.0018 (7)	0.0005 (7)

P2	0.0088 (7)	0.0092 (7)	0.0072 (7)	-0.0023 (6)	0.0003 (6)	0.0001 (6)
P3	0.0081 (7)	0.0095 (7)	0.0080 (7)	-0.0021 (6)	0.0001 (6)	0.0003 (6)
O1	0.048 (3)	0.031 (2)	0.028 (2)	-0.026 (2)	0.006 (2)	-0.0084 (19)
O2	0.023 (2)	0.051 (3)	0.027 (2)	-0.0233 (19)	-0.0095 (18)	0.0074 (19)
O3	0.011 (2)	0.012 (2)	0.013 (2)	-0.0062 (17)	-0.0001 (18)	0.0018 (17)
O4	0.023 (2)	0.010 (2)	0.011 (2)	-0.0034 (18)	-0.0045 (19)	-0.0010 (17)
O5	0.010 (2)	0.020 (2)	0.011 (2)	-0.0038 (18)	0.0011 (18)	0.0028 (18)
O6	0.010 (2)	0.010 (2)	0.015 (2)	-0.0028 (17)	-0.0031 (18)	0.0090 (17)
O7	0.020 (2)	0.012 (2)	0.011 (2)	-0.0045 (19)	-0.0062 (19)	0.0013 (17)
O8	0.010 (2)	0.009 (2)	0.012 (2)	-0.0033 (17)	0.0013 (17)	-0.0001 (16)
O9	0.011 (2)	0.013 (2)	0.014 (2)	0.0010 (18)	-0.0017 (18)	-0.0027 (18)
N1	0.009 (2)	0.011 (2)	0.007 (2)	-0.005 (2)	0.0031 (19)	-0.0031 (19)
C1	0.009 (3)	0.016 (3)	0.014 (3)	-0.007 (2)	-0.001 (2)	-0.004 (2)
C2	0.006 (3)	0.011 (3)	0.007 (3)	-0.001 (2)	0.002 (2)	-0.001 (2)
C3	0.010 (3)	0.012 (3)	0.009 (3)	-0.003 (2)	0.004 (2)	-0.006 (2)

Geometric parameters (\AA , $^\circ$)

Tl1—O8	2.555 (4)	P3—C3	1.830 (6)
Tl1—O5	2.569 (4)	O1—Tl2 ⁱⁱⁱ	2.906 (5)
Tl1—O4 ⁱ	2.778 (4)	O1—H1O	0.8200
Tl1—O3 ⁱⁱ	3.159 (4)	O2—Tl2 ^{vi}	2.965 (5)
Tl2—O5 ⁱⁱ	2.870 (4)	O2—H2O	0.8201
Tl2—O8	2.871 (4)	O3—Tl2 ⁱⁱ	2.926 (4)
Tl2—O1 ⁱⁱⁱ	2.906 (5)	O3—Tl1 ⁱⁱ	3.159 (4)
Tl2—O3	2.922 (4)	O4—Tl1 ⁱ	2.778 (4)
Tl2—O3 ⁱⁱ	2.926 (4)	O5—Tl2 ⁱⁱ	2.870 (4)
Tl2—O2 ^{iv}	2.965 (5)	O6—H6O	0.8200
Tl2—O9 ^v	3.159 (4)	O7—H7O	0.8201
P1—O3	1.501 (4)	O9—Tl2 ^v	3.159 (4)
P1—O2	1.522 (5)	N1—C3	1.506 (7)
P1—O1	1.529 (6)	N1—C1	1.510 (7)
P1—C1	1.829 (6)	N1—C2	1.517 (7)
P2—O5	1.502 (5)	N1—H1N	0.8699
P2—O4	1.512 (4)	C1—H1A	0.9900
P2—O6	1.552 (4)	C1—H1B	0.9900
P2—C2	1.823 (6)	C2—H2A	0.9900
P3—O9	1.503 (4)	C2—H2B	0.9900
P3—O8	1.507 (4)	C3—H3A	0.9900
P3—O7	1.576 (4)	C3—H3B	0.9900
O8—Tl1—O5	82.65 (14)	O9—P3—O7	109.6 (2)
O8—Tl1—O4 ⁱ	73.72 (13)	O8—P3—O7	112.4 (2)
O5—Tl1—O4 ⁱ	90.64 (13)	O9—P3—C3	105.9 (3)
O8—Tl1—O3 ⁱⁱ	84.83 (12)	O8—P3—C3	104.2 (3)
O5—Tl1—O3 ⁱⁱ	80.14 (12)	O7—P3—C3	105.6 (3)
O4 ⁱ —Tl1—O3 ⁱⁱ	157.62 (11)	P1—O1—Tl2 ⁱⁱⁱ	140.5 (3)
O8—Tl1—Tl2	42.43 (9)	P1—O1—H1O	131.2

O5—Tl1—Tl2	88.38 (10)	Tl2 ⁱⁱⁱ —O1—H1O	87.0
O4 ⁱ —Tl1—Tl2	115.70 (8)	P1—O2—Tl2 ^{vi}	142.9 (3)
O3 ⁱⁱ —Tl1—Tl2	44.31 (7)	P1—O2—H2O	122.8
O5 ⁱⁱ —Tl2—O8	153.26 (12)	Tl2 ^{vi} —O2—H2O	91.4
O5 ⁱⁱ —Tl2—O1 ⁱⁱⁱ	90.93 (14)	P1—O3—Tl2	131.5 (2)
O8—Tl2—O1 ⁱⁱⁱ	94.48 (13)	P1—O3—Tl2 ⁱⁱ	121.5 (2)
O5 ⁱⁱ —Tl2—O3	79.76 (12)	Tl2—O3—Tl2 ⁱⁱ	106.77 (13)
O8—Tl2—O3	76.90 (12)	P1—O3—Tl1 ⁱⁱ	95.14 (19)
O1 ⁱⁱⁱ —Tl2—O3	72.87 (14)	Tl2—O3—Tl1 ⁱⁱ	91.85 (11)
O5 ⁱⁱ —Tl2—O3 ⁱⁱ	76.88 (12)	Tl2 ⁱⁱ —O3—Tl1 ⁱⁱ	86.74 (11)
O8—Tl2—O3 ⁱⁱ	84.01 (11)	P2—O4—Tl1 ⁱ	130.9 (2)
O1 ⁱⁱⁱ —Tl2—O3 ⁱⁱ	145.49 (14)	P2—O5—Tl1	130.2 (2)
O3—Tl2—O3 ⁱⁱ	73.23 (13)	P2—O5—Tl2 ⁱⁱ	122.6 (2)
O5 ⁱⁱ —Tl2—O2 ^{iv}	123.69 (14)	Tl1—O5—Tl2 ⁱⁱ	106.85 (15)
O8—Tl2—O2 ^{iv}	66.41 (13)	P2—O6—H6O	119.1
O1 ⁱⁱⁱ —Tl2—O2 ^{iv}	137.28 (16)	P3—O7—H7O	111.0
O3—Tl2—O2 ^{iv}	132.04 (14)	P3—O8—Tl1	141.1 (2)
O3 ⁱⁱ —Tl2—O2 ^{iv}	73.22 (14)	P3—O8—Tl2	118.2 (2)
O5 ⁱⁱ —Tl2—O9 ^v	76.06 (12)	Tl1—O8—Tl2	100.67 (13)
O8—Tl2—O9 ^v	129.64 (11)	P3—O9—Tl2 ^v	141.9 (3)
O1 ⁱⁱⁱ —Tl2—O9 ^v	92.73 (14)	C3—N1—C1	111.4 (4)
O3—Tl2—O9 ^v	151.59 (11)	C3—N1—C2	112.6 (4)
O3 ⁱⁱ —Tl2—O9 ^v	114.72 (11)	C1—N1—C2	112.4 (4)
O2 ^{iv} —Tl2—O9 ^v	74.91 (13)	C3—N1—H1N	95.5
O5 ⁱⁱ —Tl2—Tl1	125.55 (9)	C1—N1—H1N	113.9
O8—Tl2—Tl1	36.90 (8)	C2—N1—H1N	109.9
O1 ⁱⁱⁱ —Tl2—Tl1	129.09 (10)	N1—C1—P1	110.2 (4)
O3—Tl2—Tl1	79.78 (8)	N1—C1—H1A	109.6
O3 ⁱⁱ —Tl2—Tl1	48.95 (8)	P1—C1—H1A	109.6
O2 ^{iv} —Tl2—Tl1	52.33 (11)	N1—C1—H1B	109.6
O9 ^v —Tl2—Tl1	126.82 (8)	P1—C1—H1B	109.6
O3—P1—O2	115.2 (3)	H1A—C1—H1B	108.1
O3—P1—O1	114.5 (3)	N1—C2—P2	110.9 (4)
O2—P1—O1	108.2 (3)	N1—C2—H2A	109.5
O3—P1—C1	106.0 (3)	P2—C2—H2A	109.5
O2—P1—C1	104.8 (3)	N1—C2—H2B	109.5
O1—P1—C1	107.4 (3)	P2—C2—H2B	109.5
O5—P2—O4	117.4 (3)	H2A—C2—H2B	108.1
O5—P2—O6	111.2 (2)	N1—C3—P3	110.7 (4)
O4—P2—O6	112.0 (3)	N1—C3—H3A	109.5
O5—P2—C2	106.8 (3)	P3—C3—H3A	109.5
O4—P2—C2	104.8 (3)	N1—C3—H3B	109.5
O6—P2—C2	103.2 (3)	P3—C3—H3B	109.5
O9—P3—O8	118.1 (3)	H3A—C3—H3B	108.1
O8—Tl1—Tl2—O5 ⁱⁱ	-151.22 (18)	O5—P2—O4—Tl1 ⁱ	132.5 (3)
O5—Tl1—Tl2—O5 ⁱⁱ	-70.36 (17)	O6—P2—O4—Tl1 ⁱ	-97.0 (3)
O4 ⁱ —Tl1—Tl2—O5 ⁱⁱ	-160.29 (15)	C2—P2—O4—Tl1 ⁱ	14.2 (4)

O3 ⁱⁱ —Tl1—Tl2—O5 ⁱⁱ	7.15 (15)	O4—P2—O5—Tl1	−51.5 (4)
O5—Tl1—Tl2—O8	80.86 (17)	O6—P2—O5—Tl1	177.6 (3)
O4 ⁱ —Tl1—Tl2—O8	−9.07 (17)	C2—P2—O5—Tl1	65.7 (4)
O3 ⁱⁱ —Tl1—Tl2—O8	158.37 (18)	O4—P2—O5—Tl2 ⁱⁱ	136.2 (3)
O8—Tl1—Tl2—O1 ⁱⁱⁱ	−23.9 (2)	O6—P2—O5—Tl2 ⁱⁱ	5.3 (3)
O5—Tl1—Tl2—O1 ⁱⁱⁱ	56.95 (18)	C2—P2—O5—Tl2 ⁱⁱ	−106.6 (3)
O4 ⁱ —Tl1—Tl2—O1 ⁱⁱⁱ	−32.99 (19)	O8—Tl1—O5—P2	−77.8 (3)
O3 ⁱⁱ —Tl1—Tl2—O1 ⁱⁱⁱ	134.46 (19)	O4 ⁱ —Tl1—O5—P2	−4.3 (3)
O8—Tl1—Tl2—O3	−81.77 (16)	O3 ⁱⁱ —Tl1—O5—P2	−163.8 (3)
O5—Tl1—Tl2—O3	−0.90 (12)	Tl2—Tl1—O5—P2	−120.0 (3)
O4 ⁱ —Tl1—Tl2—O3	−90.84 (13)	O8—Tl1—O5—Tl2 ⁱⁱ	95.39 (16)
O3 ⁱⁱ —Tl1—Tl2—O3	76.60 (15)	O4 ⁱ —Tl1—O5—Tl2 ⁱⁱ	168.88 (15)
O8—Tl1—Tl2—O3 ⁱⁱ	−158.37 (18)	O3 ⁱⁱ —Tl1—O5—Tl2 ⁱⁱ	9.39 (13)
O5—Tl1—Tl2—O3 ⁱⁱ	−77.51 (15)	Tl2—Tl1—O5—Tl2 ⁱⁱ	53.19 (12)
O4 ⁱ —Tl1—Tl2—O3 ⁱⁱ	−167.44 (15)	O9—P3—O8—Tl1	137.8 (3)
O8—Tl1—Tl2—O2 ^{iv}	100.75 (19)	O7—P3—O8—Tl1	8.7 (5)
O5—Tl1—Tl2—O2 ^{iv}	−178.39 (16)	C3—P3—O8—Tl1	−105.1 (4)
O4 ⁱ —Tl1—Tl2—O2 ^{iv}	91.67 (17)	O9—P3—O8—Tl2	−39.7 (3)
O3 ⁱⁱ —Tl1—Tl2—O2 ^{iv}	−100.88 (17)	O7—P3—O8—Tl2	−168.8 (2)
O8—Tl1—Tl2—O9 ^v	109.29 (17)	C3—P3—O8—Tl2	77.3 (3)
O5—Tl1—Tl2—O9 ^v	−169.85 (14)	O5—Tl1—O8—P3	86.5 (4)
O4 ⁱ —Tl1—Tl2—O9 ^v	100.22 (15)	O4 ⁱ —Tl1—O8—P3	−6.3 (4)
O3 ⁱⁱ —Tl1—Tl2—O9 ^v	−92.34 (15)	O3 ⁱⁱ —Tl1—O8—P3	167.2 (4)
O3—P1—O1—Tl2 ⁱⁱⁱ	173.5 (4)	Tl2—Tl1—O8—P3	−177.8 (5)
O2—P1—O1—Tl2 ⁱⁱⁱ	43.6 (5)	O5—Tl1—O8—Tl2	−95.68 (14)
C1—P1—O1—Tl2 ⁱⁱⁱ	−69.0 (5)	O4 ⁱ —Tl1—O8—Tl2	171.49 (16)
O3—P1—O2—Tl2 ^{vi}	−143.4 (4)	O3 ⁱⁱ —Tl1—O8—Tl2	−14.98 (12)
O1—P1—O2—Tl2 ^{vi}	−13.9 (6)	O5 ⁱⁱ —Tl2—O8—P3	−121.0 (3)
C1—P1—O2—Tl2 ^{vi}	100.4 (5)	O1 ⁱⁱⁱ —Tl2—O8—P3	−20.0 (3)
O2—P1—O3—Tl2	144.1 (3)	O3—Tl2—O8—P3	−91.2 (2)
O1—P1—O3—Tl2	17.7 (4)	O3 ⁱⁱ —Tl2—O8—P3	−165.3 (3)
C1—P1—O3—Tl2	−100.5 (3)	O2 ^{iv} —Tl2—O8—P3	120.4 (3)
O2—P1—O3—Tl2 ⁱⁱ	−42.3 (4)	O9 ^v —Tl2—O8—P3	77.3 (3)
O1—P1—O3—Tl2 ⁱⁱ	−168.6 (3)	Tl1—Tl2—O8—P3	178.4 (3)
C1—P1—O3—Tl2 ⁱⁱ	73.1 (3)	O5 ⁱⁱ —Tl2—O8—Tl1	60.5 (3)
O2—P1—O3—Tl1 ⁱⁱ	47.0 (3)	O1 ⁱⁱⁱ —Tl2—O8—Tl1	161.60 (16)
O1—P1—O3—Tl1 ⁱⁱ	−79.3 (3)	O3—Tl2—O8—Tl1	90.36 (14)
C1—P1—O3—Tl1 ⁱⁱ	162.5 (2)	O3 ⁱⁱ —Tl2—O8—Tl1	16.23 (13)
O5 ⁱⁱ —Tl2—O3—P1	−106.4 (3)	O2 ^{iv} —Tl2—O8—Tl1	−58.05 (16)
O8—Tl2—O3—P1	86.7 (3)	O9 ^v —Tl2—O8—Tl1	−101.10 (16)
O1 ⁱⁱⁱ —Tl2—O3—P1	−12.2 (3)	O8—P3—O9—Tl2 ^v	−15.8 (5)
O3 ⁱⁱ —Tl2—O3—P1	174.3 (4)	O7—P3—O9—Tl2 ^v	114.7 (4)
O2 ^{iv} —Tl2—O3—P1	127.0 (3)	C3—P3—O9—Tl2 ^v	−131.9 (4)
O9 ^v —Tl2—O3—P1	−74.5 (4)	C3—N1—C1—P1	80.8 (5)
Tl1—Tl2—O3—P1	124.3 (3)	C2—N1—C1—P1	−151.7 (4)
O5 ⁱⁱ —Tl2—O3—Tl2 ⁱⁱ	79.25 (14)	O3—P1—C1—N1	33.2 (5)
O8—Tl2—O3—Tl2 ⁱⁱ	−87.61 (13)	O2—P1—C1—N1	155.5 (4)
O1 ⁱⁱⁱ —Tl2—O3—Tl2 ⁱⁱ	173.44 (17)	O1—P1—C1—N1	−89.7 (5)

O3 ⁱⁱ —Tl2—O3—Tl2 ⁱⁱ	0.0	C3—N1—C2—P2	-152.4 (4)
O2 ^{iv} —Tl2—O3—Tl2 ⁱⁱ	-47.3 (2)	C1—N1—C2—P2	80.8 (5)
O9 ^v —Tl2—O3—Tl2 ⁱⁱ	111.2 (2)	O5—P2—C2—N1	39.1 (4)
Tl1—Tl2—O3—Tl2 ⁱⁱ	-50.01 (10)	O4—P2—C2—N1	164.3 (4)
O5 ⁱⁱ —Tl2—O3—Tl1 ⁱⁱ	-7.90 (11)	O6—P2—C2—N1	-78.3 (4)
O8—Tl2—O3—Tl1 ⁱⁱ	-174.76 (12)	C1—N1—C3—P3	-156.6 (4)
O1 ⁱⁱⁱ —Tl2—O3—Tl1 ⁱⁱ	86.28 (14)	C2—N1—C3—P3	76.0 (5)
O3 ⁱⁱ —Tl2—O3—Tl1 ⁱⁱ	-87.15 (13)	O9—P3—C3—N1	156.2 (4)
O2 ^{iv} —Tl2—O3—Tl1 ⁱⁱ	-134.49 (15)	O8—P3—C3—N1	31.0 (4)
O9 ^v —Tl2—O3—Tl1 ⁱⁱ	24.0 (3)	O7—P3—C3—N1	-87.6 (4)
Tl1—Tl2—O3—Tl1 ⁱⁱ	-137.17 (9)		

Symmetry codes: (i) $-x+1, -y, -z$; (ii) $-x+1, -y, -z+1$; (iii) $-x+1, -y+1, -z+1$; (iv) $x-1, y, z$; (v) $-x, -y+1, -z+1$; (vi) $x+1, y, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1N···O3	0.87	2.39	2.882 (5)	116
N1—H1N···O5	0.87	2.53	2.957 (6)	111
N1—H1N···O8	0.87	2.19	2.837 (7)	131
O1—H1O···O1 ⁱⁱⁱ	0.82	1.78	2.504 (8)	147
O2—H2O···O2 ^{vii}	0.82	1.68	2.497 (7)	171
O6—H6O···O9 ^{viii}	0.82	1.67	2.484 (6)	171
O7—H7O···O4 ⁱ	0.82	1.70	2.521 (6)	180

Symmetry codes: (i) $-x+1, -y, -z$; (iii) $-x+1, -y+1, -z+1$; (vii) $-x+2, -y, -z+1$; (viii) $x+1, y-1, z$.