

# Opipramol dipicrate

Jerry P. Jasinski,<sup>a\*</sup> Albert E. Pek,<sup>a</sup> B. P. Siddaraju,<sup>b</sup>  
H. S. Yathirajan<sup>c</sup> and B. Narayana<sup>d</sup>

<sup>a</sup>Department of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, <sup>b</sup>Department of Chemistry, V. V. Puram College of Science, Bangalore 560 004, India, <sup>c</sup>Department of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, and <sup>d</sup>Department of Studies in Chemistry, Mangalore University, Mangalagangothri 574 199, India  
Correspondence e-mail: jjasinski@keene.edu


Received 17 June 2010; accepted 5 July 2010

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.049;  $wR$  factor = 0.123; data-to-parameter ratio = 16.0.

In the crystal structure of the title compound,  $\text{C}_{23}\text{H}_{31}\text{N}_3\text{O}^{2+} \cdot 2\text{C}_6\text{H}_2\text{N}_3\text{O}_7^-$ , {systematic name: 1-[3-(5*H*-dibenz[*b,f*]azepin-5-yl)propyl]-4-(2-hydroxyethyl)piperazine-1,4-dium bis(2,4,6-trinitrophenolate)} the piperazine group in the opipramol dication is protonated at both N atoms. Each picrate anion interacts with the protonated N atom in the cation through a bifurcated  $\text{N}-\text{H} \cdots \text{O}$  hydrogen bond, forming an  $R_2^1(6)$  ring motif. In the cation, the dihedral angle between the mean planes of the two benzene rings is  $50.81(8)^\circ$ . Intermolecular  $\text{O}-\text{H} \cdots \text{O}$  and weak  $\text{C}-\text{H} \cdots \text{O}$  hydrogen bonds, and weak  $\pi$ -ring and  $\pi$ - $\pi$  stacking interactions dominate the crystal packing.

## Related literature

For the use of opipramol in the treatment of anxiety disorder, see: Moller *et al.* (2001). For its use in the preparation of amine derivatives, see: Shriner *et al.* (1980). For crystal engineering research, see: Desiraju *et al.* (1989). For related structures, see: Bindya *et al.* (2007); Jasinski *et al.* (2010); Yathirajan *et al.* (2007). For bond-length data, see: Allen *et al.* (1987).



## Experimental

### Crystal data

$\text{C}_{23}\text{H}_{31}\text{N}_3\text{O}^{2+} \cdot 2\text{C}_6\text{H}_2\text{N}_3\text{O}_7^-$   
 $M_r = 821.72$   
Triclinic,  $P\bar{1}$   
 $a = 7.3838(8)$  Å  
 $b = 12.0400(13)$  Å  
 $c = 22.074(2)$  Å  
 $\alpha = 74.821(1)^\circ$   
 $\beta = 84.355(2)^\circ$   
 $\gamma = 73.866(2)^\circ$   
 $V = 1818.6(3)$  Å<sup>3</sup>  
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.12$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.55 \times 0.50 \times 0.14$  mm

### Data collection

Bruker APEXII CCD diffractometer  
Absorption correction: multi-scan (SADABS; Bruker, 2008)  
 $T_{\min} = 0.937$ ,  $T_{\max} = 0.983$   
10692 measured reflections  
10692 independent reflections  
7831 reflections with  $I > 2\sigma(I)$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.123$   
 $S = 0.98$   
10692 reflections  
669 parameters  
H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.64$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.39$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N1}-\text{H1} \cdots \text{O1B}^i$	0.91 (2)	1.85 (2)	2.6901 (16)	152.6 (18)
$\text{N1}-\text{H1} \cdots \text{O7B}^i$	0.91 (2)	2.383 (19)	3.0466 (17)	130.0 (16)
$\text{N2}-\text{H2} \cdots \text{O1A}^{ii}$	0.90 (2)	1.78 (2)	2.6204 (16)	154.6 (19)
$\text{N2}-\text{H2} \cdots \text{O2A}^{ii}$	0.90 (2)	2.43 (2)	3.0711 (16)	128.2 (16)
$\text{O1}-\text{H1C} \cdots \text{O1B}^i$	0.82	2.50	3.1600 (19)	138
$\text{O1}-\text{H1C} \cdots \text{O7B}^i$	0.82	2.38	3.0841 (18)	144

Symmetry codes: (i)  $x + 1, y, z$ ; (ii)  $x, y + 1, z$ .

Table 2

$Y-X \cdots C_g$   $\pi$  ring interactions (Å).

$C_g3$  and  $C_g9$  are the centroids of the C10–C15 and C1A–C6A rings, respectively.  $C_gX \cdots \text{Perp}$  and  $C_gY \cdots \text{Perp}$  are the perpendicular distances between atoms  $X$  and  $Y$  and the ring centroid.

$Y-X \cdots C_g$	$X \cdots C_g$	$Y \cdots C_g$	$X \cdots \text{Perp}$
$\text{C1A}-\text{O1A} \cdots C_g3^i$	3.5674 (13)	3.6471 (17)	3.494
$\text{N3A}-\text{O4A} \cdots C_g9$	3.8172 (17)	3.8173 (17)	-3.357
$\text{N3B}-\text{O4B} \cdots C_g9^{ii}$	3.4320 (15)	3.9391 (15)	3.288

Symmetry codes: (i)  $x, -1 + y, z$ ; (ii)  $1 - x, -y, 1 - z$ .

Table 3

$C_g \cdots C_g$   $\pi$  stacking interactions (Å).

$C_g2$ ,  $C_g3$ ,  $C_g8$  and  $C_g9$  are the centroids of the C10–C15, C18–C23, C1A–C6A and C1B–C6B rings, respectively.  $C_gX \cdots \text{Perp}$  and  $C_gY \cdots \text{Perp}$  are the perpendicular distances between the ring centroid and the other ring.

	$C_gX \cdots C_gY$	$C_gX \cdots \text{Perp}$	$C_gY \cdots \text{Perp}$
$C_g2 \cdots C_g2^i$	3.8038 (11)	-3.5589 (7)	-3.5590 (7)
$C_g3 \cdots C_g3^i$	3.7164 (10)	-3.6624 (7)	-3.6623 (7)
$C_g8 \cdots C_g9$	3.9558 (10)	-3.2475 (6)	3.3731 (6)

Symmetry code: (i)  $2 - x, 1 - y, -z$ .

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINTE* (Bruker, 2008); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

JPJ thanks Dr Matthias Zeller and the YSU Department of Chemistry for their assistance with the data collection. The diffractometer was funded by NSF grant 0087210, by Ohio Board of Regents grant CAP-491, and by YSU. HSY thanks the UOM for research facilities and for sabbatical leave. BPS thanks R. L. Fine Chem, Bangalore, for a gift sample of opipramol dihydrochloride.

---

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2786).

---

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bindya, S., Wong, W.-T., Ashok, M. A., Yathirajan, H. S. & Rathore, R. S. (2007). *Acta Cryst. C* **63**, o546–o548.
- Bruker (2008). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Desiraju, G. R. (1989). *Crystal Engineering: The Design of Organic Solids*. Amsterdam: Elsevier.
- Jasinski, J. P., Butcher, R. J., Hakim Al-Ariqie, O. N. M., Yathirajan, H. S. & Narayana, B. (2010). *Acta Cryst. E* **66**, o411–o412.
- Moller, H. J., Volz, H. P., Reimann, I. W. & Stoll, K. D. (2001). *J. Clin. Psychopharmacol.* **21**, 59–65.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shriner, R. L., Fuson, R. C., Curtin, D. Y. & Morrill, T. C. (1980). *Qualitative Identification of Organic Compounds*, 6th ed., pp. 236–237. New York: Wiley.
- Yathirajan, H. S., Ashok, M. A., Narayana Achar, B. & Bolte, M. (2007). *Acta Cryst. E* **63**, o1693–o1695.

## supporting information

*Acta Cryst.* (2010). E66, o1979–o1980 [https://doi.org/10.1107/S1600536810026565]

## Opipramol dipicrate

Jerry P. Jasinski, Albert E. Pek, B. P. Siddaraju, H. S. Yathirajan and B. Narayana

## S1. Comment

Opipramol (systematic IUPAC name: 4-[3-(5*H*-dibenz[b,f]azepin-5-yl)propyl]-1-piperazinethanol) is an antidepressant and anxiolytic typically used in the treatment of generalized anxiety disorder (Moller *et al.*, 2001). Opipramol, a drug widely prescribed in Germany, is a tricyclic compound with no reuptake-inhibiting properties. However, it has pronounced D2-, 5-HT2-, and H1-blocking potential and high affinity to sigma receptors (sigma-1 and sigma-2). Crystalline picrates have commonly been used in the preparation of amine derivatives in qualitative organic chemistry (Shriner *et al.*, 1980). Hydrogen bonding plays a key role in molecular recognition and crystal engineering research (Desiraju *et al.*, 1989). The crystal structures of trifluoperazinium dipicrate (Yathirajan *et al.*, 2007), amitriptylinium picrate (Bindya *et al.*, 2007) and imatinibium dipicrate (Jasinski *et al.*, 2010) have been reported. The present work reports the crystal structure of the salt formed by the interaction between 4-[3-(5*H*-dibenz[b,f]azepin-5-yl)propyl]-1-piperazinethanol dihydrochloride and 2,4,6-trinitrophenol in aqueous medium.

In opipramol dipicrate,  $C_{23}H_{33}N_3O^+$ ,  $(C_6H_2N_3O_7^-)_2$ , the piperazine group in the opipramol cation is protonated at both of the N atoms. The 6-membered piperazine group (N1/C5/C6/N2/C4/C3) adopts a slightly distorted chair conformation with puckering parameters  $Q$ ,  $\theta$  and  $\varphi$  of 0.584 (7) Å, 178.40°, and 312.658 (8)°, respectively (Fig. 1). For an ideal chair  $\theta$  has a value of 0 or 180°. Bond distances and angles are in normal ranges (Allen *et al.*, 1987).  $R_2^1(6)$  graph-set motifs are formed between piperazine N1—H1 and N2—H2 groups and the picrate anions (labeled A and B) through bifurcated N—H $\cdots$ O hydrogen bonds (Fig. 2). The mean plane of the two *o*-NO<sub>2</sub> groups in the two picrate anions are twisted by 31.8 (8)°, 31.8 (8)° in both the A ring B rings with respect to the mean planes of the 6-membered benzene rings. The *p*-NO<sub>2</sub> groups in both picrate anions are nearly in the plane of the ring (torsion angles O4A—N3A—C4A—C3A = -1.7 (2)°; O4B—N3B—C4B—C3B = -12.1 (2)°). An extensive array of weak O—H $\cdots$ O and C—H $\cdots$ O intermolecular hydrogen bonds (Table 1), weak  $\pi$ -ring (Table 2) and  $\pi$ - $\pi$  (Table 3) stacking interactions dominate crystal packing in the unit cell (Fig. 3).


## S2. Experimental

Opipramol dihydrochloride (4.38 g, 0.01 mol) was dissolved in 25 ml of water and picric acid (2.4 g, 0.01 mol) was dissolved in 25 ml of water. Both the solutions were mixed and stirred in a beaker at room temperature for one hour. The mixture was kept aside for two days at room temperature. The formed salt was filtered & dried in vacuum desiccator over phosphorous pentoxide. The salt was recrystallized from DMSO by slow evaporation (m.p: 453–455 K).

## S3. Refinement

The H1C, H1 and H2 atoms were located by a Fourier map. These H atoms and the rest of the H atoms were then positioned geometrically and allowed to ride on their parent atoms with Atom—H lengths of 0.82 Å (O1), 0.91 Å (NH), 0.93 Å (CH), 0.97 Å (CH<sub>2</sub>) or (CH<sub>3</sub>). Isotropic displacement parameters for these atoms were set to 1.40 times (OH), 1.20 times (NH), 1.20 (CH) or 1.22 (CH<sub>2</sub>) times (CH<sub>3</sub>)  $U_{eq}$  of the parent atom. The highest and lowest peaks (0.64 & 0.31 e<sup>Å</sup><sup>-3</sup>)

are located 1.21 Å and 0.31 Å from atoms N1A and H1C, respectively.



**Figure 1**

Molecular structure of,  $C_{23}H_{33}N_3O^+$ ,  $(C_6H_2N_3O_7)_2$ , showing the atom labeling scheme and 30% probability displacement ellipsoids.



**Figure 2**

Diagram for the  $R_2^1(6)\cdots ab..$ graph-set motif in the cation of the title compound,  $C_{23}H_{33}N_3O^+$ ,  $(C_6H_2N_3O_7)^-$ .

**Figure 3**

Packing diagram of the title compound viewed down the  $a$  axis. Dashed lines indicate intermolecular  $N-H\cdots O$  and  $C-H\cdots O$  hydrogen bond interactions.

1-[3-(5*H*-dibenz[*b,f*]azepin-5-yl)propyl]-4-(2-hydroxyethyl)piperazine-1,4-dium bis(2,4,6-trinitrophenolate)*Crystal data* $C_{23}H_{31}N_3O^{2+} \cdot 2C_6H_5N_3O_7^-$  $M_r = 821.72$ Triclinic,  $P\bar{1}$ 

Hall symbol: -p 1

 $a = 7.3838$  (8) Å $b = 12.0400$  (13) Å $c = 22.074$  (2) Å $\alpha = 74.821$  (1)° $\beta = 84.355$  (2)° $\gamma = 73.866$  (2)° $V = 1818.6$  (3) Å<sup>3</sup> $Z = 2$  $F(000) = 856$  $D_x = 1.501$  Mg m<sup>-3</sup>Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 5178 reflections

 $\theta = 2.9$ – $30.4$ ° $\mu = 0.12$  mm<sup>-1</sup> $T = 100$  K

Plate, yellow

 $0.55 \times 0.50 \times 0.14$  mm*Data collection*

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\omega$  scans

Absorption correction: multi-scan

(SADABS; Bruker, 2008)

 $T_{\min} = 0.937$ ,  $T_{\max} = 0.983$ 

10692 measured reflections

10692 independent reflections

7831 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.000$  $\theta_{\text{max}} = 31.1$ °,  $\theta_{\text{min}} = 1.8$ ° $h = -10 \rightarrow 10$  $k = -16 \rightarrow 17$  $l = 0 \rightarrow 30$ *Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.049$  $wR(F^2) = 0.123$  $S = 0.98$ 

10692 reflections

669 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0431P)^2 + 1.3196P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\text{max}} < 0.001$  $\Delta\rho_{\text{max}} = 0.64$  e Å<sup>-3</sup> $\Delta\rho_{\text{min}} = -0.39$  e Å<sup>-3</sup>*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	1.02455 (17)	0.60675 (11)	0.37838 (6)	0.0151 (2)
N1A	0.8758 (2)	0.06526 (12)	0.19787 (7)	0.0288 (3)
N1B	0.49985 (18)	0.35440 (11)	0.34309 (6)	0.0191 (2)

N2	0.83911 (17)	0.63062 (11)	0.26315 (5)	0.0154 (2)
N2A	0.36714 (18)	-0.11954 (11)	0.30334 (6)	0.0201 (3)
N2B	0.0123 (2)	0.16475 (12)	0.46259 (7)	0.0240 (3)
N3	0.65920 (18)	0.69770 (11)	0.08383 (6)	0.0183 (2)
N3A	0.2539 (2)	0.30858 (12)	0.23707 (7)	0.0296 (3)
N3B	0.6228 (2)	-0.07470 (12)	0.40742 (6)	0.0262 (3)
O1	1.32600 (16)	0.50804 (10)	0.47765 (5)	0.0238 (2)
H1C	1.3355	0.4665	0.4528	0.036*
O1A	0.73234 (15)	-0.13584 (9)	0.23643 (5)	0.0225 (2)
O1B	0.13483 (17)	0.36702 (10)	0.41214 (6)	0.0335 (3)
O2A	0.45142 (16)	-0.22069 (9)	0.29802 (6)	0.0243 (2)
O2B	-0.12159 (18)	0.22565 (14)	0.43026 (7)	0.0439 (4)
O3A	0.23060 (17)	-0.10226 (11)	0.33974 (6)	0.0309 (3)
O3B	-0.00152 (19)	0.10754 (10)	0.51708 (6)	0.0302 (3)
O4A	0.0937 (2)	0.31534 (12)	0.25984 (7)	0.0451 (4)
O4B	0.5803 (2)	-0.16385 (10)	0.44083 (6)	0.0363 (3)
O5A	0.3147 (2)	0.39677 (10)	0.21386 (6)	0.0366 (3)
O5B	0.76309 (19)	-0.07890 (11)	0.37238 (6)	0.0350 (3)
O6A	0.9082 (2)	0.14690 (14)	0.15701 (10)	0.0721 (6)
O6B	0.64141 (15)	0.33702 (10)	0.30921 (5)	0.0240 (2)
O7A	0.99987 (19)	-0.02019 (13)	0.22406 (7)	0.0419 (3)
O7B	0.40979 (18)	0.45584 (10)	0.34673 (6)	0.0327 (3)
C1	1.2578 (2)	0.63008 (14)	0.44618 (8)	0.0226 (3)
C1A	0.6250 (2)	-0.03839 (12)	0.24106 (7)	0.0169 (3)
C1B	0.2478 (2)	0.27017 (12)	0.40804 (7)	0.0188 (3)
C2	1.0532 (2)	0.65724 (14)	0.43134 (7)	0.0201 (3)
C2A	0.4357 (2)	-0.01908 (12)	0.26842 (7)	0.0174 (3)
C2B	0.2017 (2)	0.15826 (13)	0.43494 (7)	0.0180 (3)
C3	0.8236 (2)	0.60313 (13)	0.37804 (7)	0.0158 (3)
C3A	0.3154 (2)	0.09238 (13)	0.26613 (7)	0.0203 (3)
C3B	0.3198 (2)	0.04765 (13)	0.43780 (7)	0.0203 (3)
C4	0.7918 (2)	0.55480 (12)	0.32445 (6)	0.0157 (3)
C4A	0.3782 (2)	0.19229 (13)	0.23820 (7)	0.0223 (3)
C4B	0.4978 (2)	0.04110 (12)	0.40896 (7)	0.0196 (3)
C5	1.0750 (2)	0.68021 (13)	0.31640 (7)	0.0173 (3)
C5A	0.5629 (2)	0.18292 (13)	0.21535 (7)	0.0224 (3)
C5B	0.5537 (2)	0.14175 (13)	0.37891 (7)	0.0180 (3)
C6	1.0402 (2)	0.63363 (13)	0.26278 (7)	0.0181 (3)
C6A	0.6808 (2)	0.07208 (13)	0.21871 (7)	0.0202 (3)
C6B	0.4335 (2)	0.25296 (12)	0.37819 (7)	0.0170 (3)
C7	0.8042 (2)	0.58956 (14)	0.20786 (7)	0.0208 (3)
C8	0.5958 (2)	0.61594 (14)	0.19562 (7)	0.0199 (3)
C9	0.5705 (2)	0.61345 (14)	0.12799 (7)	0.0217 (3)
C10	0.7251 (2)	0.67135 (13)	0.02498 (7)	0.0196 (3)
C11	0.6288 (3)	0.61785 (14)	-0.00504 (7)	0.0252 (3)
C12	0.7035 (3)	0.58299 (15)	-0.05966 (8)	0.0309 (4)
C13	0.8755 (3)	0.60076 (16)	-0.08394 (8)	0.0327 (4)
C14	0.9697 (3)	0.65574 (15)	-0.05503 (8)	0.0296 (4)

C15	0.8954 (2)	0.69466 (14)	-0.00106 (7)	0.0229 (3)
C16	0.9953 (2)	0.75994 (16)	0.02502 (8)	0.0280 (3)
C17	0.9182 (3)	0.84618 (16)	0.05467 (8)	0.0284 (4)
C18	0.7192 (2)	0.89186 (14)	0.07048 (7)	0.0228 (3)
C19	0.6521 (3)	1.01141 (15)	0.07358 (8)	0.0312 (4)
C20	0.4661 (3)	1.05952 (16)	0.08865 (8)	0.0359 (4)
C21	0.3421 (3)	0.98862 (17)	0.10155 (8)	0.0345 (4)
C22	0.4037 (2)	0.87034 (16)	0.09885 (7)	0.0264 (3)
C23	0.5914 (2)	0.82097 (13)	0.08345 (7)	0.0198 (3)
H1	1.096 (3)	0.5306 (18)	0.3834 (9)	0.027 (5)*
H2	0.769 (3)	0.7069 (19)	0.2581 (9)	0.032 (5)*
H1A	1.334 (3)	0.6519 (17)	0.4086 (9)	0.029 (5)*
H1B	1.265 (3)	0.6790 (17)	0.4736 (9)	0.028 (5)*
H2A	1.000 (3)	0.7399 (18)	0.4188 (9)	0.028 (5)*
H2B	0.984 (2)	0.6202 (16)	0.4667 (9)	0.019 (4)*
H3A	0.748 (2)	0.6818 (15)	0.3746 (8)	0.014 (4)*
H3B	0.797 (2)	0.5526 (16)	0.4172 (8)	0.018 (4)*
H3C	0.192 (3)	0.0976 (17)	0.2851 (9)	0.027 (5)*
H3D	0.280 (3)	-0.0219 (17)	0.4569 (9)	0.025 (5)*
H4A	0.871 (2)	0.4727 (15)	0.3266 (8)	0.014 (4)*
H4B	0.660 (3)	0.5557 (15)	0.3244 (8)	0.018 (4)*
H5A	1.205 (3)	0.6769 (16)	0.3158 (8)	0.020 (4)*
H5B	0.995 (3)	0.7623 (16)	0.3130 (8)	0.022 (5)*
H5C	0.608 (3)	0.2514 (18)	0.1990 (9)	0.031 (5)*
H5D	0.670 (3)	0.1386 (17)	0.3592 (9)	0.029 (5)*
H6A	1.064 (3)	0.6881 (17)	0.2241 (9)	0.028 (5)*
H6B	1.119 (3)	0.5537 (17)	0.2650 (8)	0.023 (5)*
H7A	0.868 (3)	0.5037 (18)	0.2157 (9)	0.028 (5)*
H7B	0.865 (3)	0.6356 (16)	0.1722 (9)	0.021 (4)*
H8A	0.530 (2)	0.6955 (15)	0.2024 (8)	0.015 (4)*
H8B	0.537 (2)	0.5596 (16)	0.2237 (8)	0.019 (4)*
H9A	0.637 (3)	0.5335 (17)	0.1205 (8)	0.023 (5)*
H9B	0.430 (3)	0.6262 (17)	0.1206 (9)	0.029 (5)*
H11	0.513 (3)	0.6024 (18)	0.0116 (10)	0.033 (5)*
H12	0.632 (3)	0.5471 (19)	-0.0806 (10)	0.038 (6)*
H13	0.932 (3)	0.5741 (19)	-0.1207 (10)	0.038 (6)*
H14	1.085 (3)	0.6724 (17)	-0.0727 (9)	0.030 (5)*
H16	1.129 (3)	0.746 (2)	0.0137 (10)	0.044 (6)*
H17	0.998 (3)	0.8897 (18)	0.0633 (10)	0.036 (6)*
H19	0.740 (3)	1.0580 (18)	0.0661 (10)	0.034 (5)*
H20	0.426 (3)	1.1418 (19)	0.0917 (10)	0.040 (6)*
H21	0.213 (3)	1.023 (2)	0.1115 (11)	0.045 (6)*
H22	0.313 (3)	0.8197 (17)	0.1082 (9)	0.029 (5)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0133 (6)	0.0123 (5)	0.0194 (6)	-0.0018 (4)	-0.0006 (4)	-0.0052 (4)



N1A	0.0266 (7)	0.0211 (6)	0.0364 (8)	-0.0088 (6)	0.0042 (6)	-0.0022 (6)
N1B	0.0171 (6)	0.0173 (6)	0.0237 (6)	-0.0042 (5)	0.0011 (5)	-0.0073 (5)
N2	0.0164 (6)	0.0132 (5)	0.0158 (5)	-0.0026 (5)	-0.0005 (4)	-0.0032 (4)
N2A	0.0165 (6)	0.0203 (6)	0.0246 (6)	-0.0040 (5)	-0.0010 (5)	-0.0083 (5)
N2B	0.0241 (7)	0.0248 (7)	0.0272 (7)	-0.0115 (6)	0.0024 (5)	-0.0092 (5)
N3	0.0225 (6)	0.0187 (6)	0.0144 (5)	-0.0071 (5)	0.0001 (5)	-0.0036 (5)
N3A	0.0403 (9)	0.0184 (6)	0.0229 (7)	0.0090 (6)	-0.0089 (6)	-0.0073 (5)
N3B	0.0341 (8)	0.0167 (6)	0.0246 (7)	0.0057 (6)	-0.0132 (6)	-0.0082 (5)
O1	0.0223 (6)	0.0215 (5)	0.0244 (5)	0.0004 (4)	-0.0065 (4)	-0.0047 (4)
O1A	0.0192 (5)	0.0133 (5)	0.0306 (6)	-0.0006 (4)	0.0033 (4)	-0.0030 (4)
O1B	0.0281 (6)	0.0145 (5)	0.0456 (7)	0.0017 (5)	0.0160 (5)	-0.0005 (5)
O2A	0.0216 (6)	0.0166 (5)	0.0340 (6)	-0.0045 (4)	0.0040 (5)	-0.0071 (5)
O2B	0.0188 (6)	0.0655 (10)	0.0417 (8)	-0.0087 (6)	-0.0041 (5)	-0.0047 (7)
O3A	0.0267 (6)	0.0318 (6)	0.0347 (7)	-0.0081 (5)	0.0110 (5)	-0.0126 (5)
O3B	0.0439 (7)	0.0259 (6)	0.0256 (6)	-0.0174 (5)	0.0102 (5)	-0.0098 (5)
O4A	0.0434 (8)	0.0295 (7)	0.0431 (8)	0.0164 (6)	0.0061 (6)	-0.0060 (6)
O4B	0.0476 (8)	0.0139 (5)	0.0443 (8)	-0.0005 (5)	-0.0108 (6)	-0.0065 (5)
O5A	0.0537 (9)	0.0153 (5)	0.0377 (7)	0.0023 (5)	-0.0136 (6)	-0.0090 (5)
O5B	0.0374 (7)	0.0278 (6)	0.0285 (6)	0.0135 (5)	-0.0034 (5)	-0.0101 (5)
O6A	0.0530 (10)	0.0301 (8)	0.1039 (15)	-0.0083 (7)	0.0356 (10)	0.0170 (9)
O6B	0.0191 (5)	0.0256 (6)	0.0279 (6)	-0.0064 (4)	0.0065 (4)	-0.0094 (5)
O7A	0.0241 (7)	0.0411 (8)	0.0543 (9)	-0.0092 (6)	-0.0031 (6)	0.0005 (7)
O7B	0.0311 (7)	0.0154 (5)	0.0498 (8)	-0.0068 (5)	0.0170 (6)	-0.0109 (5)
C1	0.0212 (8)	0.0202 (7)	0.0256 (8)	-0.0051 (6)	-0.0080 (6)	-0.0018 (6)
C1A	0.0175 (7)	0.0139 (6)	0.0171 (6)	-0.0010 (5)	-0.0033 (5)	-0.0022 (5)
C1B	0.0182 (7)	0.0141 (6)	0.0218 (7)	-0.0019 (5)	-0.0004 (5)	-0.0031 (5)
C2	0.0185 (7)	0.0190 (7)	0.0239 (7)	0.0002 (6)	-0.0065 (6)	-0.0103 (6)
C2A	0.0166 (7)	0.0159 (6)	0.0193 (6)	-0.0018 (5)	-0.0034 (5)	-0.0052 (5)
C2B	0.0180 (7)	0.0176 (6)	0.0186 (6)	-0.0057 (5)	-0.0017 (5)	-0.0034 (5)
C3	0.0123 (6)	0.0174 (6)	0.0173 (6)	-0.0030 (5)	0.0003 (5)	-0.0049 (5)
C3A	0.0182 (7)	0.0214 (7)	0.0198 (7)	0.0021 (6)	-0.0060 (5)	-0.0081 (6)
C3B	0.0285 (8)	0.0149 (6)	0.0187 (7)	-0.0060 (6)	-0.0052 (6)	-0.0042 (5)
C4	0.0173 (7)	0.0137 (6)	0.0165 (6)	-0.0053 (5)	-0.0012 (5)	-0.0028 (5)
C4A	0.0294 (8)	0.0146 (6)	0.0193 (7)	0.0042 (6)	-0.0072 (6)	-0.0061 (5)
C4B	0.0243 (8)	0.0128 (6)	0.0200 (7)	0.0020 (5)	-0.0073 (6)	-0.0060 (5)
C5	0.0144 (7)	0.0142 (6)	0.0222 (7)	-0.0039 (5)	-0.0003 (5)	-0.0021 (5)
C5A	0.0325 (9)	0.0144 (6)	0.0193 (7)	-0.0041 (6)	-0.0043 (6)	-0.0032 (5)
C5B	0.0162 (7)	0.0188 (7)	0.0186 (6)	0.0000 (5)	-0.0041 (5)	-0.0075 (5)
C6	0.0147 (7)	0.0178 (7)	0.0192 (7)	-0.0024 (5)	0.0007 (5)	-0.0026 (5)
C6A	0.0223 (7)	0.0162 (6)	0.0208 (7)	-0.0041 (6)	-0.0015 (6)	-0.0029 (5)
C6B	0.0176 (7)	0.0143 (6)	0.0192 (6)	-0.0036 (5)	-0.0011 (5)	-0.0044 (5)
C7	0.0246 (8)	0.0203 (7)	0.0162 (6)	0.0002 (6)	-0.0034 (6)	-0.0078 (6)
C8	0.0246 (8)	0.0210 (7)	0.0149 (6)	-0.0085 (6)	-0.0005 (5)	-0.0031 (5)
C9	0.0275 (8)	0.0239 (7)	0.0164 (6)	-0.0115 (6)	-0.0009 (6)	-0.0046 (6)
C10	0.0237 (7)	0.0167 (6)	0.0155 (6)	-0.0027 (6)	-0.0013 (5)	-0.0011 (5)
C11	0.0337 (9)	0.0235 (8)	0.0191 (7)	-0.0096 (7)	-0.0014 (6)	-0.0041 (6)
C12	0.0503 (11)	0.0242 (8)	0.0189 (7)	-0.0099 (8)	-0.0037 (7)	-0.0052 (6)
C13	0.0464 (11)	0.0259 (8)	0.0182 (7)	0.0012 (8)	0.0028 (7)	-0.0051 (6)

C14	0.0295 (9)	0.0263 (8)	0.0232 (8)	0.0008 (7)	0.0052 (7)	0.0000 (6)
C15	0.0220 (8)	0.0211 (7)	0.0200 (7)	-0.0004 (6)	-0.0022 (6)	-0.0001 (6)
C16	0.0195 (8)	0.0325 (9)	0.0283 (8)	-0.0076 (7)	-0.0019 (6)	0.0000 (7)
C17	0.0286 (9)	0.0300 (8)	0.0290 (8)	-0.0148 (7)	-0.0065 (7)	-0.0017 (7)
C18	0.0314 (8)	0.0204 (7)	0.0168 (7)	-0.0077 (6)	-0.0052 (6)	-0.0021 (6)
C19	0.0537 (12)	0.0206 (8)	0.0207 (8)	-0.0128 (8)	-0.0081 (7)	-0.0019 (6)
C20	0.0590 (13)	0.0199 (8)	0.0207 (8)	0.0056 (8)	-0.0101 (8)	-0.0046 (6)
C21	0.0375 (10)	0.0329 (9)	0.0219 (8)	0.0107 (8)	-0.0051 (7)	-0.0073 (7)
C22	0.0242 (8)	0.0307 (8)	0.0197 (7)	-0.0003 (7)	-0.0041 (6)	-0.0046 (6)
C23	0.0258 (8)	0.0186 (7)	0.0134 (6)	-0.0029 (6)	-0.0049 (5)	-0.0027 (5)

*Geometric parameters (Å, °)*

N1—C5	1.4973 (18)	C3B—C4B	1.394 (2)
N1—C3	1.4973 (18)	C3B—H3D	0.944 (19)
N1—C2	1.5077 (18)	C4—H4A	0.992 (17)
N1—H1	0.91 (2)	C4—H4B	0.970 (18)
N1A—O6A	1.2087 (19)	C4A—C5A	1.391 (2)
N1A—O7A	1.2250 (19)	C4B—C5B	1.372 (2)
N1A—C6A	1.455 (2)	C5—C6	1.506 (2)
N1B—O6B	1.2256 (16)	C5—H5A	0.947 (19)
N1B—O7B	1.2370 (16)	C5—H5B	0.988 (18)
N1B—C6B	1.4473 (18)	C5A—C6A	1.364 (2)
N2—C4	1.4913 (18)	C5A—H5C	0.95 (2)
N2—C6	1.4944 (19)	C5B—C6B	1.383 (2)
N2—C7	1.4996 (18)	C5B—H5D	0.92 (2)
N2—H2	0.90 (2)	C6—H6A	0.97 (2)
N2A—O3A	1.2316 (16)	C6—H6B	0.968 (19)
N2A—O2A	1.2344 (16)	C7—C8	1.519 (2)
N2A—C2A	1.4451 (19)	C7—H7A	0.99 (2)
N2B—O2B	1.2189 (19)	C7—H7B	0.983 (18)
N2B—O3B	1.2298 (18)	C8—C9	1.532 (2)
N2B—C2B	1.460 (2)	C8—H8A	0.989 (17)
N3—C10	1.4250 (18)	C8—H8B	0.960 (18)
N3—C23	1.4264 (19)	C9—H9A	1.000 (18)
N3—C9	1.4609 (19)	C9—H9B	1.03 (2)
N3A—O4A	1.229 (2)	C10—C11	1.389 (2)
N3A—O5A	1.235 (2)	C10—C15	1.400 (2)
N3A—C4A	1.4408 (19)	C11—C12	1.393 (2)
N3B—O5B	1.2277 (19)	C11—H11	0.95 (2)
N3B—O4B	1.2334 (19)	C12—C13	1.379 (3)
N3B—C4B	1.4493 (19)	C12—H12	0.98 (2)
O1—C1	1.4207 (18)	C13—C14	1.377 (3)
O1—H1C	0.8200	C13—H13	0.97 (2)
O1A—C1A	1.2431 (17)	C14—C15	1.400 (2)
O1B—C1B	1.2504 (18)	C14—H14	0.95 (2)
C1—C2	1.505 (2)	C15—C16	1.462 (2)
C1—H1A	0.97 (2)	C16—C17	1.336 (3)

C1—H1B	0.96 (2)	C16—H16	0.97 (2)
C1A—C2A	1.449 (2)	C17—C18	1.461 (2)
C1A—C6A	1.450 (2)	C17—H17	0.95 (2)
C1B—C2B	1.442 (2)	C18—C23	1.403 (2)
C1B—C6B	1.447 (2)	C18—C19	1.404 (2)
C2—H2A	0.94 (2)	C19—C20	1.379 (3)
C2—H2B	0.965 (18)	C19—H19	0.95 (2)
C2A—C3A	1.379 (2)	C20—C21	1.380 (3)
C2B—C3B	1.364 (2)	C20—H20	0.97 (2)
C3—C4	1.5114 (19)	C21—C22	1.385 (3)
C3—H3A	0.944 (17)	C21—H21	0.95 (2)
C3—H3B	0.956 (18)	C22—C23	1.395 (2)
C3A—C4A	1.381 (2)	C22—H22	1.00 (2)
C3A—H3C	0.960 (19)		
C5—N1—C3	109.19 (11)	N1—C5—H5A	108.6 (11)
C5—N1—C2	110.78 (11)	C6—C5—H5A	108.5 (11)
C3—N1—C2	110.41 (11)	N1—C5—H5B	106.6 (10)
C5—N1—H1	109.1 (12)	C6—C5—H5B	110.4 (11)
C3—N1—H1	106.7 (12)	H5A—C5—H5B	111.4 (15)
C2—N1—H1	110.5 (12)	C6A—C5A—C4A	118.59 (14)
O6A—N1A—O7A	123.06 (16)	C6A—C5A—H5C	120.0 (12)
O6A—N1A—C6A	118.13 (15)	C4A—C5A—H5C	121.4 (12)
O7A—N1A—C6A	118.74 (13)	C4B—C5B—C6B	119.73 (14)
O6B—N1B—O7B	122.01 (13)	C4B—C5B—H5D	122.4 (12)
O6B—N1B—C6B	118.63 (12)	C6B—C5B—H5D	117.8 (12)
O7B—N1B—C6B	119.32 (12)	N2—C6—C5	111.06 (12)
C4—N2—C6	109.13 (11)	N2—C6—H6A	107.4 (12)
C4—N2—C7	112.96 (11)	C5—C6—H6A	107.9 (12)
C6—N2—C7	110.53 (11)	N2—C6—H6B	108.0 (11)
C4—N2—H2	110.9 (13)	C5—C6—H6B	111.3 (11)
C6—N2—H2	106.2 (13)	H6A—C6—H6B	111.0 (16)
C7—N2—H2	106.9 (13)	C5A—C6A—C1A	124.68 (15)
O3A—N2A—O2A	121.74 (13)	C5A—C6A—N1A	117.16 (14)
O3A—N2A—C2A	118.85 (12)	C1A—C6A—N1A	118.16 (13)
O2A—N2A—C2A	119.34 (12)	C5B—C6B—C1B	123.28 (13)
O2B—N2B—O3B	124.29 (15)	C5B—C6B—N1B	116.20 (13)
O2B—N2B—C2B	118.14 (13)	C1B—C6B—N1B	120.47 (12)
O3B—N2B—C2B	117.58 (14)	N2—C7—C8	112.82 (12)
C10—N3—C23	116.26 (11)	N2—C7—H7A	107.2 (11)
C10—N3—C9	117.10 (12)	C8—C7—H7A	112.1 (11)
C23—N3—C9	118.09 (12)	N2—C7—H7B	104.2 (11)
O4A—N3A—O5A	123.17 (14)	C8—C7—H7B	109.4 (11)
O4A—N3A—C4A	118.62 (15)	H7A—C7—H7B	110.9 (15)
O5A—N3A—C4A	118.21 (15)	C7—C8—C9	109.79 (13)
O5B—N3B—O4B	123.64 (14)	C7—C8—H8A	109.1 (10)
O5B—N3B—C4B	118.44 (14)	C9—C8—H8A	109.9 (10)
O4B—N3B—C4B	117.91 (15)	C7—C8—H8B	112.2 (11)

C1—O1—H1C	109.5	C9—C8—H8B	108.8 (11)
O1—C1—C2	110.26 (13)	H8A—C8—H8B	107.1 (14)
O1—C1—H1A	111.6 (12)	N3—C9—C8	110.31 (12)
C2—C1—H1A	111.5 (12)	N3—C9—H9A	105.3 (10)
O1—C1—H1B	109.2 (12)	C8—C9—H9A	110.6 (11)
C2—C1—H1B	106.8 (12)	N3—C9—H9B	114.5 (11)
H1A—C1—H1B	107.3 (16)	C8—C9—H9B	109.6 (11)
O1A—C1A—C2A	126.28 (13)	H9A—C9—H9B	106.4 (15)
O1A—C1A—C6A	121.87 (14)	C11—C10—C15	119.85 (14)
C2A—C1A—C6A	111.85 (12)	C11—C10—N3	121.27 (14)
O1B—C1B—C2B	121.18 (14)	C15—C10—N3	118.80 (14)
O1B—C1B—C6B	127.15 (14)	C10—C11—C12	120.62 (17)
C2B—C1B—C6B	111.66 (12)	C10—C11—H11	120.9 (12)
C1—C2—N1	112.61 (12)	C12—C11—H11	118.4 (13)
C1—C2—H2A	110.8 (12)	C13—C12—C11	119.76 (17)
N1—C2—H2A	106.4 (12)	C13—C12—H12	121.0 (12)
C1—C2—H2B	110.7 (11)	C11—C12—H12	119.2 (12)
N1—C2—H2B	104.6 (11)	C14—C13—C12	119.82 (16)
H2A—C2—H2B	111.4 (16)	C14—C13—H13	119.3 (13)
C3A—C2A—N2A	116.23 (13)	C12—C13—H13	120.9 (13)
C3A—C2A—C1A	123.64 (14)	C13—C14—C15	121.58 (17)
N2A—C2A—C1A	120.05 (12)	C13—C14—H14	120.5 (12)
C3B—C2B—C1B	125.93 (14)	C15—C14—H14	117.8 (12)
C3B—C2B—N2B	117.41 (13)	C10—C15—C14	118.24 (16)
C1B—C2B—N2B	116.65 (13)	C10—C15—C16	122.62 (14)
N1—C3—C4	111.05 (11)	C14—C15—C16	119.13 (16)
N1—C3—H3A	107.0 (10)	C17—C16—C15	126.76 (16)
C4—C3—H3A	111.2 (10)	C17—C16—H16	118.1 (13)
N1—C3—H3B	107.5 (11)	C15—C16—H16	114.5 (13)
C4—C3—H3B	109.9 (11)	C16—C17—C18	127.92 (16)
H3A—C3—H3B	110.0 (14)	C16—C17—H17	117.6 (13)
C2A—C3A—C4A	119.18 (15)	C18—C17—H17	114.2 (13)
C2A—C3A—H3C	118.5 (11)	C23—C18—C19	118.16 (16)
C4A—C3A—H3C	122.3 (11)	C23—C18—C17	122.83 (14)
C2B—C3B—C4B	117.56 (14)	C19—C18—C17	119.01 (16)
C2B—C3B—H3D	121.2 (11)	C20—C19—C18	121.74 (18)
C4B—C3B—H3D	121.2 (11)	C20—C19—H19	120.6 (13)
N2—C4—C3	110.39 (11)	C18—C19—H19	117.6 (13)
N2—C4—H4A	106.7 (10)	C19—C20—C21	119.48 (17)
C3—C4—H4A	112.7 (10)	C19—C20—H20	119.4 (13)
N2—C4—H4B	108.1 (10)	C21—C20—H20	121.1 (13)
C3—C4—H4B	109.6 (10)	C20—C21—C22	120.24 (18)
H4A—C4—H4B	109.3 (14)	C20—C21—H21	119.0 (14)
C3A—C4A—C5A	121.45 (14)	C22—C21—H21	120.7 (14)
C3A—C4A—N3A	119.11 (15)	C21—C22—C23	120.70 (17)
C5A—C4A—N3A	119.27 (15)	C21—C22—H22	119.7 (11)
C5B—C4B—C3B	121.63 (13)	C23—C22—H22	119.6 (11)
C5B—C4B—N3B	119.06 (14)	C22—C23—C18	119.68 (15)

C3B—C4B—N3B	119.21 (14)	C22—C23—N3	121.32 (14)
N1—C5—C6	111.44 (12)	C18—C23—N3	118.92 (14)
O1—C1—C2—N1	-73.47 (17)	O6A—N1A—C6A—C5A	-30.2 (2)
C5—N1—C2—C1	-75.95 (15)	O7A—N1A—C6A—C5A	146.94 (16)
C3—N1—C2—C1	162.96 (13)	O6A—N1A—C6A—C1A	149.37 (18)
O3A—N2A—C2A—C3A	-16.0 (2)	O7A—N1A—C6A—C1A	-33.4 (2)
O2A—N2A—C2A—C3A	167.07 (13)	C4B—C5B—C6B—C1B	0.1 (2)
O3A—N2A—C2A—C1A	160.72 (14)	C4B—C5B—C6B—N1B	-177.32 (13)
O2A—N2A—C2A—C1A	-16.2 (2)	O1B—C1B—C6B—C5B	176.23 (16)
O1A—C1A—C2A—C3A	-172.63 (14)	C2B—C1B—C6B—C5B	-3.3 (2)
C6A—C1A—C2A—C3A	7.6 (2)	O1B—C1B—C6B—N1B	-6.4 (2)
O1A—C1A—C2A—N2A	10.9 (2)	C2B—C1B—C6B—N1B	174.07 (12)
C6A—C1A—C2A—N2A	-168.85 (12)	O6B—N1B—C6B—C5B	10.6 (2)
O1B—C1B—C2B—C3B	-174.15 (15)	O7B—N1B—C6B—C5B	-171.39 (14)
C6B—C1B—C2B—C3B	5.4 (2)	O6B—N1B—C6B—C1B	-166.93 (13)
O1B—C1B—C2B—N2B	4.6 (2)	O7B—N1B—C6B—C1B	11.1 (2)
C6B—C1B—C2B—N2B	-175.85 (12)	C4—N2—C7—C8	72.88 (16)
O2B—N2B—C2B—C3B	-127.14 (16)	C6—N2—C7—C8	-164.52 (12)
O3B—N2B—C2B—C3B	52.44 (19)	N2—C7—C8—C9	160.20 (12)
O2B—N2B—C2B—C1B	54.0 (2)	C10—N3—C9—C8	152.17 (13)
O3B—N2B—C2B—C1B	-126.42 (15)	C23—N3—C9—C8	-61.06 (17)
C5—N1—C3—C4	56.93 (14)	C7—C8—C9—N3	-56.46 (17)
C2—N1—C3—C4	178.96 (12)	C23—N3—C10—C11	-111.87 (16)
N2A—C2A—C3A—C4A	174.47 (13)	C9—N3—C10—C11	35.5 (2)
C1A—C2A—C3A—C4A	-2.1 (2)	C23—N3—C10—C15	71.41 (18)
C1B—C2B—C3B—C4B	-4.1 (2)	C9—N3—C10—C15	-141.22 (14)
N2B—C2B—C3B—C4B	177.17 (13)	C15—C10—C11—C12	2.4 (2)
C6—N2—C4—C3	58.56 (15)	N3—C10—C11—C12	-174.26 (14)
C7—N2—C4—C3	-178.06 (12)	C10—C11—C12—C13	0.6 (3)
N1—C3—C4—N2	-59.29 (15)	C11—C12—C13—C14	-1.8 (3)
C2A—C3A—C4A—C5A	-3.5 (2)	C12—C13—C14—C15	0.0 (3)
C2A—C3A—C4A—N3A	-178.66 (13)	C11—C10—C15—C14	-4.1 (2)
O4A—N3A—C4A—C3A	-1.6 (2)	N3—C10—C15—C14	172.67 (14)
O5A—N3A—C4A—C3A	177.50 (14)	C11—C10—C15—C16	174.51 (15)
O4A—N3A—C4A—C5A	-176.91 (15)	N3—C10—C15—C16	-8.7 (2)
O5A—N3A—C4A—C5A	2.2 (2)	C13—C14—C15—C10	2.9 (2)
C2B—C3B—C4B—C5B	0.2 (2)	C13—C14—C15—C16	-175.71 (16)
C2B—C3B—C4B—N3B	-176.01 (13)	C10—C15—C16—C17	-31.2 (3)
O5B—N3B—C4B—C5B	-9.7 (2)	C14—C15—C16—C17	147.44 (18)
O4B—N3B—C4B—C5B	171.44 (14)	C15—C16—C17—C18	3.0 (3)
O5B—N3B—C4B—C3B	166.71 (14)	C16—C17—C18—C23	30.6 (3)
O4B—N3B—C4B—C3B	-12.2 (2)	C16—C17—C18—C19	-150.09 (18)
C3—N1—C5—C6	-56.04 (15)	C23—C18—C19—C20	-0.2 (2)
C2—N1—C5—C6	-177.86 (12)	C17—C18—C19—C20	-179.58 (16)
C3A—C4A—C5A—C6A	2.6 (2)	C18—C19—C20—C21	0.5 (3)
N3A—C4A—C5A—C6A	177.78 (14)	C19—C20—C21—C22	-0.5 (3)
C3B—C4B—C5B—C6B	1.6 (2)	C20—C21—C22—C23	0.3 (2)

N3B—C4B—C5B—C6B	177.86 (13)	C21—C22—C23—C18	-0.1 (2)
C4—N2—C6—C5	-57.85 (15)	C21—C22—C23—N3	176.53 (14)
C7—N2—C6—C5	177.35 (12)	C19—C18—C23—C22	0.0 (2)
N1—C5—C6—N2	57.52 (15)	C17—C18—C23—C22	179.34 (15)
C4A—C5A—C6A—C1A	4.0 (2)	C19—C18—C23—N3	-176.67 (13)
C4A—C5A—C6A—N1A	-176.42 (14)	C17—C18—C23—N3	2.7 (2)
O1A—C1A—C6A—C5A	171.60 (15)	C10—N3—C23—C22	116.62 (16)
C2A—C1A—C6A—C5A	-8.7 (2)	C9—N3—C23—C22	-30.4 (2)
O1A—C1A—C6A—N1A	-8.0 (2)	C10—N3—C23—C18	-66.76 (18)
C2A—C1A—C6A—N1A	171.77 (13)	C9—N3—C23—C18	146.20 (14)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O1B <sup>i</sup>	0.91 (2)	1.85 (2)	2.6901 (16)	152.6 (18)
N1—H1...O7B <sup>i</sup>	0.91 (2)	2.383 (19)	3.0466 (17)	130.0 (16)
N2—H2...O1A <sup>ii</sup>	0.90 (2)	1.78 (2)	2.6204 (16)	154.6 (19)
N2—H2...O2A <sup>ii</sup>	0.90 (2)	2.43 (2)	3.0711 (16)	128.2 (16)
O1—H1C...O1B <sup>i</sup>	0.82	2.50	3.1600 (19)	138
O1—H1C...O7B <sup>i</sup>	0.82	2.38	3.0841 (18)	144
C2—H2A...O5B <sup>ii</sup>	0.94 (2)	2.43 (2)	3.3130 (19)	155.7 (16)
C2—H2A...O3B <sup>iii</sup>	0.94 (2)	2.60 (2)	3.2350 (19)	125.4 (15)
C3—H3B...O1 <sup>iv</sup>	0.956 (18)	2.410 (18)	3.3250 (18)	160.0 (14)
C3B—H3D...O3B <sup>v</sup>	0.944 (19)	2.503 (19)	3.318 (2)	144.7 (15)
C4—H4B...O2A <sup>ii</sup>	0.970 (18)	2.648 (17)	3.1064 (18)	109.3 (12)
C5—H5A...O2A <sup>vi</sup>	0.947 (19)	2.418 (19)	3.2683 (18)	149.3 (14)
C5—H5B...O1A <sup>ii</sup>	0.988 (18)	2.523 (18)	3.1842 (18)	124.1 (13)
C5—H5B...O5B <sup>ii</sup>	0.988 (18)	2.714 (18)	3.5844 (19)	147.2 (14)
C6—H6B...O5A <sup>i</sup>	0.968 (19)	2.492 (19)	3.3728 (19)	151.2 (15)
C7—H7A...O4A <sup>i</sup>	0.99 (2)	2.44 (2)	3.371 (2)	158.0 (16)
C8—H8A...O2A <sup>ii</sup>	0.989 (17)	2.521 (17)	3.2870 (19)	134.1 (13)
C14—H14...O6A <sup>vii</sup>	0.95 (2)	2.47 (2)	3.085 (2)	122.0 (15)

Symmetry codes: (i)  $x+1, y, z$ ; (ii)  $x, y+1, z$ ; (iii)  $-x+1, -y+1, -z+1$ ; (iv)  $-x+2, -y+1, -z+1$ ; (v)  $-x, -y, -z+1$ ; (vi)  $x+1, y+1, z$ ; (vii)  $-x+2, -y+1, -z$ .