

## 1-[2-(2-Oxo-1,3-oxazolidin-3-yl)ethyl]-4-phenyl-1*H*-1,5-benzodiazepin-2(3*H*)-one

Daouda Ballo,<sup>a</sup> Noureddine Hamou Ahabchane,<sup>a</sup> Hafid Zouihi,<sup>b</sup> El Mokhtar Essassi<sup>a</sup> and Seik Weng Ng<sup>c\*</sup>

<sup>a</sup>Laboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, <sup>b</sup>CNRST Division UATRS, Angle Allal Fassi/FAR, BP 8027 Hay Riad, Rabat, Morocco, and <sup>c</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 7 July 2010; accepted 15 July 2010

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$ ;  $R$  factor = 0.036;  $wR$  factor = 0.102; data-to-parameter ratio = 8.7.

The seven-membered ring in the title compound,  $\text{C}_{20}\text{H}_{19}\text{N}_3\text{O}_3$ , adopts a boat conformation with the two phenylene C atoms representing the stern and the methylene C atom the prow. The dihedral angle between the best plane through the seven-membered ring (r.m.s deviation = 0.358 Å) and the phenyl substituent is 55.8 (1)°. The two rings at either ends of the ethyl chain are staggered [ $\text{N}-\text{CH}_2-\text{CH}_2-\text{N}$  torsion angle = 57.5 (4)°].

### Related literature

For the background to 2,3-dihydro-1*H*-1,5-benzodiazepin-2-ones, see: Ahabchane *et al.* (1999). For a related structure, see: Ballo *et al.* (2010).



### Experimental

#### Crystal data

$\text{C}_{20}\text{H}_{19}\text{N}_3\text{O}_3$	$V = 1704.34\text{ (15) \AA}^3$
$M_r = 349.38$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 9.0163\text{ (5) \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$b = 11.6671\text{ (6) \AA}$	$T = 293\text{ K}$
$c = 16.2019\text{ (8) \AA}$	$0.25 \times 0.25 \times 0.15\text{ mm}$

#### Data collection

Bruker X8 APEXII diffractometer	1578 reflections with $I > 2\sigma(I)$
9253 measured reflections	$R_{\text{int}} = 0.039$
2053 independent reflections	

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$	235 parameters
$wR(F^2) = 0.102$	H-atom parameters constrained
$S = 0.90$	$\Delta\rho_{\text{max}} = 0.12\text{ e \AA}^{-3}$
2053 reflections	$\Delta\rho_{\text{min}} = -0.16\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2052).

### References

- Ahabchane, N. H., Keita, A. & Essassi, E. M. (1999). *C. R. Ser. IIC*, **2**, 519–523.
- Ballo, D., Ahabchane, N. H., Zouihi, H., Essassi, E. M. & Ng, S. W. (2010). *Acta Cryst. E66*, o1277.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

# supporting information

*Acta Cryst.* (2010). E66, o2080 [https://doi.org/10.1107/S160053681002828X]

## 1-[2-(2-Oxo-1,3-oxazolidin-3-yl)ethyl]-4-phenyl-1*H*-1,5-benzodiazepin-2(3*H*)-one

**Daouda Ballo, Noureddine Hamou Ahabchane, Hafid Zouihri, El Mokhtar Essassi and Seik Weng Ng**

### S1. Comment


The background to the class of 2,3-dihydro-1*H*-1,5-benzodiazepin-2-ones is given in an earlier report (Ahabchane *et al.*, 1999). A recent study presented the crystal structure of 1-allyl-4-phenyl-2,3-dihydro-1*H*-1,5-benzodiazepin-2-one (Ballo *et al.*, 2010). The present study has an oxazolidin-2-onyl-ethyl group in place of the allyl group (Scheme I, Fig. 1). The principal feature is the seven-membered ring that is fused to a phenylene ring and adopts a boat-shaped conformation, two phenylene carbons representing the stern and the methylene carbon atom the prow [r.m.s deviation 0.358 Å]. The methyl carbon deviates by 0.637 Å from the best plane. The two rings at either end of the ethyl chain are staggered [N–CH<sub>2</sub>–CH<sub>2</sub>–N torsion angle, 57.5 (4)<sup>o</sup>].

### S2. Experimental

To a solution of 4-phenyl-1*H*-1,5-benzodiazepin-2-one (2 g, 8.4 mmol) in DMF (40 ml) was added dichloroethylamine hydrochloride (0.9 g, 8.4 mmol), potassium carbonate (3 g, 22.2 mmol) and a catalytic quantity of tetra-*n*-butyl-ammonium bromide. The mixture was heated on a sand bath, the reaction monitored by thin layer chromatography. On completion of the reaction, the solvent was evaporated under reduced pressure. The residue was recrystallized from ethanol to afford the title compound as colorless crystals.

### S3. Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.93–0.97 Å) and were included in the refinement in the riding model approximation, with  $U_{\text{iso}}(\text{H})$  set to 1.2–1.5  $U_{\text{eq}}(\text{C})$ . 1486 Friedel pairs were merged in the final cycles of the refinement.

**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of the molecule of  $C_{20}H_{19}N_3O_3$  at the 50% probability level.

### 1-[2-(2-Oxo-1,3-oxazolidin-3-yl)ethyl]-4-phenyl-1*H*-1,5-benzodiazepin-2(3*H*)-one

#### Crystal data

$C_{20}H_{19}N_3O_3$   
 $M_r = 349.38$   
Orthorhombic,  $P2_12_12_1$   
Hall symbol: P 2ac 2ab  
 $a = 9.0163 (5)$  Å  
 $b = 11.6671 (6)$  Å  
 $c = 16.2019 (8)$  Å  
 $V = 1704.34 (15)$  Å<sup>3</sup>  
 $Z = 4$

$F(000) = 736$   
 $D_x = 1.362$  Mg m<sup>-3</sup>  
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 2227 reflections  
 $\theta = 2.9\text{--}21.0^\circ$   
 $\mu = 0.09$  mm<sup>-1</sup>  
 $T = 293$  K  
Prism, colorless  
0.25 × 0.25 × 0.15 mm

#### Data collection

Bruker X8 APEXII  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
9253 measured reflections  
2053 independent reflections

1578 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.039$   
 $\theta_{\text{max}} = 26.7^\circ$ ,  $\theta_{\text{min}} = 2.9^\circ$   
 $h = -11 \rightarrow 10$   
 $k = -14 \rightarrow 13$   
 $l = -20 \rightarrow 14$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.036$   
 $wR(F^2) = 0.102$   
 $S = 0.90$   
2053 reflections

235 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods  
Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0742P)^2]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.12 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.16 \text{ e \AA}^{-3}$$

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.3905 (2)	0.62919 (16)	0.33575 (12)	0.0439 (5)
N2	0.1611 (2)	0.45822 (16)	0.37488 (12)	0.0416 (5)
N3	0.4287 (2)	0.61202 (17)	0.51545 (12)	0.0458 (5)
O1	0.5951 (2)	0.51620 (17)	0.32572 (12)	0.0669 (5)
O2	0.1947 (2)	0.60716 (17)	0.57036 (13)	0.0677 (6)
O3	0.3700 (2)	0.48444 (16)	0.61065 (11)	0.0629 (5)
C1	0.2372 (3)	0.6471 (2)	0.31906 (14)	0.0434 (5)
C2	0.1935 (3)	0.7508 (2)	0.28491 (17)	0.0597 (7)
H2	0.2650	0.8049	0.2710	0.072*
C3	0.0461 (4)	0.7750 (2)	0.27117 (19)	0.0697 (8)
H3	0.0181	0.8445	0.2477	0.084*
C4	-0.0596 (3)	0.6950 (3)	0.29269 (17)	0.0646 (8)
H4	-0.1594	0.7110	0.2841	0.078*
C5	-0.0192 (3)	0.5927 (2)	0.32647 (15)	0.0520 (6)
H5	-0.0921	0.5403	0.3414	0.062*
C6	0.1301 (3)	0.5651 (2)	0.33911 (13)	0.0416 (5)
C7	0.2709 (3)	0.39924 (19)	0.34851 (13)	0.0405 (5)
C8	0.3658 (3)	0.4400 (2)	0.27737 (14)	0.0477 (6)
H8A	0.4248	0.3775	0.2554	0.057*
H8B	0.3046	0.4710	0.2335	0.057*
C9	0.4635 (3)	0.5312 (2)	0.31346 (14)	0.0467 (6)
C10	0.4696 (3)	0.7153 (2)	0.38443 (15)	0.0516 (6)
H10A	0.4560	0.7899	0.3592	0.062*
H10B	0.5748	0.6979	0.3839	0.062*
C11	0.4161 (3)	0.7200 (2)	0.47313 (15)	0.0483 (6)
H11A	0.4732	0.7772	0.5027	0.058*
H11B	0.3131	0.7440	0.4737	0.058*
C12	0.3193 (3)	0.5726 (2)	0.56469 (15)	0.0480 (6)
C13	0.5683 (3)	0.5585 (3)	0.53446 (17)	0.0590 (7)
H13A	0.6327	0.6096	0.5652	0.071*
H13B	0.6190	0.5333	0.4849	0.071*
C14	0.5192 (4)	0.4584 (3)	0.58632 (19)	0.0697 (8)
H14A	0.5224	0.3879	0.5546	0.084*
H14B	0.5826	0.4500	0.6343	0.084*
C15	0.3119 (3)	0.29286 (18)	0.39276 (14)	0.0415 (5)
C16	0.2634 (3)	0.2771 (2)	0.47373 (17)	0.0542 (6)
H16	0.2040	0.3321	0.4989	0.065*
C17	0.3038 (4)	0.1793 (3)	0.51664 (19)	0.0674 (8)
H17	0.2710	0.1690	0.5706	0.081*

C18	0.3914 (3)	0.0977 (3)	0.4807 (2)	0.0658 (8)
H18	0.4195	0.0331	0.5104	0.079*
C19	0.4373 (4)	0.1114 (3)	0.4010 (2)	0.0701 (8)
H19	0.4946	0.0551	0.3759	0.084*
C20	0.3988 (3)	0.2088 (2)	0.35785 (18)	0.0604 (7)
H20	0.4323	0.2180	0.3040	0.072*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0423 (11)	0.0491 (11)	0.0402 (10)	-0.0064 (9)	-0.0003 (9)	0.0029 (9)
N2	0.0395 (11)	0.0438 (10)	0.0415 (10)	-0.0009 (9)	-0.0001 (8)	-0.0040 (9)
N3	0.0414 (11)	0.0531 (11)	0.0428 (11)	0.0005 (9)	0.0037 (9)	0.0065 (9)
O1	0.0407 (10)	0.0837 (13)	0.0761 (13)	0.0036 (10)	0.0055 (9)	0.0061 (12)
O2	0.0511 (11)	0.0706 (12)	0.0815 (13)	0.0082 (11)	0.0214 (10)	0.0049 (10)
O3	0.0706 (12)	0.0607 (11)	0.0575 (11)	0.0065 (10)	0.0181 (10)	0.0150 (9)
C1	0.0470 (13)	0.0482 (13)	0.0351 (12)	0.0017 (11)	-0.0029 (11)	-0.0027 (11)
C2	0.0711 (18)	0.0510 (14)	0.0570 (16)	0.0016 (14)	-0.0068 (15)	0.0076 (12)
C3	0.080 (2)	0.0593 (16)	0.0695 (19)	0.0250 (16)	-0.0144 (17)	0.0092 (15)
C4	0.0580 (17)	0.0736 (18)	0.0623 (17)	0.0226 (16)	-0.0112 (15)	-0.0051 (15)
C5	0.0434 (13)	0.0644 (16)	0.0483 (14)	0.0065 (12)	-0.0018 (11)	-0.0093 (13)
C6	0.0447 (12)	0.0465 (13)	0.0335 (11)	0.0032 (10)	-0.0008 (10)	-0.0061 (10)
C7	0.0412 (13)	0.0444 (11)	0.0358 (11)	-0.0024 (10)	-0.0031 (10)	-0.0058 (10)
C8	0.0558 (14)	0.0542 (13)	0.0330 (11)	0.0057 (12)	0.0038 (11)	-0.0049 (11)
C9	0.0436 (14)	0.0599 (14)	0.0367 (12)	0.0005 (12)	0.0069 (11)	0.0080 (11)
C10	0.0551 (14)	0.0547 (13)	0.0451 (13)	-0.0174 (12)	-0.0012 (12)	0.0062 (11)
C11	0.0559 (15)	0.0455 (13)	0.0435 (13)	-0.0049 (11)	0.0009 (12)	0.0003 (10)
C12	0.0544 (15)	0.0460 (13)	0.0434 (13)	0.0008 (12)	0.0077 (11)	-0.0045 (11)
C13	0.0464 (14)	0.0794 (19)	0.0513 (15)	0.0070 (13)	0.0019 (12)	0.0094 (14)
C14	0.0655 (18)	0.0788 (19)	0.0650 (17)	0.0178 (16)	0.0061 (15)	0.0164 (16)
C15	0.0388 (11)	0.0417 (11)	0.0440 (13)	-0.0032 (10)	-0.0055 (10)	-0.0049 (10)
C16	0.0557 (15)	0.0557 (14)	0.0512 (15)	-0.0024 (12)	-0.0012 (13)	0.0025 (13)
C17	0.0756 (19)	0.0734 (18)	0.0532 (16)	-0.0086 (17)	-0.0077 (15)	0.0144 (15)
C18	0.0613 (17)	0.0592 (17)	0.077 (2)	0.0008 (14)	-0.0237 (16)	0.0153 (15)
C19	0.074 (2)	0.0600 (17)	0.077 (2)	0.0218 (15)	-0.0054 (17)	-0.0025 (15)
C20	0.0643 (18)	0.0614 (16)	0.0554 (16)	0.0138 (14)	-0.0014 (14)	-0.0048 (13)

*Geometric parameters ( $\text{\AA}$ ,  $\text{^\circ}$ )*

N1—C9	1.367 (3)	C8—C9	1.501 (3)
N1—C1	1.424 (3)	C8—H8A	0.9700
N1—C10	1.463 (3)	C8—H8B	0.9700
N2—C7	1.280 (3)	C10—C11	1.517 (4)
N2—C6	1.403 (3)	C10—H10A	0.9700
N3—C12	1.350 (3)	C10—H10B	0.9700
N3—C11	1.439 (3)	C11—H11A	0.9700
N3—C13	1.439 (3)	C11—H11B	0.9700
O1—C9	1.216 (3)	C13—C14	1.505 (4)

O2—C12	1.197 (3)	C13—H13A	0.9700
O3—C12	1.350 (3)	C13—H13B	0.9700
O3—C14	1.435 (4)	C14—H14A	0.9700
C1—C2	1.387 (3)	C14—H14B	0.9700
C1—C6	1.398 (3)	C15—C20	1.377 (3)
C2—C3	1.377 (4)	C15—C16	1.395 (4)
C2—H2	0.9300	C16—C17	1.385 (4)
C3—C4	1.378 (4)	C16—H16	0.9300
C3—H3	0.9300	C17—C18	1.367 (4)
C4—C5	1.363 (4)	C17—H17	0.9300
C4—H4	0.9300	C18—C19	1.364 (5)
C5—C6	1.399 (3)	C18—H18	0.9300
C5—H5	0.9300	C19—C20	1.379 (4)
C7—C15	1.480 (3)	C19—H19	0.9300
C7—C8	1.512 (3)	C20—H20	0.9300
C9—N1—C1	122.7 (2)	C11—C10—H10B	109.1
C9—N1—C10	118.8 (2)	H10A—C10—H10B	107.9
C1—N1—C10	118.3 (2)	N3—C11—C10	113.2 (2)
C7—N2—C6	119.6 (2)	N3—C11—H11A	108.9
C12—N3—C11	121.5 (2)	C10—C11—H11A	108.9
C12—N3—C13	111.40 (19)	N3—C11—H11B	108.9
C11—N3—C13	123.5 (2)	C10—C11—H11B	108.9
C12—O3—C14	109.1 (2)	H11A—C11—H11B	107.7
C2—C1—C6	119.6 (2)	O2—C12—O3	122.2 (2)
C2—C1—N1	118.7 (2)	O2—C12—N3	128.1 (2)
C6—C1—N1	121.7 (2)	O3—C12—N3	109.8 (2)
C3—C2—C1	121.2 (3)	N3—C13—C14	101.5 (2)
C3—C2—H2	119.4	N3—C13—H13A	111.5
C1—C2—H2	119.4	C14—C13—H13A	111.5
C2—C3—C4	119.1 (3)	N3—C13—H13B	111.5
C2—C3—H3	120.4	C14—C13—H13B	111.5
C4—C3—H3	120.4	H13A—C13—H13B	109.3
C5—C4—C3	120.7 (3)	O3—C14—C13	105.4 (2)
C5—C4—H4	119.7	O3—C14—H14A	110.7
C3—C4—H4	119.7	C13—C14—H14A	110.7
C4—C5—C6	121.1 (3)	O3—C14—H14B	110.7
C4—C5—H5	119.4	C13—C14—H14B	110.7
C6—C5—H5	119.4	H14A—C14—H14B	108.8
C1—C6—C5	118.3 (2)	C20—C15—C16	118.1 (2)
C1—C6—N2	124.5 (2)	C20—C15—C7	122.7 (2)
C5—C6—N2	117.2 (2)	C16—C15—C7	119.2 (2)
N2—C7—C15	118.8 (2)	C17—C16—C15	119.9 (3)
N2—C7—C8	121.6 (2)	C17—C16—H16	120.0
C15—C7—C8	119.5 (2)	C15—C16—H16	120.0
C9—C8—C7	104.95 (17)	C18—C17—C16	120.7 (3)
C9—C8—H8A	110.8	C18—C17—H17	119.6
C7—C8—H8A	110.8	C16—C17—H17	119.6

C9—C8—H8B	110.8	C19—C18—C17	119.8 (3)
C7—C8—H8B	110.8	C19—C18—H18	120.1
H8A—C8—H8B	108.8	C17—C18—H18	120.1
O1—C9—N1	123.2 (2)	C18—C19—C20	120.0 (3)
O1—C9—C8	122.3 (2)	C18—C19—H19	120.0
N1—C9—C8	114.4 (2)	C20—C19—H19	120.0
N1—C10—C11	112.4 (2)	C15—C20—C19	121.4 (3)
N1—C10—H10A	109.1	C15—C20—H20	119.3
C11—C10—H10A	109.1	C19—C20—H20	119.3
N1—C10—H10B	109.1		
C9—N1—C1—C2	-132.7 (2)	C9—N1—C10—C11	-108.0 (2)
C10—N1—C1—C2	52.3 (3)	C1—N1—C10—C11	67.3 (3)
C9—N1—C1—C6	49.8 (3)	C12—N3—C11—C10	-137.1 (2)
C10—N1—C1—C6	-125.2 (2)	C13—N3—C11—C10	66.2 (3)
C6—C1—C2—C3	0.7 (4)	N1—C10—C11—N3	57.5 (3)
N1—C1—C2—C3	-176.8 (3)	C14—O3—C12—O2	176.3 (3)
C1—C2—C3—C4	0.7 (5)	C14—O3—C12—N3	-4.3 (3)
C2—C3—C4—C5	-0.6 (4)	C11—N3—C12—O2	12.8 (4)
C3—C4—C5—C6	-1.0 (4)	C13—N3—C12—O2	172.0 (3)
C2—C1—C6—C5	-2.2 (3)	C11—N3—C12—O3	-166.5 (2)
N1—C1—C6—C5	175.2 (2)	C13—N3—C12—O3	-7.3 (3)
C2—C1—C6—N2	-179.1 (2)	C12—N3—C13—C14	14.8 (3)
N1—C1—C6—N2	-1.6 (3)	C11—N3—C13—C14	173.5 (2)
C4—C5—C6—C1	2.4 (4)	C12—O3—C14—C13	13.4 (3)
C4—C5—C6—N2	179.5 (2)	N3—C13—C14—O3	-16.4 (3)
C7—N2—C6—C1	-42.9 (3)	N2—C7—C15—C20	161.8 (2)
C7—N2—C6—C5	140.2 (2)	C8—C7—C15—C20	-22.5 (3)
C6—N2—C7—C15	173.44 (19)	N2—C7—C15—C16	-19.4 (3)
C6—N2—C7—C8	-2.2 (3)	C8—C7—C15—C16	156.3 (2)
N2—C7—C8—C9	76.5 (3)	C20—C15—C16—C17	0.4 (4)
C15—C7—C8—C9	-99.1 (2)	C7—C15—C16—C17	-178.4 (2)
C1—N1—C9—O1	178.2 (2)	C15—C16—C17—C18	0.2 (4)
C10—N1—C9—O1	-6.8 (3)	C16—C17—C18—C19	-1.3 (4)
C1—N1—C9—C8	-5.5 (3)	C17—C18—C19—C20	1.8 (5)
C10—N1—C9—C8	169.45 (19)	C16—C15—C20—C19	0.1 (4)
C7—C8—C9—O1	107.4 (3)	C7—C15—C20—C19	178.9 (3)
C7—C8—C9—N1	-68.8 (2)	C18—C19—C20—C15	-1.2 (5)