

Di-n-butylammonium 2-[(3,5-di-*tert*-butyl-4-hydroxybenzyl)sulfanyl]benzoate

Abeer A. Alhadi,^a Hamid Khaledi,^{a*} Hapipah Mohd Ali^a and Marilyn M. Olmstead^b

^aDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^bDepartment of Chemistry, University of California, One Shields Avenue, Davis, CA 95616, USA

Correspondence e-mail: khaledi@siswa.um.edu.my


Received 16 July 2010; accepted 23 August 2010

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.050; wR factor = 0.138; data-to-parameter ratio = 15.8.

The title salt, $\text{C}_8\text{H}_{20}\text{N}^+\cdot\text{C}_{22}\text{H}_{27}\text{O}_3\text{S}^-$, is a proton-transfer compound derived from the recently reported parent carboxylic acid [Alhadi *et al.* (2010). *Acta Cryst. E66*, o1787] by the addition of a second equivalent of di-n-butylamine, yielding the di-n-butylammonium carboxylate salt. The structure of the carboxylate anion resembles that of the parent carboxylic acid. The main difference lies in the position of the H atom in the 4-hydroxy group. In the anion the O—H bond is perpendicular, rather than parallel, to the benzyl ring. This position appears to facilitate hydrogen bonding to an O atom of the carboxylate group of a symmetry-related anion. In addition, there are three N—H···O hydrogen bonds. In contrast, the neutral species hydrogen bonds *via* a carboxylic acid dimer. The dihedral angle between the benzene rings in the anion is $79.19(7)^\circ$.

Related literature

For the structure of the parent benzoic acid, see: Alhadi *et al.* (2010). For a similar structure based on nicotinic acid, see: Mansor *et al.* (2008).



Experimental

Crystal data

$\text{C}_8\text{H}_{20}\text{N}^+\cdot\text{C}_{22}\text{H}_{27}\text{O}_3\text{S}^-$	$V = 5973.9(4)\text{ \AA}^3$
$M_r = 501.75$	$Z = 8$
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation
$a = 12.8631(5)\text{ \AA}$	$\mu = 0.14\text{ mm}^{-1}$
$b = 20.1109(9)\text{ \AA}$	$T = 296\text{ K}$
$c = 23.0930(9)\text{ \AA}$	$0.60 \times 0.40 \times 0.35\text{ mm}$

Data collection

Bruker APEXII diffractometer	44646 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	5277 independent reflections
$(SADABS; Sheldrick, 1996)$	3311 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.922$, $T_{\max} = 0.954$	$R_{\text{int}} = 0.080$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.138$	$\Delta\rho_{\text{max}} = 0.28\text{ e \AA}^{-3}$
$S = 1.01$	$\Delta\rho_{\text{min}} = -0.17\text{ e \AA}^{-3}$
5277 reflections	
333 parameters	
3 restraints	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O3—H3···O1 ⁱ	0.81 (2)	2.02 (2)	2.723 (3)	146 (3)
N1—H1A···O2	0.89 (2)	1.88 (2)	2.767 (3)	177 (3)
N1—H1B···O1 ⁱⁱ	0.89 (2)	2.00 (2)	2.806 (3)	150 (3)
N1—H1B···O2 ⁱⁱ	0.89 (2)	2.36 (2)	3.167 (3)	150 (2)
N1—H1B···O2 ⁱⁱ	0.89 (2)	2.36 (2)	3.167 (3)	150 (2)

Symmetry codes: (i) $x + 1, y, z$; (ii) $-x + 1, -y, -z + 1$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* (Bruker, 2007); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors thank the University of Malaya for funding this study (FRGS grant FP009/2008 C).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2302).

References

- Alhadi, A. A., Khaledi, H., Mohd Ali, H. & Olmstead, M. M. (2010). *Acta Cryst. E66*, o1787.
- Bruker (2007). *APEX2*, *SAINT* and *XP*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Mansor, S., Yehye, W. A., Ariffin, A., Rahman, N. A. & Ng, S. W. (2008). *Acta Cryst. E64*, o1799.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst. 43*, 920–925.

supporting information

Acta Cryst. (2010). E66, o2454 [https://doi.org/10.1107/S1600536810033921]

Di-*n*-butylammonium 2-[(3,5-di-*tert*-butyl-4-hydroxybenzyl)sulfanyl]benzoate

Abeer A. Alhadi, Hamid Khaledi, Hapipah Mohd Ali and Marilyn M. Olmstead

S1. Comment


As shown in Fig. 1, the title compound is a di-*n*-butylammonium salt. It is a proton transfer derivative of the previously reported benzoic acid analog (Alhadi *et al.*, 2010). In contrast to the structure of the parent carboxylic acid in which the C—O—H fragment was found coplanar to the aromatic ring and therefore not involved in any hydrogen bonding, in the present structure the O—H bond is perpendicular to the aromatic ring, and it participates in hydrogen bonding to the carboxylate group of the neighboring anion, forming an infinite chain along the *a* axis (Fig. 2). This arrangement is similar to that reported for the salt based on nicotinic acid (Mansor *et al.*, 2008). The dihedral angle between the aromatic rings C2/C3/C4/C5/C6/C7 and C9/C10/C11/C12/C13/C14 is 79.19 (7)°.

S2. Experimental


Thiosalicylic acid (0.154 g, 1 mmol), 2,6-di-*t*-butylphenol (2.00 g, 1 mmol) and *para*formaldehyde (0.291 g, 1 mmol) were ground into a homogenous powder and to this was added di-*n*-butylamine (0.1 ml). The slurry was heated to 353 K for 2.5 h, then cooled to 323 K. Ethanol (20 ml) was added and the mixture was stirred for 1 h at room temperature. To the resulting clear solution di-*n*-butylamine (0.1 ml) was added and the solution was set aside in the dark for 5 days whereupon the colorless crystals of the title compound were obtained.

S3. Refinement


The C-bound hydrogen atoms were placed at calculated positions (C—H 0.93–0.97 Å) and were treated as riding on their parent atoms. The nitrogen- and oxygen-bound hydrogen atoms were located in a difference map and were refined freely with distances restrained to N—H 0.86 (2) and O—H 0.82 (2) Å. For all H atoms, $U_{\text{iso}}(\text{H})$ was set to 1.2–1.5 U_{eq} (carrier atom).

**Figure 1**

Thermal ellipsoid plot of the title compound at the 30% probability level. H atoms participating in hydrogen bonding are drawn as spheres of arbitrary radius.

**Figure 2**

A view of the hydrogen bonding interactions (dashed lines) as viewed down *b*. Symmetry codes: ' = $x - 1, y, z$; " = $-x + 1, -y, -z + 1$.

Di-n-butylammonium 2-[3,5-di-*tert*-butyl-4-hydroxybenzyl]sulfanyl]benzoate*Crystal data* $M_r = 501.75$ Orthorhombic, $Pbca$

Hall symbol: -P 2ac 2ab

 $a = 12.8631 (5) \text{ \AA}$ $b = 20.1109 (9) \text{ \AA}$ $c = 23.0930 (9) \text{ \AA}$ $V = 5973.9 (4) \text{ \AA}^3$ $Z = 8$ $F(000) = 2192$ $D_x = 1.116 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3944 reflections

 $\theta = 2.4\text{--}19.4^\circ$ $\mu = 0.14 \text{ mm}^{-1}$ $T = 296 \text{ K}$

Irregular block, colourless

 $0.60 \times 0.40 \times 0.35 \text{ mm}$ *Data collection*

Bruker APEXII

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\min} = 0.922$, $T_{\max} = 0.954$

44646 measured reflections

5277 independent reflections

3311 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.080$ $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.8^\circ$ $h = -15 \rightarrow 15$ $k = -23 \rightarrow 23$ $l = -27 \rightarrow 27$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.050$ $wR(F^2) = 0.138$ $S = 1.01$

5277 reflections

333 parameters

3 restraints

0 constraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0533P)^2 + 2.2704P]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.28 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.17 \text{ e \AA}^{-3}$ *Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)*

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.4142 (2)	0.10279 (13)	0.41431 (11)	0.0573 (7)
S1	0.64187 (5)	0.14037 (4)	0.37500 (3)	0.0571 (2)
O1	0.32489 (16)	0.08674 (12)	0.42807 (11)	0.1060 (9)
O2	0.49152 (15)	0.07570 (10)	0.43550 (9)	0.0799 (6)
O3	1.11548 (13)	0.06947 (10)	0.41947 (9)	0.0676 (5)
H3	1.1701 (18)	0.0878 (16)	0.4133 (14)	0.101*
N1	0.65789 (17)	0.01880 (12)	0.49304 (11)	0.0639 (6)
H1A	0.6030 (17)	0.0360 (13)	0.4749 (11)	0.077*
H1B	0.639 (2)	-0.0157 (11)	0.5148 (11)	0.077*
C2	0.42822 (17)	0.15713 (11)	0.37050 (9)	0.0443 (6)
C3	0.34012 (19)	0.18891 (13)	0.34973 (11)	0.0552 (6)
H3A	0.2754	0.1757	0.3635	0.066*
C4	0.3448 (2)	0.23882 (15)	0.30980 (12)	0.0669 (8)

H4	0.2843	0.2591	0.2966	0.080*
C5	0.4395 (2)	0.25874 (15)	0.28939 (13)	0.0732 (9)
H5	0.4436	0.2928	0.2623	0.088*
C6	0.5294 (2)	0.22837 (14)	0.30893 (12)	0.0648 (7)
H6	0.5934	0.2422	0.2945	0.078*
C7	0.52605 (17)	0.17767 (12)	0.34970 (10)	0.0459 (6)
C8	0.74329 (17)	0.18495 (13)	0.33648 (11)	0.0579 (7)
H8A	0.7354	0.1789	0.2950	0.069*
H8B	0.7394	0.2321	0.3450	0.069*
C9	0.84580 (18)	0.15715 (13)	0.35632 (11)	0.0512 (6)
C10	0.89894 (18)	0.11082 (12)	0.32410 (11)	0.0516 (6)
H10	0.8716	0.0979	0.2885	0.062*
C11	0.99201 (17)	0.08236 (12)	0.34249 (10)	0.0482 (6)
C12	1.02979 (17)	0.10199 (12)	0.39672 (10)	0.0486 (6)
C13	0.97987 (18)	0.15064 (13)	0.43035 (10)	0.0524 (6)
C14	0.88760 (18)	0.17683 (13)	0.40858 (11)	0.0556 (7)
H14	0.8527	0.2089	0.4302	0.067*
C15	1.0201 (2)	0.17282 (17)	0.49001 (12)	0.0734 (9)
C16	0.9545 (3)	0.2305 (2)	0.51411 (17)	0.1397 (19)
H16A	0.9577	0.2675	0.4879	0.209*
H16B	0.9814	0.2436	0.5512	0.209*
H16C	0.8837	0.2164	0.5183	0.209*
C17	1.0103 (3)	0.1149 (2)	0.53240 (13)	0.1155 (15)
H17A	1.0507	0.0780	0.5186	0.173*
H17B	0.9387	0.1020	0.5355	0.173*
H17C	1.0354	0.1284	0.5697	0.173*
C18	1.1328 (2)	0.19722 (19)	0.48760 (15)	0.0960 (11)
H18A	1.1779	0.1605	0.4790	0.144*
H18B	1.1516	0.2161	0.5243	0.144*
H18C	1.1395	0.2304	0.4579	0.144*
C19	1.04977 (19)	0.03203 (13)	0.30369 (11)	0.0579 (7)
C20	0.9917 (2)	0.02028 (16)	0.24637 (13)	0.0822 (9)
H20A	1.0298	-0.0109	0.2231	0.123*
H20B	0.9854	0.0616	0.2258	0.123*
H20C	0.9237	0.0028	0.2543	0.123*
C21	1.1572 (2)	0.05973 (17)	0.28782 (13)	0.0846 (10)
H21A	1.2001	0.0610	0.3218	0.127*
H21B	1.1497	0.1039	0.2725	0.127*
H21C	1.1889	0.0317	0.2592	0.127*
C22	1.0605 (2)	-0.03551 (15)	0.33351 (14)	0.0824 (9)
H22A	1.0975	-0.0655	0.3086	0.124*
H22B	0.9926	-0.0532	0.3415	0.124*
H22C	1.0980	-0.0302	0.3691	0.124*
C23	0.7166 (4)	-0.1192 (2)	0.3170 (2)	0.1382 (17)
H23A	0.6944	-0.1593	0.3359	0.207*
H23B	0.7700	-0.1295	0.2894	0.207*
H23C	0.6585	-0.0993	0.2974	0.207*
C24	0.7569 (3)	-0.0731 (3)	0.35968 (18)	0.1261 (15)

H24A	0.7790	-0.0331	0.3397	0.151*
H24B	0.8183	-0.0928	0.3771	0.151*
C25	0.6833 (3)	-0.05329 (18)	0.40777 (15)	0.0897 (10)
H25A	0.6222	-0.0325	0.3910	0.108*
H25B	0.6608	-0.0929	0.4283	0.108*
C26	0.7327 (2)	-0.00614 (17)	0.44985 (13)	0.0771 (9)
H26A	0.7892	-0.0286	0.4696	0.093*
H26B	0.7619	0.0312	0.4288	0.093*
C27	0.7013 (2)	0.06935 (16)	0.53258 (14)	0.0783 (9)
H27A	0.7257	0.1071	0.5102	0.094*
H27B	0.7603	0.0507	0.5530	0.094*
C28	0.6206 (3)	0.09282 (17)	0.57620 (15)	0.0901 (10)
H28A	0.5647	0.1150	0.5557	0.108*
H28B	0.5914	0.0544	0.5957	0.108*
C29	0.6642 (3)	0.1391 (2)	0.62032 (18)	0.1121 (13)
H29A	0.6988	0.1755	0.6006	0.135*
H29B	0.7160	0.1157	0.6430	0.135*
C30	0.5837 (3)	0.1671 (3)	0.6604 (2)	0.1449 (18)
H30A	0.5335	0.1918	0.6384	0.217*
H30B	0.6164	0.1959	0.6880	0.217*
H30C	0.5494	0.1314	0.6804	0.217*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0437 (15)	0.0638 (17)	0.0644 (17)	0.0003 (14)	0.0004 (13)	0.0071 (14)
S1	0.0381 (3)	0.0675 (4)	0.0658 (4)	0.0052 (3)	-0.0047 (3)	0.0172 (3)
O1	0.0492 (12)	0.1187 (19)	0.150 (2)	-0.0007 (12)	0.0137 (13)	0.0679 (17)
O2	0.0560 (12)	0.0903 (15)	0.0934 (15)	0.0010 (11)	-0.0136 (10)	0.0372 (12)
O3	0.0384 (10)	0.0833 (14)	0.0810 (13)	0.0067 (10)	-0.0051 (9)	0.0132 (11)
N1	0.0461 (13)	0.0715 (17)	0.0741 (17)	-0.0010 (12)	-0.0089 (12)	0.0213 (13)
C2	0.0394 (13)	0.0497 (14)	0.0437 (13)	0.0041 (11)	-0.0048 (10)	-0.0035 (11)
C3	0.0408 (14)	0.0645 (17)	0.0603 (16)	0.0078 (12)	-0.0032 (12)	-0.0009 (14)
C4	0.0497 (16)	0.080 (2)	0.0707 (18)	0.0224 (15)	-0.0118 (13)	0.0079 (16)
C5	0.0632 (18)	0.080 (2)	0.076 (2)	0.0161 (16)	-0.0034 (15)	0.0288 (16)
C6	0.0465 (15)	0.0763 (19)	0.0715 (18)	0.0088 (14)	0.0009 (13)	0.0221 (16)
C7	0.0400 (13)	0.0513 (14)	0.0462 (14)	0.0064 (11)	-0.0047 (10)	0.0040 (12)
C8	0.0407 (14)	0.0651 (17)	0.0679 (17)	0.0015 (13)	-0.0041 (12)	0.0126 (14)
C9	0.0388 (13)	0.0586 (16)	0.0562 (15)	-0.0020 (12)	-0.0018 (11)	0.0095 (12)
C10	0.0429 (13)	0.0609 (16)	0.0510 (15)	-0.0045 (13)	-0.0015 (11)	0.0036 (13)
C11	0.0379 (13)	0.0557 (15)	0.0512 (15)	-0.0006 (11)	0.0068 (11)	0.0065 (12)
C12	0.0309 (12)	0.0614 (16)	0.0536 (15)	-0.0005 (12)	0.0056 (11)	0.0088 (12)
C13	0.0375 (13)	0.0700 (17)	0.0497 (15)	-0.0021 (12)	0.0020 (11)	0.0026 (13)
C14	0.0413 (14)	0.0644 (17)	0.0612 (17)	0.0033 (12)	0.0051 (12)	-0.0020 (13)
C15	0.0501 (16)	0.113 (2)	0.0569 (18)	0.0101 (17)	-0.0042 (13)	-0.0163 (17)
C16	0.106 (3)	0.208 (5)	0.105 (3)	0.055 (3)	-0.034 (2)	-0.092 (3)
C17	0.072 (2)	0.217 (5)	0.058 (2)	-0.007 (3)	-0.0067 (17)	0.031 (3)
C18	0.070 (2)	0.128 (3)	0.090 (2)	-0.016 (2)	-0.0125 (17)	-0.031 (2)

C19	0.0473 (15)	0.0646 (17)	0.0616 (17)	0.0025 (13)	0.0091 (12)	-0.0025 (14)
C20	0.084 (2)	0.091 (2)	0.071 (2)	0.0077 (18)	0.0033 (17)	-0.0192 (18)
C21	0.0661 (19)	0.102 (3)	0.086 (2)	-0.0117 (18)	0.0308 (16)	-0.0162 (19)
C22	0.079 (2)	0.069 (2)	0.099 (2)	0.0095 (17)	0.0052 (18)	0.0039 (18)
C23	0.154 (4)	0.123 (3)	0.138 (4)	0.040 (3)	-0.021 (3)	-0.026 (3)
C24	0.081 (3)	0.181 (5)	0.116 (3)	0.021 (3)	-0.007 (2)	-0.011 (3)
C25	0.071 (2)	0.099 (3)	0.099 (3)	0.0172 (19)	-0.0035 (19)	0.003 (2)
C26	0.0496 (17)	0.098 (2)	0.084 (2)	0.0130 (17)	-0.0005 (15)	0.0249 (19)
C27	0.0577 (18)	0.088 (2)	0.089 (2)	-0.0107 (17)	-0.0169 (16)	0.0154 (18)
C28	0.069 (2)	0.095 (3)	0.107 (3)	-0.0073 (19)	-0.0078 (19)	-0.009 (2)
C29	0.090 (3)	0.130 (3)	0.117 (3)	-0.010 (2)	-0.009 (2)	-0.021 (3)
C30	0.105 (3)	0.183 (5)	0.147 (4)	-0.021 (3)	0.022 (3)	-0.055 (4)

Geometric parameters (\AA , $^\circ$)

C1—O2	1.235 (3)	C17—H17B	0.9600
C1—O1	1.235 (3)	C17—H17C	0.9600
C1—C2	1.500 (3)	C18—H18A	0.9600
S1—C7	1.767 (2)	C18—H18B	0.9600
S1—C8	1.816 (2)	C18—H18C	0.9600
O3—C12	1.385 (3)	C19—C22	1.529 (4)
O3—H3	0.806 (18)	C19—C21	1.534 (4)
N1—C26	1.474 (4)	C19—C20	1.538 (4)
N1—C27	1.476 (4)	C20—H20A	0.9600
N1—H1A	0.890 (17)	C20—H20B	0.9600
N1—H1B	0.892 (17)	C20—H20C	0.9600
C2—C3	1.387 (3)	C21—H21A	0.9600
C2—C7	1.409 (3)	C21—H21B	0.9600
C3—C4	1.364 (4)	C21—H21C	0.9600
C3—H3A	0.9300	C22—H22A	0.9600
C4—C5	1.366 (4)	C22—H22B	0.9600
C4—H4	0.9300	C22—H22C	0.9600
C5—C6	1.383 (3)	C23—C24	1.450 (5)
C5—H5	0.9300	C23—H23A	0.9600
C6—C7	1.389 (3)	C23—H23B	0.9600
C6—H6	0.9300	C23—H23C	0.9600
C8—C9	1.504 (3)	C24—C25	1.513 (5)
C8—H8A	0.9700	C24—H24A	0.9700
C8—H8B	0.9700	C24—H24B	0.9700
C9—C10	1.374 (3)	C25—C26	1.499 (4)
C9—C14	1.379 (3)	C25—H25A	0.9700
C10—C11	1.393 (3)	C25—H25B	0.9700
C10—H10	0.9300	C26—H26A	0.9700
C11—C12	1.400 (3)	C26—H26B	0.9700
C11—C19	1.543 (3)	C27—C28	1.521 (4)
C12—C13	1.405 (3)	C27—H27A	0.9700
C13—C14	1.392 (3)	C27—H27B	0.9700
C13—C15	1.538 (4)	C28—C29	1.490 (5)

C14—H14	0.9300	C28—H28A	0.9700
C15—C17	1.526 (5)	C28—H28B	0.9700
C15—C18	1.532 (4)	C29—C30	1.498 (5)
C15—C16	1.538 (4)	C29—H29A	0.9700
C16—H16A	0.9600	C29—H29B	0.9700
C16—H16B	0.9600	C30—H30A	0.9600
C16—H16C	0.9600	C30—H30B	0.9600
C17—H17A	0.9600	C30—H30C	0.9600
O2—C1—O1	122.1 (3)	H18A—C18—H18C	109.5
O2—C1—C2	119.5 (2)	H18B—C18—H18C	109.5
O1—C1—C2	118.4 (2)	C22—C19—C21	110.4 (2)
C7—S1—C8	103.55 (11)	C22—C19—C20	107.2 (2)
C12—O3—H3	114 (3)	C21—C19—C20	106.7 (2)
C26—N1—C27	114.0 (2)	C22—C19—C11	111.4 (2)
C26—N1—H1A	109.4 (18)	C21—C19—C11	109.5 (2)
C27—N1—H1A	108.8 (18)	C20—C19—C11	111.5 (2)
C26—N1—H1B	107.3 (18)	C19—C20—H20A	109.5
C27—N1—H1B	106.9 (18)	C19—C20—H20B	109.5
H1A—N1—H1B	110 (3)	H20A—C20—H20B	109.5
C3—C2—C7	118.5 (2)	C19—C20—H20C	109.5
C3—C2—C1	118.1 (2)	H20A—C20—H20C	109.5
C7—C2—C1	123.4 (2)	H20B—C20—H20C	109.5
C4—C3—C2	122.5 (2)	C19—C21—H21A	109.5
C4—C3—H3A	118.8	C19—C21—H21B	109.5
C2—C3—H3A	118.8	H21A—C21—H21B	109.5
C3—C4—C5	119.2 (2)	C19—C21—H21C	109.5
C3—C4—H4	120.4	H21A—C21—H21C	109.5
C5—C4—H4	120.4	H21B—C21—H21C	109.5
C4—C5—C6	120.2 (3)	C19—C22—H22A	109.5
C4—C5—H5	119.9	C19—C22—H22B	109.5
C6—C5—H5	119.9	H22A—C22—H22B	109.5
C5—C6—C7	121.3 (2)	C19—C22—H22C	109.5
C5—C6—H6	119.4	H22A—C22—H22C	109.5
C7—C6—H6	119.4	H22B—C22—H22C	109.5
C6—C7—C2	118.3 (2)	C24—C23—H23A	109.5
C6—C7—S1	120.62 (18)	C24—C23—H23B	109.5
C2—C7—S1	121.05 (17)	H23A—C23—H23B	109.5
C9—C8—S1	107.30 (17)	C24—C23—H23C	109.5
C9—C8—H8A	110.3	H23A—C23—H23C	109.5
S1—C8—H8A	110.3	H23B—C23—H23C	109.5
C9—C8—H8B	110.3	C23—C24—C25	116.3 (4)
S1—C8—H8B	110.3	C23—C24—H24A	108.2
H8A—C8—H8B	108.5	C25—C24—H24A	108.2
C10—C9—C14	118.3 (2)	C23—C24—H24B	108.2
C10—C9—C8	121.6 (2)	C25—C24—H24B	108.2
C14—C9—C8	120.1 (2)	H24A—C24—H24B	107.4
C9—C10—C11	122.7 (2)	C26—C25—C24	112.1 (3)

C9—C10—H10	118.6	C26—C25—H25A	109.2
C11—C10—H10	118.6	C24—C25—H25A	109.2
C10—C11—C12	117.1 (2)	C26—C25—H25B	109.2
C10—C11—C19	120.4 (2)	C24—C25—H25B	109.2
C12—C11—C19	122.5 (2)	H25A—C25—H25B	107.9
O3—C12—C11	118.8 (2)	N1—C26—C25	112.2 (2)
O3—C12—C13	118.9 (2)	N1—C26—H26A	109.2
C11—C12—C13	122.2 (2)	C25—C26—H26A	109.2
C14—C13—C12	117.0 (2)	N1—C26—H26B	109.2
C14—C13—C15	120.0 (2)	C25—C26—H26B	109.2
C12—C13—C15	122.9 (2)	H26A—C26—H26B	107.9
C9—C14—C13	122.7 (2)	N1—C27—C28	111.4 (2)
C9—C14—H14	118.7	N1—C27—H27A	109.3
C13—C14—H14	118.7	C28—C27—H27A	109.3
C17—C15—C18	110.2 (3)	N1—C27—H27B	109.3
C17—C15—C13	109.0 (3)	C28—C27—H27B	109.3
C18—C15—C13	112.2 (2)	H27A—C27—H27B	108.0
C17—C15—C16	107.4 (3)	C29—C28—C27	113.0 (3)
C18—C15—C16	106.9 (3)	C29—C28—H28A	109.0
C13—C15—C16	111.0 (2)	C27—C28—H28A	109.0
C15—C16—H16A	109.5	C29—C28—H28B	109.0
C15—C16—H16B	109.5	C27—C28—H28B	109.0
H16A—C16—H16B	109.5	H28A—C28—H28B	107.8
C15—C16—H16C	109.5	C28—C29—C30	113.4 (3)
H16A—C16—H16C	109.5	C28—C29—H29A	108.9
H16B—C16—H16C	109.5	C30—C29—H29A	108.9
C15—C17—H17A	109.5	C28—C29—H29B	108.9
C15—C17—H17B	109.5	C30—C29—H29B	108.9
H17A—C17—H17B	109.5	H29A—C29—H29B	107.7
C15—C17—H17C	109.5	C29—C30—H30A	109.5
H17A—C17—H17C	109.5	C29—C30—H30B	109.5
H17B—C17—H17C	109.5	H30A—C30—H30B	109.5
C15—C18—H18A	109.5	C29—C30—H30C	109.5
C15—C18—H18B	109.5	H30A—C30—H30C	109.5
H18A—C18—H18B	109.5	H30B—C30—H30C	109.5
C15—C18—H18C	109.5		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O3—H3···O1 ⁱ	0.81 (2)	2.02 (2)	2.723 (3)	146 (3)
N1—H1A···O2	0.89 (2)	1.88 (2)	2.767 (3)	177 (3)
N1—H1B···O1 ⁱⁱ	0.89 (2)	2.00 (2)	2.806 (3)	150 (3)
N1—H1B···O2 ⁱⁱ	0.89 (2)	2.36 (2)	3.167 (3)	150 (2)
N1—H1B···O2 ⁱⁱ	0.89 (2)	2.36 (2)	3.167 (3)	150 (2)

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+1, -y, -z+1$.