

Bromido(quinolin-8-ol- $\kappa^2 N,O$)(quinolin-8-olato- $\kappa^2 N,O$)zinc(II) methanol monosolvate

Ezzatollah Najafi,^a Mostafa M. Amini^a and Seik Weng Ng^{b*}

^aDepartment of Chemistry, General Campus, Shahid Beheshti University, Tehran 1983963113, Iran, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 12 September 2010; accepted 14 September 2010

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.032; wR factor = 0.103; data-to-parameter ratio = 17.0.

The title compound, $[\text{ZnBr}(\text{C}_9\text{H}_6\text{NO})(\text{C}_9\text{H}_7\text{NO})]\cdot\text{CH}_3\text{OH}$, has its metal atom N,O -chelated by a neutral and a deprotonated 8-hydroxyquinoline ligand. The hydroxy unit of the neutral ligand is a hydrogen-bond donor to the methanol O atom and the alkoxy O atom of the monoanionic ligand is a hydrogen-bond acceptor to the methanol O atom. In the crystal, adjacent molecules are linked by these two hydrogen bonds, generating a chain running along the a axis.

Related literature

For a related structure, see: Najafi *et al.* (2010).



Experimental

Crystal data

$[\text{ZnBr}(\text{C}_9\text{H}_6\text{NO})(\text{C}_9\text{H}_7\text{NO})]\cdot\text{CH}_3\text{O}$	$\gamma = 109.470 (1)^\circ$
$M_r = 466.63$	$V = 884.81 (12)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 8.4485 (7)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 8.6968 (7)\text{ \AA}$	$\mu = 3.67\text{ mm}^{-1}$
$c = 13.1868 (10)\text{ \AA}$	$T = 100\text{ K}$
$\alpha = 97.241 (1)^\circ$	$0.30 \times 0.20 \times 0.10\text{ mm}$
$\beta = 99.209 (1)^\circ$	

Data collection

Bruker SMART APEX	8392 measured reflections
diffractometer	4022 independent reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	3354 reflections with $I > 2\sigma(I)$
$R_{\text{int}} = 0.028$	
$T_{\min} = 0.406$, $T_{\max} = 0.711$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$	237 parameters
$wR(F^2) = 0.103$	H-atom parameters constrained
$S = 1.10$	$\Delta\rho_{\max} = 0.78\text{ e \AA}^{-3}$
4022 reflections	$\Delta\rho_{\min} = -0.85\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$O2-\text{H}2\cdots O3^i$	0.84	1.90	2.585 (4)	137
$O3-\text{H}3\cdots O1$	0.84	1.71	2.551 (4)	178

Symmetry code: (i) $x - 1, y, z$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank Shahid Beheshti University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5356).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Najafi, E., Amini, M. M. & Ng, S. W. (2010). *Acta Cryst. E66*, m1276.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2010). E66, m1278 [doi:10.1107/S160053681003672X]

Bromido(quinolin-8-ol- κ^2N,O)(quinolin-8-olato- κ^2N,O)zinc(II) methanol monosolvate

Ezzatollah Najafi, Mostafa M. Amini and Seik Weng Ng

S1. Comment


An earlier study reported $C_{10}H_{10}NO^+ZnBr_2(C_{10}H_8NO)CH_3OH$, which feature cations, tetrahedral anions and solvent molecules linked by N···O, O···O and O···Br hydrogen bonds into a linear chain. The salt was synthesized by reacting zinc bromide and 2-methyl-8-hydroxyquinoline in methanol; no base was added (Najafi *et al.*, 2010). The present study uses 8-hydroxyquinoline instead of 2-methyl-8-hydroxyquinoline as the organic reactant. The product is a mono-solvated neutral molecule (Scheme I, Fig. 1). The methanol-solvated compound, $ZnBr(C_9H_6NO)(C_9H_7NO)CH_3OH$, has its metal atom *N,O*-chelated by a neutral and deprotonated 8-hydroxyquinoline ligand. The hydroxy unit of the neutral ligand is hydrogen-bond donor methanol O atom and the alkoxy O atom of the monoanionic ligand is hydrogen-bond acceptor to methanol O atom. Adjacent molecules are linked by these two hydrogen bonds to generate a linear chain running along the *a*-axis of the triclinic unit cell (Fig. 2).

S2. Experimental


Zinc bromide (0.19 g, 0.75 mmol) and 8-hydroxyquinoline (0.22 g, 1.5 mmol) were loaded into a convection tube; the tube was filled with dry methanol and kept at 333 K. Crystals were collected from the side arm after several days.

S3. Refinement

Hydrogen atoms were placed in calculated positions (C–H 0.95–0.98, O–H 0.84 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to 1.2–1.5 $U_{eq}(C,O)$.

**Figure 1**

Anisotropic displacement ellipsoid plot (Barbour, 2001) of the title compound at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

**Figure 2**

Hydrogen bonded chain structure.

Bromido(quinolin-8-ol- κ^2 N,O)(quinolin-8-olato- κ^2 N,O)zinc(II) methanol monosolvate

Crystal data

$[\text{ZnBr}(\text{C}_9\text{H}_6\text{NO})(\text{C}_9\text{H}_7\text{NO})]\cdot\text{CH}_4\text{O}$
 $M_r = 466.63$

Triclinic, $P\bar{1}$
Hall symbol: -P 1

$a = 8.4485 (7)$ Å
 $b = 8.6968 (7)$ Å
 $c = 13.1868 (10)$ Å
 $\alpha = 97.241 (1)^\circ$
 $\beta = 99.209 (1)^\circ$
 $\gamma = 109.470 (1)^\circ$
 $V = 884.81 (12)$ Å³
 $Z = 2$
 $F(000) = 468$

$D_x = 1.751$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 3867 reflections
 $\theta = 2.5\text{--}28.2^\circ$
 $\mu = 3.67$ mm⁻¹
 $T = 100$ K
Prism, yellow
 $0.30 \times 0.20 \times 0.10$ mm

Data collection

Bruker SMART APEX
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.406$, $T_{\max} = 0.711$

8392 measured reflections
4022 independent reflections
3354 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.5^\circ$
 $h = -10 \rightarrow 10$
 $k = -11 \rightarrow 11$
 $l = -15 \rightarrow 17$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.103$
 $S = 1.10$
4022 reflections
237 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0457P)^2 + 1.9422P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.78$ e Å⁻³
 $\Delta\rho_{\min} = -0.85$ e Å⁻³

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.45338 (5)	0.64466 (5)	0.71737 (3)	0.01184 (12)
Br1	0.58054 (4)	0.92415 (4)	0.81357 (3)	0.01654 (11)
O1	0.6864 (3)	0.6465 (3)	0.64187 (19)	0.0152 (5)
O2	0.2092 (3)	0.5306 (3)	0.73396 (19)	0.0145 (5)
H2	0.1230	0.5461	0.7024	0.022*
O3	0.9908 (3)	0.6800 (3)	0.7299 (2)	0.0209 (6)
H3	0.8908	0.6707	0.7016	0.031*
N1	0.3868 (4)	0.6687 (4)	0.5653 (2)	0.0119 (6)
N2	0.5025 (4)	0.4869 (3)	0.8093 (2)	0.0121 (6)
C1	0.5180 (4)	0.7280 (4)	0.5148 (3)	0.0110 (6)
C2	0.6796 (4)	0.7212 (4)	0.5573 (3)	0.0128 (7)
C3	0.8163 (4)	0.7842 (4)	0.5120 (3)	0.0150 (7)
H3A	0.9255	0.7819	0.5415	0.018*
C4	0.7946 (5)	0.8531 (5)	0.4209 (3)	0.0177 (7)
H4	0.8897	0.8950	0.3892	0.021*
C5	0.6384 (4)	0.8603 (4)	0.3777 (3)	0.0151 (7)
H5	0.6264	0.9080	0.3172	0.018*

C6	0.4957 (5)	0.7963 (4)	0.4238 (3)	0.0141 (7)
C7	0.3290 (5)	0.7915 (4)	0.3821 (3)	0.0153 (7)
H7	0.3085	0.8350	0.3208	0.018*
C8	0.1969 (5)	0.7238 (5)	0.4304 (3)	0.0173 (7)
H8	0.0835	0.7166	0.4020	0.021*
C9	0.2328 (4)	0.6652 (4)	0.5232 (3)	0.0156 (7)
H9	0.1412	0.6208	0.5571	0.019*
C10	0.3596 (4)	0.4025 (4)	0.8429 (3)	0.0109 (6)
C11	0.2035 (4)	0.4283 (4)	0.8012 (3)	0.0138 (7)
C12	0.0569 (4)	0.3427 (4)	0.8339 (3)	0.0150 (7)
H12	-0.0481	0.3560	0.8074	0.018*
C13	0.0599 (5)	0.2365 (5)	0.9055 (3)	0.0165 (7)
H13	-0.0433	0.1800	0.9263	0.020*
C14	0.2083 (5)	0.2119 (4)	0.9464 (3)	0.0165 (7)
H14	0.2079	0.1400	0.9951	0.020*
C15	0.3606 (4)	0.2956 (4)	0.9146 (3)	0.0128 (7)
C16	0.5206 (5)	0.2803 (4)	0.9519 (3)	0.0145 (7)
H16	0.5282	0.2085	0.9996	0.017*
C17	0.6636 (5)	0.3682 (4)	0.9194 (3)	0.0153 (7)
H17	0.7716	0.3604	0.9457	0.018*
C18	0.6497 (4)	0.4706 (4)	0.8467 (3)	0.0134 (7)
H18	0.7494	0.5301	0.8235	0.016*
C19	1.0704 (5)	0.8335 (5)	0.8036 (3)	0.0271 (9)
H19A	1.0933	0.8107	0.8743	0.041*
H19B	0.9936	0.8967	0.8004	0.041*
H19C	1.1789	0.8984	0.7867	0.041*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0117 (2)	0.0145 (2)	0.0111 (2)	0.00508 (15)	0.00377 (15)	0.00635 (15)
Br1	0.0208 (2)	0.01368 (17)	0.01481 (19)	0.00510 (14)	0.00314 (14)	0.00568 (13)
O1	0.0162 (12)	0.0189 (12)	0.0138 (12)	0.0078 (10)	0.0056 (10)	0.0078 (10)
O2	0.0105 (11)	0.0194 (12)	0.0163 (13)	0.0080 (10)	0.0012 (9)	0.0076 (10)
O3	0.0130 (12)	0.0228 (13)	0.0256 (15)	0.0072 (11)	-0.0002 (11)	0.0035 (11)
N1	0.0106 (13)	0.0158 (14)	0.0101 (14)	0.0054 (11)	0.0026 (11)	0.0030 (11)
N2	0.0159 (14)	0.0129 (13)	0.0080 (13)	0.0037 (11)	0.0045 (11)	0.0049 (11)
C1	0.0104 (15)	0.0115 (15)	0.0109 (16)	0.0041 (12)	0.0033 (12)	-0.0001 (12)
C2	0.0159 (17)	0.0138 (15)	0.0117 (16)	0.0083 (13)	0.0044 (13)	0.0042 (13)
C3	0.0115 (16)	0.0187 (17)	0.0171 (18)	0.0080 (13)	0.0034 (13)	0.0043 (14)
C4	0.0172 (18)	0.0193 (17)	0.0155 (18)	0.0026 (14)	0.0076 (14)	0.0052 (14)
C5	0.0169 (17)	0.0161 (16)	0.0107 (16)	0.0044 (14)	0.0015 (13)	0.0030 (13)
C6	0.0185 (17)	0.0126 (15)	0.0114 (16)	0.0046 (13)	0.0061 (14)	0.0027 (13)
C7	0.0180 (17)	0.0180 (17)	0.0099 (16)	0.0072 (14)	-0.0003 (13)	0.0043 (13)
C8	0.0159 (17)	0.0236 (18)	0.0126 (17)	0.0076 (14)	0.0010 (14)	0.0061 (14)
C9	0.0120 (16)	0.0209 (17)	0.0161 (18)	0.0077 (14)	0.0045 (14)	0.0045 (14)
C10	0.0124 (15)	0.0120 (15)	0.0089 (15)	0.0056 (12)	0.0025 (12)	0.0011 (12)
C11	0.0145 (16)	0.0158 (16)	0.0118 (16)	0.0062 (13)	0.0025 (13)	0.0034 (13)

C12	0.0108 (16)	0.0208 (17)	0.0123 (17)	0.0054 (13)	0.0003 (13)	0.0038 (13)
C13	0.0146 (17)	0.0199 (17)	0.0152 (18)	0.0041 (14)	0.0063 (14)	0.0062 (14)
C14	0.0206 (18)	0.0153 (16)	0.0152 (18)	0.0056 (14)	0.0069 (14)	0.0066 (14)
C15	0.0151 (16)	0.0123 (15)	0.0082 (16)	0.0029 (13)	0.0004 (13)	-0.0002 (12)
C16	0.0197 (18)	0.0145 (16)	0.0127 (17)	0.0105 (14)	0.0021 (14)	0.0048 (13)
C17	0.0143 (17)	0.0208 (17)	0.0127 (17)	0.0093 (14)	0.0009 (13)	0.0043 (14)
C18	0.0121 (16)	0.0171 (16)	0.0143 (17)	0.0081 (13)	0.0049 (13)	0.0045 (13)
C19	0.0193 (19)	0.027 (2)	0.029 (2)	0.0051 (16)	0.0007 (17)	-0.0014 (17)

Geometric parameters (\AA , $\text{^{\circ}}$)

Zn1—O2	2.030 (2)	C7—C8	1.372 (5)
Zn1—N2	2.040 (3)	C7—H7	0.9500
Zn1—N1	2.050 (3)	C8—C9	1.411 (5)
Zn1—O1	2.340 (2)	C8—H8	0.9500
Zn1—Br1	2.3911 (5)	C9—H9	0.9500
O1—C2	1.362 (4)	C10—C15	1.409 (5)
O2—C11	1.328 (4)	C10—C11	1.445 (5)
O2—H2	0.8400	C11—C12	1.383 (5)
O3—C19	1.430 (5)	C12—C13	1.404 (5)
O3—H3	0.8400	C12—H12	0.9500
N1—C9	1.319 (4)	C13—C14	1.378 (5)
N1—C1	1.372 (4)	C13—H13	0.9500
N2—C18	1.322 (4)	C14—C15	1.410 (5)
N2—C10	1.361 (4)	C14—H14	0.9500
C1—C2	1.414 (5)	C15—C16	1.417 (5)
C1—C6	1.418 (5)	C16—C17	1.364 (5)
C2—C3	1.368 (5)	C16—H16	0.9500
C3—C4	1.422 (5)	C17—C18	1.406 (5)
C3—H3A	0.9500	C17—H17	0.9500
C4—C5	1.376 (5)	C18—H18	0.9500
C4—H4	0.9500	C19—H19A	0.9800
C5—C6	1.416 (5)	C19—H19B	0.9800
C5—H5	0.9500	C19—H19C	0.9800
C6—C7	1.412 (5)		
O2—Zn1—N2	82.51 (11)	C6—C7—H7	120.1
O2—Zn1—N1	95.88 (11)	C7—C8—C9	118.8 (3)
N2—Zn1—N1	143.75 (11)	C7—C8—H8	120.6
O2—Zn1—O1	150.98 (10)	C9—C8—H8	120.6
N2—Zn1—O1	90.06 (10)	N1—C9—C8	123.4 (3)
N1—Zn1—O1	73.91 (10)	N1—C9—H9	118.3
O2—Zn1—Br1	112.41 (7)	C8—C9—H9	118.3
N2—Zn1—Br1	109.66 (8)	N2—C10—C15	122.9 (3)
N1—Zn1—Br1	104.40 (8)	N2—C10—C11	116.5 (3)
O1—Zn1—Br1	96.52 (6)	C15—C10—C11	120.7 (3)
C2—O1—Zn1	108.4 (2)	O2—C11—C12	124.4 (3)
C11—O2—Zn1	111.3 (2)	O2—C11—C10	118.4 (3)

C11—O2—H2	124.4	C12—C11—C10	117.2 (3)
Zn1—O2—H2	124.4	C11—C12—C13	121.4 (3)
C19—O3—H3	109.5	C11—C12—H12	119.3
C9—N1—C1	118.4 (3)	C13—C12—H12	119.3
C9—N1—Zn1	122.8 (2)	C14—C13—C12	121.9 (3)
C1—N1—Zn1	117.2 (2)	C14—C13—H13	119.0
C18—N2—C10	119.0 (3)	C12—C13—H13	119.0
C18—N2—Zn1	129.8 (2)	C13—C14—C15	118.6 (3)
C10—N2—Zn1	110.8 (2)	C13—C14—H14	120.7
N1—C1—C2	117.6 (3)	C15—C14—H14	120.7
N1—C1—C6	122.0 (3)	C10—C15—C14	120.1 (3)
C2—C1—C6	120.3 (3)	C10—C15—C16	116.4 (3)
O1—C2—C3	124.0 (3)	C14—C15—C16	123.5 (3)
O1—C2—C1	116.0 (3)	C17—C16—C15	120.2 (3)
C3—C2—C1	119.9 (3)	C17—C16—H16	119.9
C2—C3—C4	119.9 (3)	C15—C16—H16	119.9
C2—C3—H3A	120.0	C16—C17—C18	119.3 (3)
C4—C3—H3A	120.0	C16—C17—H17	120.4
C5—C4—C3	121.2 (3)	C18—C17—H17	120.4
C5—C4—H4	119.4	N2—C18—C17	122.2 (3)
C3—C4—H4	119.4	N2—C18—H18	118.9
C4—C5—C6	119.7 (3)	C17—C18—H18	118.9
C4—C5—H5	120.2	O3—C19—H19A	109.5
C6—C5—H5	120.2	O3—C19—H19B	109.5
C7—C6—C5	123.5 (3)	H19A—C19—H19B	109.5
C7—C6—C1	117.5 (3)	O3—C19—H19C	109.5
C5—C6—C1	118.9 (3)	H19A—C19—H19C	109.5
C8—C7—C6	119.8 (3)	H19B—C19—H19C	109.5
C8—C7—H7	120.1		
O2—Zn1—O1—C2	94.6 (3)	C4—C5—C6—C7	-177.0 (3)
N2—Zn1—O1—C2	169.1 (2)	C4—C5—C6—C1	0.9 (5)
N1—Zn1—O1—C2	22.0 (2)	N1—C1—C6—C7	-4.1 (5)
Br1—Zn1—O1—C2	-81.1 (2)	C2—C1—C6—C7	176.7 (3)
N2—Zn1—O2—C11	6.3 (2)	N1—C1—C6—C5	177.9 (3)
N1—Zn1—O2—C11	149.9 (2)	C2—C1—C6—C5	-1.3 (5)
O1—Zn1—O2—C11	82.7 (3)	C5—C6—C7—C8	178.5 (3)
Br1—Zn1—O2—C11	-101.9 (2)	C1—C6—C7—C8	0.6 (5)
O2—Zn1—N1—C9	21.6 (3)	C6—C7—C8—C9	2.0 (5)
N2—Zn1—N1—C9	106.9 (3)	C1—N1—C9—C8	-1.9 (5)
O1—Zn1—N1—C9	173.8 (3)	Zn1—N1—C9—C8	162.7 (3)
Br1—Zn1—N1—C9	-93.4 (3)	C7—C8—C9—N1	-1.4 (6)
O2—Zn1—N1—C1	-173.6 (2)	C18—N2—C10—C15	-1.2 (5)
N2—Zn1—N1—C1	-88.3 (3)	Zn1—N2—C10—C15	-174.5 (3)
O1—Zn1—N1—C1	-21.3 (2)	C18—N2—C10—C11	178.6 (3)
Br1—Zn1—N1—C1	71.4 (2)	Zn1—N2—C10—C11	5.2 (4)
O2—Zn1—N2—C18	-178.6 (3)	Zn1—O2—C11—C12	175.2 (3)
N1—Zn1—N2—C18	91.6 (3)	Zn1—O2—C11—C10	-5.4 (4)

O1—Zn1—N2—C18	29.5 (3)	N2—C10—C11—O2	0.1 (5)
Br1—Zn1—N2—C18	-67.5 (3)	C15—C10—C11—O2	179.8 (3)
O2—Zn1—N2—C10	-6.2 (2)	N2—C10—C11—C12	179.6 (3)
N1—Zn1—N2—C10	-96.0 (3)	C15—C10—C11—C12	-0.7 (5)
O1—Zn1—N2—C10	-158.1 (2)	O2—C11—C12—C13	-179.9 (3)
Br1—Zn1—N2—C10	104.9 (2)	C10—C11—C12—C13	0.7 (5)
C9—N1—C1—C2	-176.1 (3)	C11—C12—C13—C14	-0.2 (6)
Zn1—N1—C1—C2	18.4 (4)	C12—C13—C14—C15	-0.3 (5)
C9—N1—C1—C6	4.7 (5)	N2—C10—C15—C14	179.9 (3)
Zn1—N1—C1—C6	-160.8 (3)	C11—C10—C15—C14	0.2 (5)
Zn1—O1—C2—C3	161.4 (3)	N2—C10—C15—C16	0.5 (5)
Zn1—O1—C2—C1	-20.0 (3)	C11—C10—C15—C16	-179.2 (3)
N1—C1—C2—O1	3.7 (4)	C13—C14—C15—C10	0.3 (5)
C6—C1—C2—O1	-177.1 (3)	C13—C14—C15—C16	179.7 (3)
N1—C1—C2—C3	-177.7 (3)	C10—C15—C16—C17	1.0 (5)
C6—C1—C2—C3	1.6 (5)	C14—C15—C16—C17	-178.5 (3)
O1—C2—C3—C4	177.1 (3)	C15—C16—C17—C18	-1.7 (5)
C1—C2—C3—C4	-1.4 (5)	C10—N2—C18—C17	0.4 (5)
C2—C3—C4—C5	1.0 (6)	Zn1—N2—C18—C17	172.3 (3)
C3—C4—C5—C6	-0.8 (5)	C16—C17—C18—N2	1.0 (5)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O2—H2···O3 ⁱ	0.84	1.90	2.585 (4)	137
O3—H3···O1	0.84	1.71	2.551 (4)	178

Symmetry code: (i) $x-1, y, z$.