

**Bis(2-aminopyrimidine- $\kappa N^1$ )aqua(nitrato- $\kappa O$ )(nitrato- $\kappa^2 O,O'$ )zinc(II)**Shan Gao<sup>a</sup> and Seik Weng Ng<sup>b\*</sup>

<sup>a</sup>College of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 30 August 2010; accepted 14 September 2010

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$ ;  $R$  factor = 0.030;  $wR$  factor = 0.081; data-to-parameter ratio = 14.1.

The water-coordinated Zn atom in the title monoqua zinc nitrate adduct of 2-aminopyrimidine,  $[\text{Zn}(\text{NO}_3)_2(\text{C}_4\text{H}_5\text{N}_3)_2(\text{H}_2\text{O})]$ , is bonded to a monodentate nitrate ion and is chelated by the other nitrate ion. The heterocyclic ligands coordinate through ring *N*-donor sites. The coordination geometry about the Zn(II) atom is a distorted octahedron. Intramolecular N—H···O hydrogen bonds occur. In the crystal, adjacent adduct molecules are linked by O—H···O, O—H···N and N—H···O hydrogen bonds into a layer motif parallel to (001).

**Related literature**

The aquazinc nitrate adduct is isotypic with its Co and Ni analogs, see: Pike *et al.* (2006). The copper nitrate adduct is anhydrous, see: Albada *et al.* (2002).

**Experimental***Crystal data* $M_r = 397.63$ Monoclinic,  $C2/c$  $a = 13.2742(4)\text{ \AA}$  $b = 8.0142(2)\text{ \AA}$  $c = 28.6204(7)\text{ \AA}$  $\beta = 101.335(1)^\circ$  $V = 2985.31(14)\text{ \AA}^3$ 

$Z = 8$   
Mo  $K\alpha$  radiation  
 $\mu = 1.70\text{ mm}^{-1}$

$T = 293\text{ K}$   
 $0.22 \times 0.18 \times 0.12\text{ mm}$

*Data collection*

Rigaku R-AXIS RAPID diffractometer  
Absorption correction: multi-scan (*ABSCOR*; Higashi, 1995)  
 $T_{\min} = 0.706$ ,  $T_{\max} = 0.822$

14113 measured reflections  
3401 independent reflections  
3006 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.039$

*Refinement*

$R[F^2 > 2\sigma(F^2)] = 0.030$   
 $wR(F^2) = 0.081$   
 $S = 1.04$   
3401 reflections  
241 parameters  
6 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.40\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.43\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1w—H11···O2 <sup>i</sup>	0.83 (1)	1.99 (2)	2.776 (2)	158 (3)
O1w—H12···N2 <sup>ii</sup>	0.84 (1)	1.94 (1)	2.754 (2)	165 (3)
N3—H31···O1	0.87 (1)	2.23 (2)	2.989 (3)	146 (2)
N3—H32···O5 <sup>iii</sup>	0.86 (1)	2.34 (2)	3.133 (3)	152 (3)
N6—H61···O1	0.87 (1)	2.37 (3)	3.010 (2)	131 (3)
N6—H61···O5	0.87 (1)	2.43 (2)	3.122 (3)	137 (3)
N6—H62···O1 <sup>iv</sup>	0.87 (1)	2.41 (2)	3.192 (2)	150 (3)
N6—H62···O3 <sup>iv</sup>	0.87 (1)	2.45 (2)	3.265 (3)	156 (3)

Symmetry codes: (i)  $x + \frac{1}{2}, y + \frac{1}{2}, z$ ; (ii)  $x + \frac{1}{2}, y - \frac{1}{2}, z$ ; (iii)  $x - \frac{1}{2}, y + \frac{1}{2}, z$ ; (iv)  $-x + 1, y, -z + \frac{3}{2}$ .

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSC and Rigaku, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank the Key Project of the Natural Science Foundation of Heilongjiang Province (No. ZD200903), the Innovation Team of the Education Bureau of Heilongjiang Province (No. 2010t d03), Heilongjiang University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2204).

**References**

- Albada, G. A., Mutikainen, I., Turpeinen, U. & Reedijk, J. (2002). *Acta Cryst. E58*, m55–m57.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Pike, R. D., Lim, M. J., Wilcox, E. A. L. & Tronic, T. A. (2006). *J. Chem. Crystallogr.* **11**, 781–791.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC and Rigaku (2002). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

# supporting information

*Acta Cryst.* (2010). E66, m1279 [doi:10.1107/S1600536810036731]

## **Bis(2-aminopyrimidine- $\kappa N^1$ )aqua(nitrato- $\kappa O$ )(nitrato- $\kappa^2 O,O'$ )zinc(II)**

**Shan Gao and Seik Weng Ng**

### **S1. Comment**

The cobalt, nickel and copper adducts of 2-aminopyrimidine have been reported; the first two are monoaqua complexes (Pike *et al.*, 2006) whereas the copper complex is anhydrous (Albada *et al.*, 2002). In the aqua complexes, one nitrate is monodentate and the other is chelating; the heterocyclic ligand coordinates through a ring donor site. The present zinc analog (Scheme I, Fig. 1) is isostructural to the cobalt and nickel adducts, whose structures have been described in detail. Adjacent molecules are linked by O—H $\cdots$ O and N—H $\cdots$ O hydrogen bonds into a layer motif (Fig. 2).


### **S2. Experimental**

Zinc nitrate (1 mmol) and 2-aminopyrimidine (1 mmol) were dissolved in a small volume of water to give a colorless solution. Colorless prismatic crystals separated from the solution after a few days.


### **S3. Refinement**

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 Å) and were included in the refinement in the riding model approximation, with  $U(H)$  set to 1.2 $U(C)$ .

The amino and water H-atoms were located in a difference Fourier map, and were refined with a distance restraint of N—H 0.88 $\pm$ 0.01 and O—H 0.84 $\pm$ 0.01 Å; their temperature factors were freely refined.

**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of  $\text{Zn}(\text{H}_2\text{O})(\text{NO}_3)_2(\text{C}_4\text{H}_5\text{N}_3)_2$  at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

**Figure 2**

Hydrogen-bonded layer structure.

**Bis(2-aminopyrimidine- $\kappa N^1$ )aqua(nitrato- $\kappa O$ )(nitrato- $\kappa^2 O, O'$ )zinc(II)***Crystal data* $[Zn(NO_3)_2(C_4H_5N_3)_2(H_2O)]$  $M_r = 397.63$ Monoclinic,  $C2/c$ 

Hall symbol: -C 2yc

 $a = 13.2742 (4) \text{ \AA}$  $b = 8.0142 (2) \text{ \AA}$  $c = 28.6204 (7) \text{ \AA}$  $\beta = 101.335 (1)^\circ$  $V = 2985.31 (14) \text{ \AA}^3$  $Z = 8$  $F(000) = 1616$  $D_x = 1.769 \text{ Mg m}^{-3}$ Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$ 

Cell parameters from 11713 reflections

 $\theta = 3.0\text{--}27.4^\circ$  $\mu = 1.70 \text{ mm}^{-1}$  $T = 293 \text{ K}$ 

Prism, colorless

 $0.22 \times 0.18 \times 0.12 \text{ mm}$ *Data collection*

Rigaku R-AXIS RAPID

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 10.000 pixels  $\text{mm}^{-1}$  $\omega$  scans

Absorption correction: multi-scan

(ABSCOR; Higashi, 1995)

 $T_{\min} = 0.706$ ,  $T_{\max} = 0.822$ 

14113 measured reflections

3401 independent reflections

3006 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.039$  $\theta_{\max} = 27.4^\circ$ ,  $\theta_{\min} = 3.0^\circ$  $h = -17 \rightarrow 17$  $k = -10 \rightarrow 10$  $l = -34 \rightarrow 37$ *Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.030$  $wR(F^2) = 0.081$  $S = 1.04$ 

3401 reflections

241 parameters

6 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0479P)^2 + 2.0445P]$   
where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.001$  $\Delta\rho_{\max} = 0.40 \text{ e \AA}^{-3}$  $\Delta\rho_{\min} = -0.43 \text{ e \AA}^{-3}$ *Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.498530 (14)	0.66515 (3)	0.616791 (7)	0.02883 (9)
O1	0.39847 (10)	0.55960 (19)	0.66269 (5)	0.0379 (3)
O2	0.28031 (17)	0.4531 (3)	0.60891 (6)	0.0822 (7)
O3	0.30770 (17)	0.3540 (3)	0.67905 (8)	0.0742 (6)
O4	0.46872 (15)	0.4598 (2)	0.55893 (9)	0.0764 (7)
O5	0.56026 (15)	0.4082 (3)	0.62668 (7)	0.0614 (5)
O6	0.5176 (2)	0.2078 (2)	0.57552 (7)	0.0745 (6)
O1W	0.62147 (10)	0.7246 (2)	0.58475 (5)	0.0377 (3)
N1	0.39392 (11)	0.8157 (2)	0.57361 (5)	0.0301 (3)
N2	0.24514 (12)	0.9882 (2)	0.55462 (6)	0.0398 (4)
N3	0.29718 (15)	0.8839 (3)	0.63018 (7)	0.0492 (5)
N4	0.55915 (12)	0.8053 (2)	0.67664 (5)	0.0306 (3)

N5	0.64072 (15)	0.8410 (2)	0.75816 (6)	0.0437 (4)
N6	0.60734 (16)	0.5800 (2)	0.72650 (7)	0.0460 (4)
N7	0.32868 (12)	0.4509 (2)	0.64962 (6)	0.0371 (4)
N8	0.51533 (15)	0.3535 (2)	0.58614 (8)	0.0466 (4)
C1	0.40689 (15)	0.8333 (3)	0.52831 (7)	0.0351 (4)
H1	0.4624	0.7805	0.5192	0.042*
C2	0.34156 (16)	0.9258 (3)	0.49524 (7)	0.0415 (5)
H2	0.3512	0.9366	0.4641	0.050*
C3	0.26084 (16)	1.0021 (3)	0.51044 (7)	0.0418 (5)
H3	0.2154	1.0662	0.4888	0.050*
C4	0.31217 (13)	0.8944 (3)	0.58546 (7)	0.0329 (4)
C5	0.56075 (18)	0.9717 (3)	0.67104 (8)	0.0447 (5)
H5	0.5335	1.0169	0.6413	0.054*
C6	0.6010 (2)	1.0766 (3)	0.70754 (10)	0.0586 (6)
H6	0.6024	1.1916	0.7033	0.070*
C7	0.63959 (19)	1.0039 (3)	0.75104 (8)	0.0534 (6)
H7	0.6661	1.0729	0.7766	0.064*
C8	0.60164 (13)	0.7461 (3)	0.72035 (6)	0.0317 (4)
H11	0.6657 (17)	0.789 (3)	0.5994 (9)	0.061 (8)*
H12	0.653 (2)	0.640 (2)	0.5780 (11)	0.067 (9)*
H31	0.3355 (18)	0.817 (3)	0.6501 (8)	0.053 (8)*
H32	0.2393 (12)	0.920 (3)	0.6356 (10)	0.057 (8)*
H61	0.567 (2)	0.515 (3)	0.7066 (9)	0.074 (9)*
H62	0.624 (2)	0.548 (4)	0.7560 (5)	0.071 (9)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Zn1	0.02802 (13)	0.03286 (14)	0.02437 (13)	0.00355 (8)	0.00208 (8)	-0.00019 (8)
O1	0.0338 (7)	0.0452 (8)	0.0339 (7)	-0.0098 (6)	0.0044 (5)	-0.0037 (6)
O2	0.0895 (14)	0.1073 (18)	0.0386 (9)	-0.0538 (13)	-0.0145 (9)	0.0059 (10)
O3	0.0787 (14)	0.0756 (14)	0.0671 (13)	-0.0275 (10)	0.0110 (11)	0.0237 (10)
O4	0.0554 (10)	0.0468 (10)	0.1130 (18)	0.0005 (8)	-0.0180 (11)	0.0186 (11)
O5	0.0691 (11)	0.0716 (12)	0.0484 (10)	-0.0027 (9)	0.0233 (8)	-0.0121 (9)
O6	0.1327 (19)	0.0331 (9)	0.0651 (12)	0.0095 (10)	0.0380 (12)	-0.0043 (9)
O1W	0.0294 (7)	0.0403 (8)	0.0446 (8)	-0.0014 (6)	0.0102 (6)	-0.0080 (7)
N1	0.0258 (7)	0.0392 (9)	0.0245 (7)	0.0043 (6)	0.0032 (6)	0.0006 (6)
N2	0.0365 (8)	0.0476 (10)	0.0336 (8)	0.0150 (7)	0.0028 (7)	0.0005 (7)
N3	0.0460 (10)	0.0729 (13)	0.0310 (9)	0.0265 (10)	0.0133 (8)	0.0088 (9)
N4	0.0319 (7)	0.0340 (8)	0.0257 (7)	-0.0031 (6)	0.0051 (6)	0.0003 (6)
N5	0.0488 (10)	0.0522 (11)	0.0279 (8)	-0.0134 (8)	0.0022 (7)	-0.0063 (7)
N6	0.0556 (11)	0.0426 (10)	0.0327 (9)	-0.0036 (8)	-0.0089 (8)	0.0065 (8)
N7	0.0328 (8)	0.0439 (9)	0.0345 (8)	-0.0068 (7)	0.0066 (6)	0.0001 (7)
N8	0.0512 (11)	0.0366 (10)	0.0561 (12)	0.0037 (8)	0.0210 (9)	-0.0023 (8)
C1	0.0344 (9)	0.0444 (11)	0.0272 (9)	0.0067 (8)	0.0075 (7)	0.0022 (7)
C2	0.0451 (11)	0.0533 (13)	0.0256 (9)	0.0084 (9)	0.0056 (8)	0.0066 (9)
C3	0.0412 (10)	0.0475 (12)	0.0330 (10)	0.0103 (9)	-0.0018 (8)	0.0042 (9)
C4	0.0296 (9)	0.0401 (10)	0.0281 (9)	0.0053 (7)	0.0030 (7)	-0.0011 (7)

C5	0.0589 (13)	0.0350 (11)	0.0386 (11)	-0.0020 (9)	0.0055 (9)	0.0039 (9)
C6	0.0835 (18)	0.0343 (12)	0.0553 (14)	-0.0100 (11)	0.0072 (13)	-0.0061 (10)
C7	0.0634 (14)	0.0536 (14)	0.0417 (12)	-0.0178 (12)	0.0068 (10)	-0.0162 (11)
C8	0.0287 (8)	0.0404 (11)	0.0252 (8)	-0.0058 (7)	0.0035 (7)	-0.0001 (7)

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

Zn1—N1	2.0583 (15)	N3—H31	0.869 (10)
Zn1—N4	2.0740 (16)	N3—H32	0.864 (10)
Zn1—O1W	2.0782 (14)	N4—C5	1.343 (3)
Zn1—O5	2.214 (2)	N4—C8	1.353 (2)
Zn1—O1	2.2117 (14)	N5—C7	1.320 (3)
Zn1—O4	2.313 (2)	N5—C8	1.341 (2)
O1—N7	1.273 (2)	N6—C8	1.342 (3)
O2—N7	1.215 (2)	N6—H61	0.871 (10)
O3—N7	1.218 (3)	N6—H62	0.869 (10)
O4—N8	1.235 (3)	C1—C2	1.369 (3)
O5—N8	1.274 (3)	C1—H1	0.9300
O6—N8	1.209 (2)	C2—C3	1.376 (3)
O1W—H11	0.831 (10)	C2—H2	0.9300
O1W—H12	0.840 (10)	C3—H3	0.9300
N1—C1	1.348 (2)	C5—C6	1.366 (3)
N1—C4	1.355 (2)	C5—H5	0.9300
N2—C3	1.326 (3)	C6—C7	1.379 (4)
N2—C4	1.351 (2)	C6—H6	0.9300
N3—C4	1.336 (3)	C7—H7	0.9300
N1—Zn1—N4	106.55 (6)	C8—N6—H62	115 (2)
N1—Zn1—O1W	95.52 (6)	H61—N6—H62	118 (3)
N4—Zn1—O1W	91.68 (6)	O3—N7—O2	121.54 (19)
N1—Zn1—O5	144.20 (7)	O3—N7—O1	119.02 (18)
N4—Zn1—O5	108.93 (7)	O2—N7—O1	119.28 (18)
O1W—Zn1—O5	88.09 (6)	O6—N8—O4	122.8 (2)
N1—Zn1—O1	99.66 (6)	O6—N8—O5	122.1 (2)
N4—Zn1—O1	84.11 (6)	O4—N8—O5	115.1 (2)
O1W—Zn1—O1	164.82 (6)	N1—C1—C2	122.55 (18)
O5—Zn1—O1	79.56 (6)	N1—C1—H1	118.7
N1—Zn1—O4	89.25 (6)	C2—C1—H1	118.7
N4—Zn1—O4	163.88 (6)	C1—C2—C3	116.67 (19)
O1W—Zn1—O4	83.45 (7)	C1—C2—H2	121.7
O5—Zn1—O4	55.71 (7)	C3—C2—H2	121.7
O1—Zn1—O4	96.60 (7)	N2—C3—C2	122.81 (18)
N7—O1—Zn1	125.05 (12)	N2—C3—H3	118.6
N8—O4—Zn1	92.60 (16)	C2—C3—H3	118.6
N8—O5—Zn1	96.21 (14)	N3—C4—N2	117.25 (17)
Zn1—O1W—H11	117 (2)	N3—C4—N1	119.20 (17)
Zn1—O1W—H12	113 (2)	N2—C4—N1	123.53 (17)
H11—O1W—H12	106 (3)	N4—C5—C6	122.2 (2)

C1—N1—C4	116.87 (16)	N4—C5—H5	118.9
C1—N1—Zn1	116.11 (12)	C6—C5—H5	118.9
C4—N1—Zn1	126.99 (13)	C5—C6—C7	116.8 (2)
C3—N2—C4	117.57 (17)	C5—C6—H6	121.6
C4—N3—H31	119.3 (18)	C7—C6—H6	121.6
C4—N3—H32	117.1 (19)	N5—C7—C6	123.2 (2)
H31—N3—H32	121 (3)	N5—C7—H7	118.4
C5—N4—C8	116.43 (17)	C6—C7—H7	118.4
C5—N4—Zn1	116.84 (13)	N6—C8—N5	117.00 (17)
C8—N4—Zn1	126.66 (13)	N6—C8—N4	118.12 (17)
C7—N5—C8	116.47 (19)	N5—C8—N4	124.86 (19)
C8—N6—H61	120 (2)		
N1—Zn1—O1—N7	72.86 (16)	O5—Zn1—N4—C8	-19.60 (17)
N4—Zn1—O1—N7	178.70 (16)	O1—Zn1—N4—C8	57.21 (15)
O1W—Zn1—O1—N7	-106.8 (2)	O4—Zn1—N4—C8	-36.2 (3)
O5—Zn1—O1—N7	-70.77 (15)	Zn1—O1—N7—O3	150.33 (18)
O4—Zn1—O1—N7	-17.50 (16)	Zn1—O1—N7—O2	-34.3 (3)
N1—Zn1—O4—N8	-168.46 (15)	Zn1—O4—N8—O6	173.3 (2)
N4—Zn1—O4—N8	22.8 (4)	Zn1—O4—N8—O5	-5.9 (2)
O1W—Zn1—O4—N8	95.89 (15)	Zn1—O5—N8—O6	-173.0 (2)
O5—Zn1—O4—N8	3.73 (13)	Zn1—O5—N8—O4	6.2 (2)
O1—Zn1—O4—N8	-68.82 (15)	C4—N1—C1—C2	0.0 (3)
N1—Zn1—O5—N8	9.79 (19)	Zn1—N1—C1—C2	178.16 (17)
N4—Zn1—O5—N8	-178.14 (12)	N1—C1—C2—C3	0.3 (3)
O1W—Zn1—O5—N8	-87.01 (13)	C4—N2—C3—C2	0.0 (3)
O1—Zn1—O5—N8	101.86 (13)	C1—C2—C3—N2	-0.3 (4)
O4—Zn1—O5—N8	-3.63 (13)	C3—N2—C4—N3	178.6 (2)
N4—Zn1—N1—C1	126.33 (14)	C3—N2—C4—N1	0.4 (3)
O1W—Zn1—N1—C1	32.92 (15)	C1—N1—C4—N3	-178.6 (2)
O5—Zn1—N1—C1	-61.49 (19)	Zn1—N1—C4—N3	3.6 (3)
O1—Zn1—N1—C1	-147.00 (14)	C1—N1—C4—N2	-0.4 (3)
O4—Zn1—N1—C1	-50.43 (15)	Zn1—N1—C4—N2	-178.24 (15)
N4—Zn1—N1—C4	-55.78 (17)	C8—N4—C5—C6	-1.3 (3)
O1W—Zn1—N1—C4	-149.19 (16)	Zn1—N4—C5—C6	-178.4 (2)
O5—Zn1—N1—C4	116.40 (17)	N4—C5—C6—C7	-0.6 (4)
O1—Zn1—N1—C4	30.90 (17)	C8—N5—C7—C6	-0.1 (4)
O4—Zn1—N1—C4	127.46 (17)	C5—C6—C7—N5	1.3 (4)
N1—Zn1—N4—C5	-27.69 (17)	C7—N5—C8—N6	176.5 (2)
O1W—Zn1—N4—C5	68.58 (16)	C7—N5—C8—N4	-2.1 (3)
O5—Zn1—N4—C5	157.14 (15)	C5—N4—C8—N6	-175.85 (19)
O1—Zn1—N4—C5	-126.05 (16)	Zn1—N4—C8—N6	0.9 (3)
O4—Zn1—N4—C5	140.6 (3)	C5—N4—C8—N5	2.8 (3)
N1—Zn1—N4—C8	155.57 (14)	Zn1—N4—C8—N5	179.52 (15)
O1W—Zn1—N4—C8	-108.16 (15)		

*Hydrogen-bond geometry (Å, °)*

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
O1w—H11···O2 <sup>i</sup>	0.83 (1)	1.99 (2)	2.776 (2)	158 (3)
O1w—H12···N2 <sup>ii</sup>	0.84 (1)	1.94 (1)	2.754 (2)	165 (3)
N3—H31···O1	0.87 (1)	2.23 (2)	2.989 (3)	146 (2)
N3—H32···O5 <sup>iii</sup>	0.86 (1)	2.34 (2)	3.133 (3)	152 (3)
N6—H61···O1	0.87 (1)	2.37 (3)	3.010 (2)	131 (3)
N6—H61···O5	0.87 (1)	2.43 (2)	3.122 (3)	137 (3)
N6—H62···O1 <sup>iv</sup>	0.87 (1)	2.41 (2)	3.192 (2)	150 (3)
N6—H62···O3 <sup>iv</sup>	0.87 (1)	2.45 (2)	3.265 (3)	156 (3)

Symmetry codes: (i)  $x+1/2, y+1/2, z$ ; (ii)  $x+1/2, y-1/2, z$ ; (iii)  $x-1/2, y+1/2, z$ ; (iv)  $-x+1, y, -z+3/2$ .