

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Redetermination of 3-methylisoquinoline at 150 K

Andrew D. Bond

University of Southern Denmark, Department of Physics and Chemistry, Campusvej 55, 5230 Odense, Denmark

Correspondence e-mail: adb@chem.sdu.dk


Received 30 September 2010; accepted 5 October 2010

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.042; wR factor = 0.119; data-to-parameter ratio = 18.3.

The structure of the title compound, $\text{C}_{10}\text{H}_9\text{N}$, has been redetermined at 150 K. The redetermination is of significantly higher precision than a previous room-temperature structure [Ribar *et al.* (1974). *Cryst. Struct. Commun.* **3**, 323–325]. The C–N bond lengths for this redetermination are much closer to those observed in comparable structures, and the orientation of the methyl group with respect to the isoquinoline plane is clarified. Intermolecular weak C–H \cdots N contacts are present in the crystal.

Related literature

For the structure at room temperature, see: Ribar *et al.* (1974). For the structure of the parent compound isoquinoline, see: Hensen *et al.* (1999). The C–N bond length in the structure of Ribar *et al.* (1974) clearly lies outside of the main distribution for 19 relevant structural fragments in the Cambridge Structural Database, being the second shortest bond in the sample [one shorter bond exists for refcode SAKCIQ, but this structure has $R1 = 14.2\%$ (Trumpp-Kallmeyer *et al.*, 1998)]. The corresponding C–N bond length in this redetermination lies exactly at the mean of the CSD sample.



Experimental

Crystal data

$\text{C}_{10}\text{H}_9\text{N}$
 $M_r = 143.18$
 Monoclinic, $P2_1/c$
 $a = 6.1991$ (4) Å
 $b = 7.4176$ (6) Å
 $c = 16.5421$ (12) Å
 $\beta = 93.438$ (2)°
 $V = 759.28$ (10) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.07$ mm⁻¹
 $T = 150$ K
 $0.25 \times 0.15 \times 0.12$ mm

Data collection

Bruker–Nonius X8 APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2003)
 $T_{\min} = 0.826$, $T_{\max} = 0.991$
 9801 measured reflections
 1844 independent reflections
 1171 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.119$
 $S = 1.06$
 1844 reflections
 101 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.24$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.20$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C5}-\text{H5A}\cdots\text{N2}^i$	0.95	2.88	3.6891 (14)	144
$\text{C6}-\text{H6A}\cdots\text{N2}^{ii}$	0.95	2.64	3.5813 (15)	170

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2003); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

The Danish Natural Sciences Research Council and the Carlsberg Foundation are acknowledged for provision of the X-ray equipment.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5670).

References

- Bruker (2003). SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2004). APEX2. Bruker–Nonius BV, Delft, The Netherlands.
 Hensen, K., Mayr-Stein, R. & Bolte, M. (1999). *Acta Cryst.* **C55**, 1565–1567.
 Ribar, B., Divjakovic, V., Janic, I., Argay, G., Kalman, A. & Djuric, S. (1974). *Cryst. Struct. Commun.* **3**, 323–325.
 Sheldrick, G. M. (2003). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Trumpp-Kallmeyer, S., Rubin, J. R., Humblet, C., Hamby, J. M. & Hollis Showalter, H. D. (1998). *J. Med. Chem.* **41**, 1752–1763.

supporting information

Acta Cryst. (2010). E66, o2768 [https://doi.org/10.1107/S1600536810039838]

Redetermination of 3-methylisoquinoline at 150 K

Andrew D. Bond

S1. Comment

The structure of 3-iso-methylquinoline at room temperature has been reported by Ribar *et al.* (1974). This redetermination at 150 K provides significantly improved precision, and more regular positions for the H atoms.

Considering the C11—N2 bond length: the CCDC *Mogul* package identifies 19 relevant structural fragments in the CSD, with a mean bond length of 1.314 (10) Å. The structure of Ribar *et al.* [C—N = 1.300 (5) Å] lies clearly outside of the main distribution, being the second shortest bond in the sample (one shorter bond of 1.292 Å exists for refcode SAKCIQ, but this structure has $R1 = 14.2\%$ (Trumpp-Kallmeyer *et al.*, 1998). By contrast, the C1—N2 bond length of 1.3144 (13) Å in this redetermination corresponds exactly with the mean value. Alternation is also more clearly seen for the bond lengths C5—C6, C6—C7 and C7—C8 (1.3649 (16), 1.4093 (16) and 1.3646 (15) Å, respectively), compared to the previous structure.

Concerning the H atoms, the orientation of the methyl group in particular is clarified: in the structure of Ribar *et al.*, the H—C(methyl)—H angles are irregular (range 94.8–112.8 °) and the orientation of the group is such that one C—H bond is twisted from the isoquinoline plane with a C—C—C(methyl)—H torsion angle *ca* 22 °. In the redetermination, the refined orientation of the methyl group places one C—H bond much more clearly in the isoquinoline plane (torsion angle 5.8 (1) °). This also has an influence on the geometry observed for the intermolecular contact between the methyl group and a neighbouring isoquinoline molecule. In the redetermination, atom H11B lies over the centroid of the C5—C10 ring with H11B...Cg = 2.95 Å and C11—H11B...Cg = 131.9 Å.

S2. Experimental

The colourless block of (I) used for structure determination was taken directly from the sample as supplied by Aldrich Chemical Company.

S3. Refinement

H atoms bound to C(sp^2) were positioned geometrically with C—H = 0.95 Å and refined as riding with $U_{iso}(H) = 1.2 U_{eq}(C)$. The H atoms of the methyl group were positioned with C—H = 0.98 Å and refined as riding with $U_{iso}(H) = 1.5 U_{eq}(C)$, and with rotation about the local 3-fold axis.




Figure 1

Molecular structure showing displacement ellipsoids at 50% probability for non-H atoms.




Figure 2

Unit-cell contents.

3-methylisoquinoline

Crystal data

$C_{10}H_9N$

$M_r = 143.18$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1ybc$

$a = 6.1991(4)\ \text{\AA}$

$b = 7.4176(6)\ \text{\AA}$

$c = 16.5421(12)\ \text{\AA}$

$\beta = 93.438(2)^\circ$

$V = 759.28(10)\ \text{\AA}^3$

$Z = 4$

$F(000) = 304$

$D_x = 1.253\ \text{Mg m}^{-3}$

Melting point = 336–338 K

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1780 reflections

$\theta = 2.5\text{--}25.4^\circ$

$\mu = 0.07\ \text{mm}^{-1}$

$T = 150\ \text{K}$

Block, colourless

$0.25 \times 0.15 \times 0.12\ \text{mm}$

Data collection

Bruker–Nonius X8 APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
thin-slice ω and φ scans
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2003)
 $T_{\min} = 0.826$, $T_{\max} = 0.991$

9801 measured reflections
1844 independent reflections
1171 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$
 $\theta_{\max} = 28.4^\circ$, $\theta_{\min} = 3.7^\circ$
 $h = -7 \rightarrow 8$
 $k = -9 \rightarrow 9$
 $l = -21 \rightarrow 20$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.119$
 $S = 1.06$
1844 reflections
101 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0637P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.24 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.11212 (16)	0.91552 (15)	0.31390 (6)	0.0232 (3)
H1A	-0.0145	0.9742	0.3301	0.028*
N2	0.26422 (14)	0.87917 (12)	0.37033 (5)	0.0247 (3)
C3	0.44782 (16)	0.79419 (15)	0.34720 (7)	0.0234 (3)
C4	0.47647 (16)	0.75059 (15)	0.26814 (7)	0.0232 (3)
H4A	0.6070	0.6944	0.2544	0.028*
C5	0.32884 (18)	0.74255 (15)	0.12395 (7)	0.0257 (3)
H5A	0.4564	0.6873	0.1066	0.031*
C6	0.16084 (19)	0.77766 (16)	0.06908 (7)	0.0294 (3)
H6A	0.1709	0.7434	0.0141	0.035*
C7	-0.02738 (18)	0.86413 (16)	0.09309 (7)	0.0289 (3)
H7A	-0.1420	0.8894	0.0540	0.035*
C8	-0.04638 (16)	0.91186 (15)	0.17202 (7)	0.0247 (3)
H8A	-0.1734	0.9708	0.1877	0.030*
C9	0.12341 (16)	0.87349 (14)	0.23056 (6)	0.0205 (3)
C10	0.31404 (16)	0.78813 (14)	0.20670 (7)	0.0209 (3)

C11	0.61079 (18)	0.74985 (17)	0.41473 (7)	0.0317 (3)
H11A	0.6495	0.8598	0.4451	0.048*
H11B	0.7403	0.6991	0.3923	0.048*
H11C	0.5493	0.6617	0.4510	0.048*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0222 (6)	0.0219 (7)	0.0259 (6)	0.0014 (5)	0.0044 (5)	0.0006 (5)
N2	0.0262 (5)	0.0249 (6)	0.0232 (5)	0.0014 (4)	0.0022 (4)	0.0010 (4)
C3	0.0235 (6)	0.0196 (7)	0.0271 (7)	-0.0011 (4)	0.0003 (5)	0.0040 (5)
C4	0.0198 (5)	0.0212 (6)	0.0289 (7)	0.0011 (4)	0.0050 (5)	0.0025 (5)
C5	0.0300 (6)	0.0221 (6)	0.0258 (7)	0.0000 (5)	0.0089 (5)	-0.0002 (5)
C6	0.0402 (7)	0.0277 (7)	0.0206 (6)	-0.0059 (5)	0.0043 (5)	0.0007 (5)
C7	0.0287 (6)	0.0299 (7)	0.0274 (7)	-0.0049 (5)	-0.0050 (5)	0.0048 (5)
C8	0.0216 (6)	0.0233 (7)	0.0290 (7)	-0.0006 (4)	0.0001 (5)	0.0028 (5)
C9	0.0211 (5)	0.0172 (6)	0.0233 (6)	-0.0019 (4)	0.0028 (4)	0.0021 (5)
C10	0.0223 (6)	0.0173 (6)	0.0235 (6)	-0.0028 (4)	0.0042 (4)	0.0016 (5)
C11	0.0303 (6)	0.0335 (8)	0.0307 (7)	0.0015 (5)	-0.0040 (5)	0.0053 (6)

Geometric parameters (Å, °)

C1—N2	1.3144 (13)	C6—C7	1.4093 (16)
C1—C9	1.4194 (15)	C6—H6A	0.950
C1—H1A	0.950	C7—C8	1.3646 (15)
N2—C3	1.3753 (13)	C7—H7A	0.950
C3—C4	1.3690 (15)	C8—C9	1.4157 (14)
C3—C11	1.4971 (15)	C8—H8A	0.950
C4—C10	1.4148 (15)	C9—C10	1.4174 (15)
C4—H4A	0.950	C11—H11A	0.980
C5—C6	1.3649 (16)	C11—H11B	0.980
C5—C10	1.4184 (16)	C11—H11C	0.980
C5—H5A	0.950		
N2—C1—C9	124.73 (10)	C8—C7—H7A	119.7
N2—C1—H1A	117.6	C6—C7—H7A	119.7
C9—C1—H1A	117.6	C7—C8—C9	119.93 (10)
C1—N2—C3	117.83 (9)	C7—C8—H8A	120.0
C4—C3—N2	122.08 (10)	C9—C8—H8A	120.0
C4—C3—C11	122.72 (10)	C8—C9—C10	119.83 (10)
N2—C3—C11	115.20 (10)	C8—C9—C1	122.84 (10)
C3—C4—C10	120.81 (10)	C10—C9—C1	117.33 (10)
C3—C4—H4A	119.6	C4—C10—C9	117.21 (10)
C10—C4—H4A	119.6	C4—C10—C5	124.19 (10)
C6—C5—C10	120.36 (10)	C9—C10—C5	118.59 (10)
C6—C5—H5A	119.8	C3—C11—H11A	109.5
C10—C5—H5A	119.8	C3—C11—H11B	109.5
C5—C6—C7	120.73 (11)	H11A—C11—H11B	109.5

C5—C6—H6A	119.6	C3—C11—H11C	109.5
C7—C6—H6A	119.6	H11A—C11—H11C	109.5
C8—C7—C6	120.54 (11)	H11B—C11—H11C	109.5
C9—C1—N2—C3	-0.04 (17)	N2—C1—C9—C8	178.66 (10)
C1—N2—C3—C4	1.19 (16)	N2—C1—C9—C10	-0.81 (17)
C1—N2—C3—C11	-177.61 (9)	C3—C4—C10—C9	0.55 (16)
N2—C3—C4—C10	-1.46 (17)	C3—C4—C10—C5	-178.11 (10)
C11—C3—C4—C10	177.24 (10)	C8—C9—C10—C4	-178.97 (9)
C10—C5—C6—C7	1.80 (17)	C1—C9—C10—C4	0.51 (15)
C5—C6—C7—C8	-1.03 (18)	C8—C9—C10—C5	-0.23 (16)
C6—C7—C8—C9	-0.38 (17)	C1—C9—C10—C5	179.26 (9)
C7—C8—C9—C10	0.99 (16)	C6—C5—C10—C4	177.49 (10)
C7—C8—C9—C1	-178.46 (10)	C6—C5—C10—C9	-1.16 (16)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C5—H5A...N2 ⁱ	0.95	2.88	3.6891 (14)	144
C6—H6A...N2 ⁱⁱ	0.95	2.64	3.5813 (15)	170

Symmetry codes: (i) $-x+1, y-1/2, -z+1/2$; (ii) $x, -y+3/2, z-1/2$.