

2-(2-Oxothiolan-3-yl)isoindoline-1,3-dione

Abdul Rauf Raza,^a Aisha Saddiq,^a M. Nawaz Tahir^{b*} and Sadia Saddiq^a

^aDepartment of Chemistry, University of Sargodha, Sargodha, Pakistan, and

^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 24 October 2010; accepted 25 October 2010

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.055; wR factor = 0.151; data-to-parameter ratio = 12.6.

In the title compound, $\text{C}_{12}\text{H}_9\text{NO}_3\text{S}$, the isoindoline-1,3-dione group is almost planar, with an r.m.s. deviation of 0.020 \AA , whereas the heterocyclic ring approximates to an envelope with the methylene group not adjacent to the S atom in the flap position. A short intramolecular C–H···O contact generates an S(6) ring motif. In the crystal structure, weak aromatic π – π stacking interactions occur between the centroids of the benzene rings at a distance of $3.558(2)\text{ \AA}$.

Related literature

For background to isocoumarins, see: Hussain *et al.* (2001); Lee *et al.* (2001); Nozawa *et al.* (1981). For related crystal structures, see: Beck *et al.* (2007); Freer & Kraut (1965). For graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{12}\text{H}_9\text{NO}_3\text{S}$	$V = 1095.2(3)\text{ \AA}^3$
$M_r = 247.26$	$Z = 4$
Monoclinic, $P2_1/c$	$\text{Mo } K\alpha$ radiation
$a = 8.0601(13)\text{ \AA}$	$\mu = 0.29\text{ mm}^{-1}$
$b = 6.9860(11)\text{ \AA}$	$T = 296\text{ K}$
$c = 19.709(3)\text{ \AA}$	$0.24 \times 0.10 \times 0.08\text{ mm}$
$\beta = 99.296(9)\text{ }^\circ$	

Data collection

Bruker Kappa APEXII CCD diffractometer	14781 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	1934 independent reflections
$T_{\min} = 0.968$, $T_{\max} = 0.978$	1105 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.093$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$	154 parameters
$wR(F^2) = 0.151$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\max} = 0.40\text{ e } \text{\AA}^{-3}$
1934 reflections	$\Delta\rho_{\min} = -0.31\text{ e } \text{\AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C10—H10B···O1	0.97	2.52	3.149 (5)	122

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the provision of funds for the purchase of the diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5705).

References

- Beck, B., Srivastava, S. & Dömling, A. (2007). *Heterocycles*, **73**, 177–182.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.*, **34**, 1555–1573.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.*, **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.*, **32**, 837–838.
- Freer, S. T. & Kraut, J. (1965). *Acta Cryst.*, **19**, 992–1002.
- Hussain, M. T., Rama, N. H. & Malik, A. (2001). *Indian J. Chem. Sect. B*, **40**, 372–376.
- Lee, J. H., Park, Y. J., Kim, H. S., Hong, Y. S., Kim, K.-W. & Lee, J. J. (2001). *J. Antibiot.*, **54**, 463–466.
- Nozawa, K., Yamada, M., Tsuda, Y., Kawai, K. & Nakajima, S. (1981). *Chem. Pharm. Bull.*, **29**, 2689–2691.
- Sheldrick, G. M. (2008). *Acta Cryst. A*, **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D*, **65**, 148–155.

supporting information

Acta Cryst. (2010). E66, o2979 [https://doi.org/10.1107/S1600536810043400]

2-(2-Oxothiolan-3-yl)isoindoline-1,3-dione

Abdul Rauf Raza, Aisha Saddiq, M. Nawaz Tahir and Sadia Saddiq

S1. Comment

Isocoumarins are important class of naturally occurring compounds. Their structure is similar to coumarin but with inverted lactone ring. They exhibit a wide range of biological activities such as antimicrobial (Hussain *et al.*, 2001), anti-fungal (Nozawa *et al.*, 1981), anti-angiogenic (Lee *et al.*, 2001). The title compound (**I**, Fig. 1) was obtained as an interesting side-product during the synthesis of isocoumarin.

The crystal structure of D,L-homocysteine thiolactone hydrochloride (Freer & Kraut, 1965) and (*R*^{*})-2-(4-Chlorophenyl)-N-(hept-4-yl)-2-((S^{*})-2-oxotetrahydrothiophen-3-ylamino)acetamide (Beck *et al.*, 2007) have been reported which contain the heterocyclic ring.


The title compound essentially consists of monomers. In (**I**), the 2-benzoazole-1,3-dione group (C1–C8/N1/O1/O2) is planar with r.m.s. deviation of 0.0196 Å. The heterocyclic ring (C9/C10/C11/S1/C12) is not planar as the r.m.s. deviation of the plane is 0.1511 Å. The dihedral angle between these two groups is 88.05 (10)^o. There exist weak intramolecular H-bondings of C—H···O type (Table 1, Fig. 1) completing S(5) and S(6) ring motifs (Bernstein *et al.*, 1995). There exist π–π interaction between the centroids of benzene rings at a distance of 3.558 (2) Å [symmetry: 1 - *x*, - *y*, - *z*].

S2. Experimental

Homophthalic acid (1.0 g, 5.5 mmol, 1 eq) was added to 4-(methylthio)-2-(1,3-dioxoisindolin-2-yl)butanoyl chloride (6.5 g, 0.022 mol, 4 eq) and the mixture was heated at 473 K with continuous stirring for 6 h. The crude product was added to chilled water (20 ml), partitioned with EtOAc (3 × 25 ml), organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The column chromatographic separation using EtOAc and n-hexane (3:7) as mobile phase afforded the title compound (**I**) as colourless needles.

S3. Refinement

The H atoms were positioned geometrically with C—H = 0.93–0.97 Å and were included in the refinement in the riding model approximation, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.2$ for all H atoms.

**Figure 1**

View of (I) with displacement ellipsoids drawn at the 50% probability level.

2-(2-Oxothiolan-3-yl)isoindoline-1,3-dione*Crystal data*

$C_{12}H_9NO_3S$
 $M_r = 247.26$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 8.0601 (13)$ Å
 $b = 6.9860 (11)$ Å
 $c = 19.709 (3)$ Å
 $\beta = 99.296 (9)$ °
 $V = 1095.2 (3)$ Å³
 $Z = 4$

$F(000) = 512$
 $D_x = 1.500 \text{ Mg m}^{-3}$
 $Mo K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 1105 reflections
 $\theta = 2.1\text{--}25.0$ °
 $\mu = 0.29 \text{ mm}^{-1}$
 $T = 296 \text{ K}$
Needle, colourless
 $0.24 \times 0.10 \times 0.08$ mm

Data collection

Bruker Kappa APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
Detector resolution: 8.2 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2005)
 $T_{\min} = 0.968$, $T_{\max} = 0.978$

14781 measured reflections
1934 independent reflections
1105 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.093$
 $\theta_{\max} = 25.0$ °, $\theta_{\min} = 2.1$ °
 $h = -9 \rightarrow 9$
 $k = -8 \rightarrow 8$
 $l = -23 \rightarrow 23$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.151$
 $S = 1.02$
1934 reflections
154 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0651P)^2 + 0.3208P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.40 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.31 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	1.27260 (14)	0.17317 (15)	0.21281 (6)	0.0666 (4)
O1	1.0043 (4)	0.2417 (4)	0.01824 (15)	0.0690 (11)
O2	0.6135 (4)	0.3134 (4)	0.15927 (16)	0.0694 (11)
O3	1.0050 (3)	-0.0304 (4)	0.16664 (14)	0.0619 (11)
N1	0.8391 (4)	0.2932 (4)	0.10189 (16)	0.0463 (11)
C1	0.8677 (5)	0.2558 (5)	0.0348 (2)	0.0496 (14)
C2	0.6994 (5)	0.2417 (5)	-0.0077 (2)	0.0445 (14)
C3	0.6538 (6)	0.2073 (5)	-0.0763 (2)	0.0600 (16)
C4	0.4821 (7)	0.1996 (6)	-0.1015 (2)	0.0724 (19)
C5	0.3638 (6)	0.2223 (6)	-0.0585 (3)	0.073 (2)
C6	0.4108 (5)	0.2546 (5)	0.0105 (3)	0.0580 (16)
C7	0.5801 (4)	0.2646 (5)	0.0350 (2)	0.0469 (14)
C8	0.6676 (5)	0.2926 (5)	0.1061 (2)	0.0486 (14)
C9	0.9686 (4)	0.3132 (5)	0.1613 (2)	0.0492 (12)
C10	1.0927 (5)	0.4715 (5)	0.1568 (2)	0.0537 (16)
C11	1.2487 (5)	0.4309 (5)	0.2093 (2)	0.0553 (16)
C12	1.0662 (5)	0.1259 (5)	0.17663 (18)	0.0444 (12)
H3	0.73362	0.18977	-0.10491	0.0718*
H4	0.44640	0.17879	-0.14821	0.0870*
H5	0.25006	0.21559	-0.07677	0.0880*
H6	0.33150	0.26915	0.03956	0.0696*
H9	0.91266	0.34070	0.20085	0.0591*
H10A	1.04437	0.59374	0.16649	0.0646*
H10B	1.12157	0.47652	0.11092	0.0646*
H11A	1.23521	0.48081	0.25402	0.0667*
H11B	1.34670	0.49028	0.19538	0.0667*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0529 (7)	0.0572 (7)	0.0828 (9)	0.0032 (5)	-0.0094 (6)	0.0022 (6)
O1	0.0456 (18)	0.086 (2)	0.078 (2)	-0.0061 (15)	0.0178 (16)	-0.0142 (16)
O2	0.0619 (19)	0.075 (2)	0.076 (2)	-0.0060 (16)	0.0255 (17)	-0.0102 (16)
O3	0.0674 (19)	0.0400 (16)	0.078 (2)	-0.0076 (14)	0.0106 (16)	-0.0004 (14)
N1	0.0420 (19)	0.0466 (18)	0.049 (2)	0.0011 (14)	0.0038 (16)	0.0021 (15)
C1	0.040 (2)	0.042 (2)	0.067 (3)	-0.0027 (18)	0.009 (2)	0.0031 (18)

C2	0.043 (2)	0.037 (2)	0.052 (3)	-0.0082 (16)	0.003 (2)	0.0049 (17)
C3	0.069 (3)	0.051 (2)	0.058 (3)	-0.013 (2)	0.004 (2)	0.009 (2)
C4	0.091 (4)	0.056 (3)	0.059 (3)	-0.017 (3)	-0.022 (3)	0.010 (2)
C5	0.054 (3)	0.044 (3)	0.111 (5)	-0.006 (2)	-0.019 (3)	0.018 (3)
C6	0.041 (2)	0.037 (2)	0.092 (4)	-0.0023 (18)	-0.001 (2)	0.004 (2)
C7	0.039 (2)	0.029 (2)	0.069 (3)	-0.0025 (16)	-0.002 (2)	0.0051 (17)
C8	0.042 (2)	0.038 (2)	0.068 (3)	-0.0018 (17)	0.016 (2)	0.0010 (19)
C9	0.046 (2)	0.048 (2)	0.053 (2)	-0.0043 (18)	0.006 (2)	-0.0034 (19)
C10	0.061 (3)	0.041 (2)	0.058 (3)	-0.005 (2)	0.006 (2)	-0.0018 (18)
C11	0.056 (3)	0.053 (2)	0.056 (3)	-0.007 (2)	0.006 (2)	-0.009 (2)
C12	0.048 (2)	0.042 (2)	0.044 (2)	-0.0006 (18)	0.0098 (19)	0.0016 (17)

Geometric parameters (\AA , $\text{^{\circ}}$)

S1—C11	1.811 (4)	C6—C7	1.374 (6)
S1—C12	1.733 (4)	C7—C8	1.477 (5)
O1—C1	1.202 (5)	C9—C10	1.503 (5)
O2—C8	1.207 (5)	C9—C12	1.532 (5)
O3—C12	1.201 (5)	C10—C11	1.521 (6)
N1—C1	1.403 (5)	C3—H3	0.9300
N1—C8	1.398 (5)	C4—H4	0.9300
N1—C9	1.444 (5)	C5—H5	0.9300
C1—C2	1.478 (6)	C6—H6	0.9300
C2—C3	1.364 (5)	C9—H9	0.9800
C2—C7	1.386 (5)	C10—H10A	0.9700
C3—C4	1.394 (7)	C10—H10B	0.9700
C4—C5	1.383 (7)	C11—H11A	0.9700
C5—C6	1.370 (8)	C11—H11B	0.9700
C11—S1—C12	94.86 (18)	S1—C12—O3	125.6 (3)
C1—N1—C8	111.7 (3)	S1—C12—C9	110.3 (2)
C1—N1—C9	125.1 (3)	O3—C12—C9	124.1 (3)
C8—N1—C9	123.0 (3)	C2—C3—H3	121.00
O1—C1—N1	124.6 (4)	C4—C3—H3	122.00
O1—C1—C2	129.6 (4)	C3—C4—H4	119.00
N1—C1—C2	105.7 (3)	C5—C4—H4	119.00
C1—C2—C3	130.5 (4)	C4—C5—H5	119.00
C1—C2—C7	108.2 (3)	C6—C5—H5	119.00
C3—C2—C7	121.3 (4)	C5—C6—H6	121.00
C2—C3—C4	116.9 (4)	C7—C6—H6	121.00
C3—C4—C5	121.4 (4)	N1—C9—H9	107.00
C4—C5—C6	121.3 (5)	C10—C9—H9	107.00
C5—C6—C7	117.2 (4)	C12—C9—H9	107.00
C2—C7—C6	121.9 (4)	C9—C10—H10A	110.00
C2—C7—C8	108.7 (3)	C9—C10—H10B	110.00
C6—C7—C8	129.5 (4)	C11—C10—H10A	110.00
O2—C8—N1	123.4 (4)	C11—C10—H10B	110.00
O2—C8—C7	131.0 (4)	H10A—C10—H10B	108.00

N1—C8—C7	105.6 (3)	S1—C11—H11A	110.00
N1—C9—C10	115.1 (3)	S1—C11—H11B	110.00
N1—C9—C12	111.0 (3)	C10—C11—H11A	110.00
C10—C9—C12	108.3 (3)	C10—C11—H11B	110.00
C9—C10—C11	107.9 (3)	H11A—C11—H11B	109.00
S1—C11—C10	106.4 (2)		
C12—S1—C11—C10	-19.3 (3)	C1—C2—C7—C6	-178.6 (3)
C11—S1—C12—O3	-178.8 (4)	C1—C2—C7—C8	-0.4 (4)
C11—S1—C12—C9	-0.9 (3)	C3—C2—C7—C6	-0.3 (6)
C8—N1—C1—O1	-177.2 (3)	C3—C2—C7—C8	178.0 (3)
C8—N1—C1—C2	3.5 (4)	C2—C3—C4—C5	-1.2 (6)
C9—N1—C1—O1	-2.4 (6)	C3—C4—C5—C6	0.5 (6)
C9—N1—C1—C2	178.3 (3)	C4—C5—C6—C7	0.4 (6)
C1—N1—C8—O2	176.8 (3)	C5—C6—C7—C2	-0.5 (5)
C1—N1—C8—C7	-3.8 (4)	C5—C6—C7—C8	-178.3 (4)
C9—N1—C8—O2	1.9 (5)	C2—C7—C8—O2	-178.1 (4)
C9—N1—C8—C7	-178.7 (3)	C2—C7—C8—N1	2.5 (4)
C1—N1—C9—C10	59.6 (4)	C6—C7—C8—O2	-0.1 (7)
C1—N1—C9—C12	-63.9 (4)	C6—C7—C8—N1	-179.5 (4)
C8—N1—C9—C10	-126.2 (3)	N1—C9—C10—C11	-160.7 (3)
C8—N1—C9—C12	110.3 (4)	C12—C9—C10—C11	-35.8 (4)
O1—C1—C2—C3	0.8 (7)	N1—C9—C12—S1	148.7 (2)
O1—C1—C2—C7	178.9 (4)	N1—C9—C12—O3	-33.4 (5)
N1—C1—C2—C3	-180.0 (4)	C10—C9—C12—S1	21.4 (4)
N1—C1—C2—C7	-1.8 (4)	C10—C9—C12—O3	-160.7 (4)
C1—C2—C3—C4	179.0 (4)	C9—C10—C11—S1	34.4 (4)
C7—C2—C3—C4	1.1 (5)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C10—H10B···O1	0.97	2.52	3.149 (5)	122