

10-Methyl-9-[2-(propan-2-yl)phenoxy-carbonyl]acridinium trifluoromethane-sulfonate

Damian Trzybiński, Karol Krzymiński and Jerzy Błażejowski*‡

Faculty of Chemistry, University of Gdańsk, J. Sobieskiego 18, 80-952 Gdańsk, Poland

Correspondence e-mail: bla@chem.univ.gda.pl


Received 30 September 2010; accepted 4 October 2010

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.045; wR factor = 0.116; data-to-parameter ratio = 13.1.

In the crystal of the title compound, $\text{C}_{24}\text{H}_{22}\text{NO}_2^+\cdot\text{CF}_3\text{SO}_3^-$, adjacent cations and anions are connected through $\text{C}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\text{F}$ and $\text{S}-\text{O}\cdots\pi$ interactions, while neighboring cations via $\pi-\pi$ interactions [centroid–centroid distance = $3.962(2)\text{ \AA}$]. The acridine and benzene ring systems are oriented at a dihedral angle of $14.6(1)^\circ$. The carboxyl group is twisted at an angle of $87.6(1)^\circ$ relative to the acridine skeleton. The mean planes of adjacent acridine units are parallel or inclined at an angle of $13.4(1)^\circ$ in the crystal structure.

Related literature

For background to the chemiluminogenic properties of 9-phenoxy carbonyl-10-methylacridinium trifluoromethane-sulfonates, see: Natrajan *et al.* (2010); Brown *et al.* (2009); King *et al.* (2007); Rak *et al.* (1999); Roda *et al.* (2003); Zomer & Jacquemijns (2001). For related structures, see: Sikorski *et al.* (2006, 2007); Trzybiński *et al.* (2010). For intermolecular interactions, see: Bianchi *et al.* (2004); Dorn *et al.* (2005); Hunter *et al.* (2001); Lyssenko & Antipin (2004); Novoa *et al.* (2006). For the synthesis, see: Sato (1996); Trzybiński *et al.* (2010).



Experimental

Crystal data

$\text{C}_{24}\text{H}_{22}\text{NO}_2^+\cdot\text{CF}_3\text{SO}_3^-$	$V = 2340.47(17)\text{ \AA}^3$
$M_r = 505.51$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 14.4346(7)\text{ \AA}$	$\mu = 0.20\text{ mm}^{-1}$
$b = 12.9677(5)\text{ \AA}$	$T = 295\text{ K}$
$c = 13.0862(5)\text{ \AA}$	$0.32 \times 0.20 \times 0.05\text{ mm}$
$\beta = 107.160(5)^\circ$	

Data collection

Oxford Diffraction Gemini R Ultra Ruby CCD diffractometer	17556 measured reflections
Absorption correction: multi-scan (<i>CrysAlis RED</i> ; Oxford Diffraction, 2008)	4169 independent reflections
$T_{\min} = 0.955$, $T_{\max} = 1.000$	2436 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$	319 parameters
$wR(F^2) = 0.116$	H-atom parameters constrained
$S = 0.93$	$\Delta\rho_{\max} = 0.34\text{ e \AA}^{-3}$
4169 reflections	$\Delta\rho_{\min} = -0.26\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}4-\text{H}4\cdots\text{O}29^i$	0.93	2.48	3.363 (3)	159
$\text{C}27-\text{H}27\text{C}\cdots\text{F}35^i$	0.96	2.51	3.250 (4)	134

Symmetry code: (i) $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$.

Table 2
 $\text{S}-\text{O}\cdots\pi$ interactions (\AA , $^\circ$).

$\text{Cg}2$ is the centroids of the $\text{C}1-\text{C}4/\text{C}11/\text{C}12$ ring.

X	I	J	$I\cdots J$	$X\cdots J$	$X-I\cdots J$
S28	O31	$\text{Cg}2^{ii}$	3.208 (2)	4.128 (2)	120.3 (2)

Symmetry code: (ii) $-x + 2, -y + 1, -z + 1$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2008); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2008); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

‡ to whom correspondence should be addressed

This study was financed by the State Funds for Scientific Research (grant No. N204 123 32/3143 – contract No. 3143/H03/2007/32 of the Polish Ministry of Research and Higher Education for the period 2007–2010 – and DS/8820-4-0087-0).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5040).

References

- Bianchi, R., Forni, A. & Pilati, T. (2004). *Acta Cryst.* **B60**, 559–568.
- Brown, R. C., Li, Z., Rutter, A. J., Mu, X., Weeks, O. H., Smith, K. & Weeks, I. (2009). *Org. Biomol. Chem.* **7**, 386–394.
- Dorn, T., Janiak, C. & Abu-Shandi, K. (2005). *CrystEngComm*, **7**, 633–641.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Hunter, C. A., Lawson, K. R., Perkins, J. & Urch, C. J. (2001). *J. Chem. Soc. Perkin Trans. 2*, pp. 651–669.
- King, D. W., Cooper, W. J., Rusak, S. A., Peake, B. M., Kiddie, J. J., O'Sullivan, D. W., Melamed, M. L., Morgan, C. R. & Theberge, S. M. (2007). *Anal. Chem.* **79**, 4169–4176.
- Lyssenko, K. A. & Antipin, M. Y. (2004). *Russ. Chem. Bull. Int. Ed.* **53**, 10–17.
- Natrajan, A., Sharpe, D., Costello, J. & Jiang, Q. (2010). *Anal. Biochem.* **406**, 204–213.
- Novoa, J. J., Mota, F. & D'Oria, E. (2006). *Hydrogen Bonding – New Insights*, edited by S. Grabowski, pp. 193–244. The Netherlands: Springer.
- Oxford Diffraction (2008). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
- Rak, J., Skurski, P. & Błażejowski, J. (1999). *J. Org. Chem.* **64**, 3002–3008.
- Roda, A., Guardigli, M., Michelini, E., Mirasoli, M. & Pasini, P. (2003). *Anal. Chem.* **75**, 462–470.
- Sato, N. (1996). *Tetrahedron Lett.* **37**, 8519–8522.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sikorski, A., Krzymiński, K., Białońska, A., Lis, T. & Błażejowski, J. (2006). *Acta Cryst. E62*, o822–o824.
- Sikorski, A., Krzymiński, K., Malecha, P., Lis, T. & Błażejowski, J. (2007). *Acta Cryst. E63*, o4484–o4485.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Trzybiński, D., Krzymiński, K., Sikorski, A., Malecha, P. & Błażejowski, J. (2010). *Acta Cryst. E66*, o826–o827.
- Zomer, G. & Jacquemyns, M. (2001). *Chemiluminescence in Analytical Chemistry*, edited by A. M. Garcia-Campana & W. R. G. Baeyens, pp. 529–549. New York: Marcel Dekker.

supporting information

Acta Cryst. (2010). E66, o2773–o2774 [https://doi.org/10.1107/S160053681003953X]

10-Methyl-9-[2-(propan-2-yl)phenoxy carbonyl]acridinium trifluoromethane-sulfonate

Damian Trzybiński, Karol Krzymiński and Jerzy Błażejowski

S1. Comment

9-(Phenoxy carbonyl)-10-methylacridinium salts have long been known as chemiluminescent indicators or the chemiluminogenic fragments of chemiluminescent labels widely used in assays of biologically and environmentally important entities such as antigens, antibodies, enzymes or DNA fragments (Zomer & Jacquemijns, 2001; Roda *et al.*, 2003; King *et al.*, 2007; Brown *et al.*, 2009; Natrajan *et al.*, 2010). The cations of these salts are oxidized with H₂O₂ in alkaline media, a reaction that produces light. The latter process is accompanied by the removal of the phenoxy carbonyl fragment and the conversion of the remaining part of the molecules to electronically excited, light-emitting 10-methyl-9-acridinone (Rak *et al.*, 1999). The efficiency of chemiluminescence – crucial for analytical applications – is affected by the structure of the phenyl fragment (Zomer & Jacquemijns, 2001; Natrajan *et al.*, 2010). In the search for efficient chemiluminogens we synthesized 9-(phenoxy carbonyl)-10-methylacridinium derivatives alkyl substituted in the *ortho* position of the phenyl fragment. Here we present the structure of 9-(2-*i*-propylphenoxy carbonyl)-10-methylacridinium trifluoromethanesulfonate.

In the cation of the title compound (Fig. 1), the bond lengths and angles characterizing the geometry of the acridinium moiety are typical of acridine-based derivatives (Sikorski *et al.*, 2006; Sikorski *et al.*, 2007; Trzybiński *et al.*, 2010). With respective average deviations from planarity of 0.0127 (3) Å and 0.0030 (3) Å, the acridine and benzene ring systems are oriented at a dihedral angle of 14.6 (1)°. The carboxyl group is twisted at an angle of 87.6 (1)° relative to the acridine skeleton. The mean planes of the adjacent acridine moieties are parallel (remain at an angle 0.0 (1)°) or inclined at an angle of 87.6 (1)° in the lattice. In the series of 9-phenoxy carbonyl-10-methylacridinium trifluoromethanesulfonates substituted in the *ortho* position of the phenyl fragment with Me (Sikorski *et al.*, 2006), Et (Trzybiński *et al.*, 2010), *i*-Pr (this work) and *t*-Bu (Sikorski *et al.*, 2007), the dihedral angle between acridine and the benzene ring, and that between the carboxyl group and the acridine skeleton, increase in the order 2-Et < 2-*i*-Pr < 2-Me < 2-*t*-Bu, and 2-*t*-Bu < 2-Et < 2-*i*-Pr < 2-Me, respectively. This implies that increasing size of the alkyl substituent in the *ortho* position does not systematically influence the mutual arrangement of the above mentioned fragments of the molecules.

In the crystal structure, each anion is connected to the adjacent cations through C–H···O (Table 1, Fig. 2), C–H···F (Table 1, Fig. 2) and S–O···π (Table 2, Fig. 2) interactions. Neighboring cations contact each other via π–π (Table 3, Fig. 2) interactions. The C–H···O (Novoa *et al.*, 2006) and C–H···F (Bianchi *et al.*, 2004; Lyssenko & Antipin, 2004) interactions are of the hydrogen bond type. The S–O···π (Dorn *et al.*, 2005) and the π–π (Hunter *et al.*, 2001) interactions should be of an attractive nature. The crystal structure is stabilized by a network of these short-range specific interactions and by long-range electrostatic interactions between ions.

S2. Experimental

9-(Chlorocarbonyl)acridine, obtained by treating acridine-9-carboxylic acid with a tenfold molar excess of thionyl chloride, was first esterified with 2-*i*-propylphenol in anhydrous dichloromethane in the presence of *N,N*-diethyl-ethanamine and a catalytic amount of *N,N*-dimethyl-4-pyridinamine (room temperature, 15 h) (Sato, 1996) to obtain 2-*i*-propylphenylacridine-9-carboxylate (purified chromatographically (SiO_2 , cyclohexane/ethyl acetate, 1/1 v/v)). The latter compound was then quaternarized with a fivefold molar excess of methyl trifluoromethanesulfonate dissolved in anhydrous dichloromethane (Trzybiński *et al.*, 2010). The crude 9-(*i*-propylphenoxy carbonyl)-10-methylacridinium trifluoromethanesulfonate was dissolved in a small amount of ethanol, filtered and precipitated with 20 v/v excess of diethyl ether. Yellow crystals suitable for X-ray investigations were grown from absolute ethanol solution (m.p. 464–466 K).

S3. Refinement

H atoms were positioned geometrically, with C—H = 0.93 Å and 0.96 Å for the aromatic and alkyl H atoms, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.2$ for the aromatic and $x = 1.5$ for the alkyl H atoms.





Figure 1


The molecular structure of the title compound showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 25% probability level and H atoms are shown as small spheres of arbitrary radius. $Cg1$, $Cg2$, and $Cg3$ denote the ring centroids.

**Figure 2**

The arrangement of the ions in the crystal structure. The C–H···O and C–H···F interactions are represented by dashed lines, the S–O··· π and π ··· π contacts by dotted lines. H atoms not involved in interactions have been omitted. [Symmetry codes: (i) $-x + 2, y + 1/2, -z + 1/2$; (ii) $-x + 2, -y + 1, -z + 1$; (iii) $-x + 1, -y + 2, -z + 1$.]

10-Methyl-9-[2-(propan-2-yl)phenoxy carbonyl]acridinium trifluoromethanesulfonate

Crystal data



$M_r = 505.51$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 14.4346 (7)$ Å

$b = 12.9677 (5)$ Å

$c = 13.0862 (5)$ Å

$\beta = 107.160 (5)^\circ$

$V = 2340.47 (17)$ Å³

$Z = 4$

$F(000) = 1048$

$D_x = 1.435$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5919 reflections

$\theta = 3.0\text{--}29.2^\circ$

$\mu = 0.20$ mm⁻¹

$T = 295$ K

Prism, yellow

$0.32 \times 0.20 \times 0.05$ mm

Data collection

Oxford Diffraction Gemini R Ultra Ruby CCD diffractometer
 Radiation source: Enhanced (Mo) X-ray Source
 Graphite monochromator
 Detector resolution: 10.4002 pixels mm⁻¹
 ω scans
 Absorption correction: multi-scan
 (*CrysAlis RED*; Oxford Diffraction, 2008)
 $T_{\min} = 0.955$, $T_{\max} = 1.000$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.116$
 $S = 0.93$
 4169 reflections
 319 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods

17556 measured reflections
 4169 independent reflections
 2436 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$
 $\theta_{\max} = 25.1^\circ$, $\theta_{\min} = 3.0^\circ$
 $h = -17 \rightarrow 15$
 $k = -15 \rightarrow 15$
 $l = -15 \rightarrow 14$

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0687P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.34 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.79663 (19)	0.93471 (19)	0.23882 (19)	0.0538 (6)
H1	0.7585	0.9653	0.2766	0.065*
C2	0.83868 (19)	0.9943 (2)	0.1808 (2)	0.0597 (7)
H2	0.8295	1.0653	0.1783	0.072*
C3	0.89659 (19)	0.9483 (2)	0.12396 (19)	0.0611 (7)
H3	0.9250	0.9899	0.0836	0.073*
C4	0.91234 (18)	0.8453 (2)	0.12618 (18)	0.0543 (7)
H4	0.9516	0.8173	0.0883	0.065*
C5	0.85827 (19)	0.50582 (19)	0.25407 (18)	0.0529 (6)
H5	0.8967	0.4761	0.2162	0.064*
C6	0.8169 (2)	0.4459 (2)	0.3131 (2)	0.0630 (7)
H6	0.8270	0.3750	0.3146	0.076*
C7	0.7591 (2)	0.4880 (2)	0.3722 (2)	0.0615 (7)
H7	0.7319	0.4452	0.4127	0.074*
C8	0.74323 (19)	0.58985 (19)	0.37013 (18)	0.0511 (6)
H8	0.7052	0.6173	0.4098	0.061*
C9	0.76717 (16)	0.76264 (17)	0.30277 (17)	0.0416 (6)
N10	0.88297 (13)	0.67612 (15)	0.18987 (14)	0.0445 (5)
C11	0.80948 (17)	0.82595 (17)	0.24349 (17)	0.0429 (6)

C12	0.86933 (16)	0.78034 (18)	0.18595 (17)	0.0429 (6)
C13	0.78353 (16)	0.65678 (17)	0.30837 (16)	0.0415 (6)
C14	0.84322 (17)	0.61298 (17)	0.24995 (17)	0.0433 (6)
C15	0.70287 (18)	0.80997 (17)	0.3623 (2)	0.0450 (6)
O16	0.61194 (12)	0.81571 (14)	0.29809 (12)	0.0569 (5)
O17	0.72938 (13)	0.83794 (15)	0.45268 (14)	0.0667 (5)
C18	0.54272 (18)	0.8721 (2)	0.33483 (17)	0.0516 (6)
C19	0.48032 (18)	0.8207 (2)	0.37892 (18)	0.0529 (7)
C20	0.4108 (2)	0.8824 (2)	0.4042 (2)	0.0648 (8)
H20	0.3666	0.8514	0.4338	0.078*
C21	0.4053 (2)	0.9861 (3)	0.3872 (2)	0.0688 (8)
H21	0.3582	1.0245	0.4057	0.083*
C22	0.4685 (2)	1.0341 (2)	0.3430 (2)	0.0718 (8)
H22	0.4648	1.1049	0.3313	0.086*
C23	0.5384 (2)	0.9758 (2)	0.3157 (2)	0.0652 (8)
H23	0.5817	1.0070	0.2849	0.078*
C24	0.4878 (2)	0.7061 (2)	0.4025 (2)	0.0665 (8)
H24	0.5264	0.6755	0.3599	0.080*
C25	0.5415 (3)	0.6867 (3)	0.5193 (3)	0.1123 (14)
H25A	0.6038	0.7197	0.5370	0.168*
H25B	0.5500	0.6138	0.5315	0.168*
H25C	0.5047	0.7143	0.5632	0.168*
C26	0.3890 (3)	0.6532 (3)	0.3699 (3)	0.1142 (14)
H26A	0.3566	0.6672	0.2959	0.171*
H26B	0.3506	0.6789	0.4130	0.171*
H26C	0.3976	0.5801	0.3803	0.171*
C27	0.9445 (2)	0.6310 (2)	0.1287 (2)	0.0645 (7)
H27A	0.9342	0.6677	0.0625	0.097*
H27B	0.9277	0.5597	0.1139	0.097*
H27C	1.0114	0.6362	0.1698	0.097*
S28	0.91241 (5)	0.22174 (5)	0.57658 (5)	0.0556 (2)
O29	0.94541 (16)	0.30956 (14)	0.53324 (15)	0.0809 (6)
O30	0.87168 (17)	0.23980 (15)	0.66093 (15)	0.0838 (7)
O31	0.97784 (15)	0.13502 (16)	0.59437 (16)	0.0819 (6)
C32	0.8122 (2)	0.1755 (2)	0.4689 (2)	0.0603 (7)
F33	0.73755 (14)	0.24221 (15)	0.44537 (15)	0.0955 (6)
F34	0.77668 (13)	0.08743 (12)	0.49175 (13)	0.0840 (5)
F35	0.83430 (13)	0.16316 (15)	0.37865 (11)	0.0895 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0500 (17)	0.0489 (16)	0.0643 (15)	0.0028 (13)	0.0195 (14)	-0.0015 (12)
C2	0.0544 (18)	0.0509 (16)	0.0716 (17)	-0.0002 (13)	0.0151 (15)	0.0075 (14)
C3	0.0533 (19)	0.069 (2)	0.0583 (16)	-0.0096 (14)	0.0130 (14)	0.0150 (14)
C4	0.0429 (16)	0.0720 (19)	0.0499 (14)	-0.0010 (13)	0.0166 (12)	0.0044 (13)
C5	0.0505 (17)	0.0518 (16)	0.0532 (14)	0.0114 (13)	0.0101 (13)	-0.0025 (12)
C6	0.073 (2)	0.0457 (16)	0.0642 (16)	0.0109 (14)	0.0110 (15)	0.0028 (13)

C7	0.073 (2)	0.0538 (17)	0.0577 (15)	0.0001 (15)	0.0192 (15)	0.0078 (13)
C8	0.0516 (17)	0.0532 (16)	0.0500 (13)	0.0041 (12)	0.0175 (12)	0.0016 (12)
C9	0.0353 (15)	0.0456 (14)	0.0421 (12)	0.0012 (11)	0.0088 (11)	-0.0043 (10)
N10	0.0327 (12)	0.0512 (13)	0.0496 (11)	0.0048 (9)	0.0122 (10)	-0.0056 (9)
C11	0.0354 (14)	0.0459 (14)	0.0451 (12)	-0.0009 (11)	0.0079 (11)	-0.0033 (10)
C12	0.0313 (14)	0.0520 (16)	0.0419 (12)	0.0005 (11)	0.0057 (11)	-0.0020 (11)
C13	0.0350 (14)	0.0469 (14)	0.0400 (12)	0.0022 (11)	0.0072 (11)	-0.0027 (10)
C14	0.0352 (15)	0.0474 (15)	0.0433 (12)	0.0038 (11)	0.0053 (11)	-0.0011 (10)
C15	0.0448 (17)	0.0431 (14)	0.0491 (14)	0.0012 (11)	0.0168 (13)	-0.0002 (11)
O16	0.0371 (11)	0.0851 (13)	0.0496 (9)	0.0067 (9)	0.0144 (9)	-0.0141 (8)
O17	0.0565 (12)	0.0881 (14)	0.0508 (10)	0.0133 (10)	0.0083 (9)	-0.0208 (9)
C18	0.0415 (16)	0.0713 (18)	0.0416 (13)	0.0115 (13)	0.0117 (12)	-0.0051 (12)
C19	0.0424 (16)	0.0733 (18)	0.0438 (13)	0.0045 (13)	0.0141 (12)	-0.0088 (12)
C20	0.0462 (19)	0.088 (2)	0.0656 (16)	0.0073 (16)	0.0254 (14)	-0.0050 (15)
C21	0.053 (2)	0.092 (2)	0.0599 (17)	0.0258 (17)	0.0146 (15)	-0.0015 (16)
C22	0.073 (2)	0.075 (2)	0.0639 (17)	0.0230 (17)	0.0137 (16)	0.0102 (14)
C23	0.062 (2)	0.081 (2)	0.0571 (16)	0.0124 (16)	0.0234 (15)	0.0095 (14)
C24	0.062 (2)	0.0676 (19)	0.0780 (19)	-0.0062 (15)	0.0340 (16)	-0.0155 (15)
C25	0.169 (4)	0.072 (2)	0.092 (2)	-0.007 (2)	0.032 (3)	0.0148 (18)
C26	0.084 (3)	0.097 (3)	0.177 (4)	-0.025 (2)	0.062 (3)	-0.056 (3)
C27	0.0563 (19)	0.0703 (18)	0.0781 (17)	0.0122 (14)	0.0374 (15)	-0.0050 (14)
S28	0.0708 (5)	0.0511 (4)	0.0508 (4)	-0.0092 (4)	0.0269 (3)	-0.0037 (3)
O29	0.1012 (17)	0.0671 (12)	0.0837 (13)	-0.0311 (11)	0.0416 (13)	0.0000 (10)
O30	0.128 (2)	0.0756 (13)	0.0675 (11)	-0.0128 (12)	0.0596 (13)	-0.0138 (9)
O31	0.0703 (15)	0.0821 (14)	0.0873 (13)	0.0193 (12)	0.0142 (11)	0.0035 (11)
C32	0.061 (2)	0.0637 (18)	0.0655 (17)	0.0012 (15)	0.0326 (16)	0.0020 (13)
F33	0.0703 (13)	0.1043 (14)	0.1154 (14)	0.0249 (11)	0.0328 (11)	0.0228 (11)
F34	0.0793 (13)	0.0644 (11)	0.1129 (13)	-0.0209 (9)	0.0354 (11)	-0.0061 (9)
F35	0.0815 (13)	0.1338 (16)	0.0590 (9)	-0.0108 (11)	0.0300 (9)	-0.0252 (9)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.347 (3)	C18—C19	1.377 (3)
C1—C11	1.422 (3)	C19—C20	1.397 (3)
C1—H1	0.9300	C19—C24	1.515 (4)
C2—C3	1.405 (3)	C20—C21	1.362 (4)
C2—H2	0.9300	C20—H20	0.9300
C3—C4	1.354 (4)	C21—C22	1.365 (4)
C3—H3	0.9300	C21—H21	0.9300
C4—C12	1.412 (3)	C22—C23	1.389 (4)
C4—H4	0.9300	C22—H22	0.9300
C5—C6	1.353 (3)	C23—H23	0.9300
C5—C14	1.405 (3)	C24—C25	1.516 (4)
C5—H5	0.9300	C24—C26	1.527 (4)
C6—C7	1.405 (4)	C24—H24	0.9800
C6—H6	0.9300	C25—H25A	0.9600
C7—C8	1.340 (3)	C25—H25B	0.9600
C7—H7	0.9300	C25—H25C	0.9600

C8—C13	1.422 (3)	C26—H26A	0.9600
C8—H8	0.9300	C26—H26B	0.9600
C9—C11	1.389 (3)	C26—H26C	0.9600
C9—C13	1.391 (3)	C27—H27A	0.9600
C9—C15	1.507 (3)	C27—H27B	0.9600
N10—C12	1.365 (3)	C27—H27C	0.9600
N10—C14	1.373 (3)	S28—O30	1.4148 (17)
N10—C27	1.481 (3)	S28—O29	1.4161 (18)
C11—C12	1.431 (3)	S28—O31	1.4443 (2)
C13—C14	1.428 (3)	S28—C32	1.799 (3)
C15—O17	1.188 (3)	C32—F35	1.321 (3)
C15—O16	1.335 (3)	C32—F34	1.322 (3)
O16—C18	1.431 (3)	C32—F33	1.344 (3)
C18—C23	1.366 (4)		
C2—C1—C11	121.2 (2)	C18—C19—C24	123.0 (2)
C2—C1—H1	119.4	C20—C19—C24	121.8 (2)
C11—C1—H1	119.4	C21—C20—C19	122.6 (3)
C1—C2—C3	119.5 (2)	C21—C20—H20	118.7
C1—C2—H2	120.2	C19—C20—H20	118.7
C3—C2—H2	120.2	C20—C21—C22	120.3 (3)
C4—C3—C2	122.0 (2)	C20—C21—H21	119.9
C4—C3—H3	119.0	C22—C21—H21	119.9
C2—C3—H3	119.0	C21—C22—C23	119.2 (3)
C3—C4—C12	120.1 (2)	C21—C22—H22	120.4
C3—C4—H4	119.9	C23—C22—H22	120.4
C12—C4—H4	119.9	C18—C23—C22	119.0 (3)
C6—C5—C14	120.1 (2)	C18—C23—H23	120.5
C6—C5—H5	120.0	C22—C23—H23	120.5
C14—C5—H5	120.0	C19—C24—C25	110.7 (2)
C5—C6—C7	121.7 (2)	C19—C24—C26	112.3 (3)
C5—C6—H6	119.2	C25—C24—C26	111.4 (3)
C7—C6—H6	119.2	C19—C24—H24	107.4
C8—C7—C6	119.9 (2)	C25—C24—H24	107.4
C8—C7—H7	120.1	C26—C24—H24	107.4
C6—C7—H7	120.1	C24—C25—H25A	109.5
C7—C8—C13	121.1 (2)	C24—C25—H25B	109.5
C7—C8—H8	119.4	H25A—C25—H25B	109.5
C13—C8—H8	119.4	C24—C25—H25C	109.5
C11—C9—C13	121.1 (2)	H25A—C25—H25C	109.5
C11—C9—C15	119.2 (2)	H25B—C25—H25C	109.5
C13—C9—C15	119.70 (19)	C24—C26—H26A	109.5
C12—N10—C14	122.16 (18)	C24—C26—H26B	109.5
C12—N10—C27	118.31 (19)	H26A—C26—H26B	109.5
C14—N10—C27	119.5 (2)	C24—C26—H26C	109.5
C9—C11—C1	122.5 (2)	H26A—C26—H26C	109.5
C9—C11—C12	118.9 (2)	H26B—C26—H26C	109.5
C1—C11—C12	118.6 (2)	N10—C27—H27A	109.5

N10—C12—C4	122.0 (2)	N10—C27—H27B	109.5
N10—C12—C11	119.5 (2)	H27A—C27—H27B	109.5
C4—C12—C11	118.5 (2)	N10—C27—H27C	109.5
C9—C13—C8	122.7 (2)	H27A—C27—H27C	109.5
C9—C13—C14	119.0 (2)	H27B—C27—H27C	109.5
C8—C13—C14	118.3 (2)	O30—S28—O29	116.47 (12)
N10—C14—C5	121.7 (2)	O30—S28—O31	113.97 (13)
N10—C14—C13	119.3 (2)	O29—S28—O31	114.20 (13)
C5—C14—C13	118.9 (2)	O30—S28—C32	104.08 (13)
O17—C15—O16	125.3 (2)	O29—S28—C32	104.00 (12)
O17—C15—C9	124.9 (2)	O31—S28—C32	101.75 (13)
O16—C15—C9	109.79 (19)	F35—C32—F34	108.1 (2)
C15—O16—C18	118.12 (17)	F35—C32—F33	105.2 (2)
C23—C18—C19	123.6 (2)	F34—C32—F33	105.7 (2)
C23—C18—O16	116.1 (2)	F35—C32—S28	112.96 (19)
C19—C18—O16	120.1 (2)	F34—C32—S28	112.72 (19)
C18—C19—C20	115.2 (2)	F33—C32—S28	111.7 (2)
C11—C1—C2—C3	-0.2 (4)	C8—C13—C14—N10	-179.9 (2)
C1—C2—C3—C4	-0.5 (4)	C9—C13—C14—C5	-178.8 (2)
C2—C3—C4—C12	0.6 (4)	C8—C13—C14—C5	1.2 (3)
C14—C5—C6—C7	-0.6 (4)	C11—C9—C15—O17	-92.5 (3)
C5—C6—C7—C8	0.5 (4)	C13—C9—C15—O17	87.2 (3)
C6—C7—C8—C13	0.4 (4)	C11—C9—C15—O16	87.4 (2)
C13—C9—C11—C1	-178.0 (2)	C13—C9—C15—O16	-92.8 (2)
C15—C9—C11—C1	1.7 (3)	O17—C15—O16—C18	8.9 (3)
C13—C9—C11—C12	1.2 (3)	C9—C15—O16—C18	-171.01 (19)
C15—C9—C11—C12	-179.1 (2)	C15—O16—C18—C23	86.1 (3)
C2—C1—C11—C9	179.9 (2)	C15—O16—C18—C19	-98.8 (3)
C2—C1—C11—C12	0.7 (4)	C23—C18—C19—C20	-0.3 (4)
C14—N10—C12—C4	178.4 (2)	O16—C18—C19—C20	-175.1 (2)
C27—N10—C12—C4	-0.4 (3)	C23—C18—C19—C24	-178.2 (2)
C14—N10—C12—C11	-1.9 (3)	O16—C18—C19—C24	7.0 (4)
C27—N10—C12—C11	179.4 (2)	C18—C19—C20—C21	-0.4 (4)
C3—C4—C12—N10	179.7 (2)	C24—C19—C20—C21	177.5 (3)
C3—C4—C12—C11	-0.1 (3)	C19—C20—C21—C22	0.6 (4)
C9—C11—C12—N10	0.5 (3)	C20—C21—C22—C23	0.0 (4)
C1—C11—C12—N10	179.7 (2)	C19—C18—C23—C22	0.8 (4)
C9—C11—C12—C4	-179.8 (2)	O16—C18—C23—C22	175.8 (2)
C1—C11—C12—C4	-0.6 (3)	C21—C22—C23—C18	-0.6 (4)
C11—C9—C13—C8	178.6 (2)	C18—C19—C24—C25	97.9 (3)
C15—C9—C13—C8	-1.2 (3)	C20—C19—C24—C25	-79.8 (3)
C11—C9—C13—C14	-1.4 (3)	C18—C19—C24—C26	-137.0 (3)
C15—C9—C13—C14	178.9 (2)	C20—C19—C24—C26	45.3 (3)
C7—C8—C13—C9	178.8 (2)	O30—S28—C32—F35	174.13 (19)
C7—C8—C13—C14	-1.2 (3)	O29—S28—C32—F35	51.7 (2)
C12—N10—C14—C5	-179.5 (2)	O31—S28—C32—F35	-67.2 (2)
C27—N10—C14—C5	-0.8 (3)	O30—S28—C32—F34	-63.0 (2)

C12—N10—C14—C13	1.6 (3)	O29—S28—C32—F34	174.54 (18)
C27—N10—C14—C13	−179.6 (2)	O31—S28—C32—F34	55.6 (2)
C6—C5—C14—N10	−179.2 (2)	O30—S28—C32—F33	55.7 (2)
C6—C5—C14—C13	−0.3 (3)	O29—S28—C32—F33	−66.7 (2)
C9—C13—C14—N10	0.0 (3)	O31—S28—C32—F33	174.42 (17)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C4—H4···O29 ⁱ	0.93	2.48	3.363 (3)	159
C27—H27C···F35 ⁱ	0.96	2.51	3.250 (4)	134

Symmetry code: (i) $-x+2, y+1/2, -z+1/2$.