

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Piperazine-2,3,5,6-tetraone. Corrigendum

Jing-Jing Jia,^a Xiu-Jin Meng,^a Shi-Zhang Liang,^a Shu-Hua Zhang^b and Yi-Min Jiang^{a*}

^aCollege of Chemistry and Chemical Engineering, Guangxi Normal University, Guilin, Guangxi 541004, People's Republic of China, and ^bCollege of Chemistry and Bioengineering, Guilin University of Technology, Guilin, Guangxi 541004, People's Republic of China

Correspondence e-mail: 19768069@qq.com

Received 14 December 2010; accepted 15 December 2010

The correspondence author in the paper by Jia *et al.* [*Acta Cryst.* (2010), **E66**, o3315] is corrected.

In the paper by Jia *et al.* (2010), the correspondence author is incorrectly given as 'Jing-Jing Jia'. The correct correspondence author is 'Yi-Min Jiang', as given above.

References

Jia, J.-J., Meng, X.-J., Liang, S.-Z., Zhang, S.-H. & Jiang, Y.-M. (2010). *Acta Cryst.* **E66**, o3315.

Piperazine-2,3,5,6-tetraone

Jing-Jing Jia,^{a*} Xiu-Jin Meng,^a Shi-Zhang Liang,^a Shu-Hua Zhang^b and Yi-Min Jiang^a^aCollege of Chemistry and Chemical Engineering, Guangxi Normal University, Guilin, Guangxi 541004, People's Republic of China, and ^bCollege of Chemistry and Bioengineering, Guilin University of Technology, Guilin, Guangxi 541004, People's Republic of China

Correspondence e-mail: 19768069@qq.com


Received 2 November 2010; accepted 18 November 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.048; wR factor = 0.092; data-to-parameter ratio = 9.5.

The molecule of the title compound, $\text{C}_4\text{H}_2\text{N}_2\text{O}_4$, is located around an inversion center and the four O atoms are in the 2,3,5,6-positions of the piperazine ring. In the crystal, bifurcated $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into a corrugated layer parallel to (101).

Related literature

For the synthesis of tetraone, see: Norcross *et al.* (2008). For related structures, see Sletten *et al.* (1970, 1980); Sarangarajan *et al.* (2005); Norcross *et al.* (2008); Jin *et al.* (1998); Sanner *et al.* (1992); Ongania *et al.* (1985).



Experimental

Crystal data

$\text{C}_4\text{H}_2\text{N}_2\text{O}_4$
 $M_r = 142.08$
 Monoclinic, $P2_1/n$
 $a = 5.163$ (1) Å
 $b = 8.6220$ (17) Å
 $c = 5.6540$ (11) Å
 $\beta = 105.25$ (3)°

$V = 242.83$ (8) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.18$ mm⁻¹
 $T = 293$ K
 $0.42 \times 0.32 \times 0.12$ mm

Data collection

Siemens P4 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.930$, $T_{\max} = 0.978$

1357 measured reflections
 438 independent reflections
 383 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.092$
 $S = 1.23$
 438 reflections

46 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.20$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.21$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}^i$	0.86	2.48	3.060 (2)	125
$\text{N1}-\text{H1}\cdots\text{O2}^{ii}$	0.86	2.23	3.035 (2)	157

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: XSCANS (Siemens, 1994); cell refinement: XSCANS; data reduction: XSCANS; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEPIII (Burnett & Johnson, 1996), ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: SHELXTL (Sheldrick, 2008).

This work was funded by the Guangxi Science Foundation, Guangxi Zhuang Autonomous Region of the People's Republic of China (grant No. 2010GXNSFD013017).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2617).

References

- Burnett, M. N. & Johnson, C. K. (1996). ORTEPIII. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Jin, G. Z., Kim, Y., Chung, J. H., Sok, D. E. & Ahn, B. Z. (1998). *Arch. Pharm. (Weinheim)*, **331**, 380–4.
 Norcross, N. R., Melbardis, J. P., Solera, M. F., Sephton, M. A., Kilner, C., Zakharov, L. N., Astles, P. C., Warriner, S. L. & Blakemore, P. R. (2008). *J. Org. Chem.* **73**, 7939–7951.
 Ongania, K. H., Granozzi, G., Busetti, V., Casarin, M. & Ajò, D. (1985). *Tetrahedron*, **41**, 2015–2018.
 Sanner, M. A., Weigelt, C., Stansberry, M., Killeen, K., Michne, W. F., Kessler, D. W. & Kullnig, R. K. (1992). *J. Org. Chem.* **57**, 5264–5268.
 Sarangarajan, T. R., Panchanatheswaran, K., Low, J. N. & Glidewell, C. (2005). *Acta Cryst.* **C61**, o118–o121.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Siemens (1994). XSCANS. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Sletten, E. (1970). *J. Am. Chem. Soc.* **92**, 172–177.
 Sletten, J. (1980). *Acta Chem. Scand. Ser. A*, **34**, 593–595.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2010). E66, o3315 [https://doi.org/10.1107/S1600536810048075]

Piperazine-2,3,5,6-tetraone

Jing-Jing Jia, Xiu-Jin Meng, Shi-Zhang Liang, Shu-Hua Zhang and Yi-Min Jiang

S1. Comment

The synthesis and antitumor activity of some tetraone compounds have been widely studied (Jin *et al.*, 1998; Sanner *et al.*, 1992). Most tetraone compounds were found from a naturally occurring alkaloid in a variety of leguminous plant and tree species, including broom, lupin, gorse, and laburnum (Norcross *et al.*, 2008). As part of our interest in the synthesis of tetraone derivatives, we report here the structure of the title compound.

The molecule of the title compound is located around inversion center and the four O atoms are in the 2,3,5,6 position on the piperazine ring (Fig. 1). The molecule is planar with rms deviation of 0.013 Å. The bond distances and angles are similar to those found in related piperazine derivatives (Sletten *et al.*, 1970; Sarangarajan *et al.*, 2005; Sletten *et al.*, 1980; Ongania *et al.*, 1985).

The N—H donor and the C—O acceptor groups participate in the hydrogen bonding forming corrugated layers parallel to the (1 0 1) plane through bifurcated N-H...O hydrogen bonds (Table 1, Fig. 2).

S2. Experimental

For the preparation of the title compound, the 2-mercaptopyrazine (10 mmol, 1.1200 g) was dissolved in ethanol (50 ml) at 358 K and a solution of 30% H₂O₂ (10 ml) was added. The resulting solution was stirred at 358 K for 4 h, then concentrating at 388 K, until 3 ml solution remained. Colourless-block crystal suitable for X-ray diffraction were obtained by slow evaporation at room temperature after several days in 55% yield.

S3. Refinement

H atom attached to N atom was positioned geometrically and treated as riding on the parent atom with N-H = 0.86 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$.




Figure 1

Molecular view of compound I with the atom labeling scheme. Ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii. [Symmetry code: (i) $-x+1, -y+1, -z+1$]




Figure 2

Partial packing view showing the corrugated layer parallel to the (1 0 1) plane. H bonds are shown as dashed lines.

[Symmetry codes: (ii) $-x+1/2, y-1/2, -z+3/2$; (iii) $x-1/2, -y+1/2, z+1/2$]

Piperazine-2,3,5,6-tetraone

Crystal data

$C_4H_2N_2O_4$
 $M_r = 142.08$
 Monoclinic, $P2_1/n$
 Hall symbol: $-P\ 2_1n$
 $a = 5.163\ (1)\ \text{\AA}$
 $b = 8.6220\ (17)\ \text{\AA}$
 $c = 5.6540\ (11)\ \text{\AA}$
 $\beta = 105.25\ (3)^\circ$
 $V = 242.83\ (8)\ \text{\AA}^3$
 $Z = 2$

$F(000) = 144$
 $D_x = 1.943\ \text{Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$
 Cell parameters from 438 reflections
 $\theta = 4.4\text{--}25.3^\circ$
 $\mu = 0.18\ \text{mm}^{-1}$
 $T = 293\ \text{K}$
 Block, colourless
 $0.42 \times 0.32 \times 0.12\ \text{mm}$

Data collection

Siemens P4
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan
 (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.930, T_{\max} = 0.978$

1357 measured reflections
 438 independent reflections
 383 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$
 $\theta_{\max} = 25.3^\circ, \theta_{\min} = 4.4^\circ$
 $h = -6 \rightarrow 6$
 $k = -10 \rightarrow 10$
 $l = -6 \rightarrow 6$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.048$	H-atom parameters constrained
$wR(F^2) = 0.092$	$w = 1/[\sigma^2(F_o^2) + (0.0355P)^2 + 0.1147P]$
$S = 1.23$	where $P = (F_o^2 + 2F_c^2)/3$
438 reflections	$(\Delta/\sigma)_{\max} < 0.001$
46 parameters	$\Delta\rho_{\max} = 0.20 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.1613 (3)	0.55814 (19)	0.7646 (3)	0.0343 (5)
O2	0.8069 (3)	0.25100 (19)	0.6060 (3)	0.0342 (5)
N1	0.4775 (4)	0.3995 (2)	0.6856 (3)	0.0261 (5)
H1	0.4603	0.3362	0.7981	0.031*
C1	0.6621 (5)	0.3635 (2)	0.5614 (4)	0.0233 (5)
C2	0.3169 (4)	0.5281 (2)	0.6461 (4)	0.0233 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0357 (10)	0.0355 (10)	0.0373 (10)	0.0010 (8)	0.0197 (9)	-0.0034 (8)
O2	0.0348 (10)	0.0268 (9)	0.0402 (10)	0.0099 (8)	0.0086 (8)	0.0048 (7)
N1	0.0330 (11)	0.0229 (10)	0.0248 (11)	0.0004 (9)	0.0120 (9)	0.0053 (8)
C1	0.0221 (12)	0.0205 (11)	0.0256 (12)	-0.0019 (10)	0.0034 (10)	-0.0017 (9)
C2	0.0224 (12)	0.0225 (11)	0.0239 (12)	-0.0027 (10)	0.0042 (10)	-0.0038 (9)

Geometric parameters (\AA , $^\circ$)

O1—C2	1.202 (3)	N1—C2	1.368 (3)
O2—C1	1.210 (3)	N1—H1	0.8600
N1—C1	1.360 (3)	C1—C2 ⁱ	1.526 (3)
C1—N1—C2	125.31 (19)	N1—C1—C2 ⁱ	117.28 (19)
C1—N1—H1	117.3	O1—C2—N1	123.3 (2)
C2—N1—H1	117.3	O1—C2—C1 ⁱ	119.3 (2)

O2—C1—N1	123.6 (2)	N1—C2—C1 ⁱ	117.35 (18)
O2—C1—C2 ⁱ	119.10 (19)		

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1...O1 ⁱⁱ	0.86	2.48	3.060 (2)	125
N1—H1...O2 ⁱⁱⁱ	0.86	2.23	3.035 (2)	157

Symmetry codes: (ii) $-x+1/2, y-1/2, -z+3/2$; (iii) $x-1/2, -y+1/2, z+1/2$.