

7-Bromo-1-(4-fluorophenylsulfonyl)-2-methylnaphtho[2,1-*b*]furan

Hong Dae Choi,^a Pil Ja Seo,^a Byeng Wha Son^b and Uk Lee^{b*}

^aDepartment of Chemistry, Dongeui University, San 24 Kaya-dong Busanjin-gu, Busan 614-714, Republic of Korea, and ^bDepartment of Chemistry, Pukyong National University, 599-1 Daeyeon 3-dong, Nam-gu, Busan 608-737, Republic of Korea
Correspondence e-mail: uklee@pknu.ac.kr


Received 9 December 2010; accepted 26 December 2010

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.031; wR factor = 0.078; data-to-parameter ratio = 16.8.

In the title compound, $\text{C}_{19}\text{H}_{12}\text{BrFO}_3\text{S}$, the 4-fluorophenyl ring makes a dihedral angle of $80.32(5)^\circ$ with the mean plane of the naphthofuran fragment. In the crystal, molecules are linked by weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\pi$ interactions. The crystal structure also exhibits aromatic $\pi-\pi$ interactions between the central benzene rings of neighbouring molecules [centroid–centroid distance = $3.564(3)\text{ \AA}$].

Related literature

For the pharmacological activity of naphthofuran compounds, see: Einhorn *et al.* (1984); Hranjec *et al.* (2003); Mahadevan & Vaidya (2003). For our previous structural studies of related 1-arylsulfonyl-7-bromo-2-methylnaphtho[2,1-*b*]furan derivatives, see: Choi *et al.* (2008a,b).



Experimental

Crystal data

$\text{C}_{19}\text{H}_{12}\text{BrFO}_3\text{S}$

$M_r = 419.26$

Monoclinic, $P2_1/n$
 $a = 12.0415(2)\text{ \AA}$
 $b = 8.1579(1)\text{ \AA}$
 $c = 17.4578(3)\text{ \AA}$
 $\beta = 105.325(1)^\circ$
 $V = 1653.96(5)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 2.64\text{ mm}^{-1}$
 $T = 173\text{ K}$
 $0.25 \times 0.21 \times 0.12\text{ mm}$

Data collection

Bruker SMART APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2009)
 $S = 0.553$, $T_{\max} = 0.744$

14803 measured reflections
3803 independent reflections
3026 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.078$
 $S = 1.02$
3803 reflections

227 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.42\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.41\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$ is the centroid of the C14–C19 4-fluorophenyl ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$C5-\text{H}5\cdots O2^i$	0.95	2.57	3.312 (3)	135
$C18-\text{H}18\cdots O2^{ii}$	0.95	2.36	3.266 (3)	160
$C19-\text{H}19\cdots O3^i$	0.95	2.57	3.113 (2)	117
$C10-\text{H}10\cdots Cg1^{iii}$	0.95	2.69	3.609 (2)	152

Symmetry codes: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, y + 1, z$; (iii) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 1998); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2255).

References

- Brandenburg, K. (1998). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2009). *APEX2*, *SADABS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2008a). *Acta Cryst. E64*, o944.
- Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2008b). *Acta Cryst. E64*, o1158.
- Einhorn, J., Demerseman, P., Royer, R., Cavier, R. & Gayral, P. (1984). *Eur. J. Med. Chem.* **19**, 405–410.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Hranjec, M., Grdisa, M., Pavelic, K., Boykin, D. W. & Karminski-Zamola, G. (2003). *Farmac.* **58**, 1319–1324.
- Mahadevan, K. M. & Vaidya, V. P. (2003). *Indian J. Pharm. Sci.* **65**, 128–134.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

supporting information

Acta Cryst. (2011). E67, o280 [doi:10.1107/S160053681005436X]

7-Bromo-1-(4-fluorophenylsulfonyl)-2-methylnaphtho[2,1-*b*]furan

Hong Dae Choi, Pil Ja Seo, Byeng Wha Son and Uk Lee

S1. Comment

Many compounds involving a naphthofuran ring have attracted much attention in view of their diverse pharmacological properties such as antibacterial, antitumor and anthelmintic activities (Einhorn *et al.*, 1984, Hranjec *et al.*, 2003, Mahadevan & Vaidya, 2003). As a part of our ongoing studies of the substituent effect on the solid state structures of 1-arylsulfonyl-7-bromo-2-methylnaphtho[2,1-*b*]furan analogues (Choi *et al.*, 2008*a,b*), we report herein on the crystal structure of the title compound.


In the title molecule (Fig. 1), the naphthofuran moiety is essentially planar, with a mean deviation of 0.011 (2) Å from the least-squares plane defined by the thirteen constituent atoms. The dihedral angle formed by the mean plane of the naphthofuran system and the 4-fluorophenyl ring is 80.32 (5)°. The crystal packing (Fig. 2) is stabilized by weak intermolecular C–H···O hydrogen bonds; the first one between a benzene H atom and the oxygen of the O=S=O unit (Table 1; C5–H5···O2ⁱ), and the second one between the 4-fluorophenyl H atom and the oxygen of the O=S=O unit (Table 1; C18–H18···O2ⁱⁱ), and the third one between the 4-fluorophenyl H and the oxygen of the O=S=O unit (Table 1; C19–H19···O3ⁱ). The molecular packing (Fig. 3) is further stabilized by an intermolecular C–H···π interaction between a benzene H atom and the 4-fluorophenyl ring (Table 1; C10–H10···Cg1ⁱⁱⁱ, Cg1 is the centroid of the C14–C19 4-fluorophenyl ring). In addition, the crystal packing (Fig. 3) exhibits an aromatic π–π interaction between the central benzene rings of neighbouring molecules. The Cg2···Cg2^{vi} distance is 3.564 (3) Å (Cg2 is the centroid of the C2/C3/C8–C11 benzene ring).

S2. Experimental


3-Chloroperoxybenzoic acid (77%) (404 mg, 1.8 mmol) was added in small portions to a stirred solution of 7-bromo-1-(4-fluorophenylsulfonyl)-2-methylnaphtho[2,1-*b*]furan (348 mg, 0.9 mmol) in dichloromethane (40 mL) at 273 K. After being stirred at room temperature for 10 h, the mixture was washed with saturated sodium bicarbonate solution and the organic layer was separated, dried over magnesium sulfate, filtered and concentrated at reduced pressure. The residue was purified by column chromatography (hexane–ethyl acetate, 4:1 v/v) to afford the title compound as a colourless solid [yield 73%, m.p. 485–486 K; R_f = 0.52 (hexane–ethyl acetate, 4:1 v/v)]. Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of the title compound in acetone at room temperature.

S3. Refinement


All H atoms were positioned geometrically and refined using a riding model, with C–H = 0.95 Å for aryl and 0.98 Å for methyl H atoms. $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aryl and $1.5U_{\text{eq}}(\text{C})$ for methyl H atoms.

**Figure 1**

The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as a small spheres of arbitrary radius.

**Figure 2**

A view of C–H \cdots O interactions (dotted lines) in the crystal structure of the title compound. Symmetry codes: (i) $-x+1/2$, $y+1/2$, $-z+1/2$; (ii) x , $y+1$, z ; (iv) $-x+1/2$, $y-1/2$, $-z+1/2$; (v) x , $y-1$, z .

**Figure 3**

A view of C–H \cdots π and π – π interactions (dotted lines) in the crystal structure of the title compound. Symmetry codes: (iii) $x+1/2$, $-y+1/2$, $z+1/2$; (vi) $-x+1$, $-y$, $-z + 1$; (vii) $x-1/2$, $-y+1/2$, $z-1/2$.

7-Bromo-1-(4-fluorophenylsulfonyl)-2-methylnaphtho[2,1-*b*]furan

Crystal data

$C_{19}H_{12}BrFO_3S$
 $M_r = 419.26$
Monoclinic, $P2_1/n$

Hall symbol: -P 2yn
 $a = 12.0415 (2)$ Å
 $b = 8.1579 (1)$ Å

$c = 17.4578 (3)$ Å
 $\beta = 105.325 (1)^\circ$
 $V = 1653.96 (5)$ Å³
 $Z = 4$
 $F(000) = 840$
 $D_x = 1.684$ Mg m⁻³
 Melting point = 485–486 K

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 4448 reflections
 $\theta = 2.4\text{--}27.4^\circ$
 $\mu = 2.64$ mm⁻¹
 $T = 173$ K
 Block, colourless
 $0.25 \times 0.21 \times 0.12$ mm

Data collection

Bruker SMART APEXII CCD
 diffractometer
 Radiation source: rotating anode
 Graphite multilayer monochromator
 Detector resolution: 10.0 pixels mm⁻¹
 φ and ω scans
 Absorption correction: multi-scan
 (*SADABS*; Bruker, 2009)
 $T_{\min} = 0.553$, $T_{\max} = 0.744$

14803 measured reflections
 3803 independent reflections
 3026 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$
 $\theta_{\max} = 27.6^\circ$, $\theta_{\min} = 1.9^\circ$
 $h = -15 \rightarrow 15$
 $k = -10 \rightarrow 10$
 $l = -22 \rightarrow 22$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.078$
 $S = 1.02$
 3803 reflections
 227 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0385P)^2 + 0.6504P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.42$ e Å⁻³
 $\Delta\rho_{\min} = -0.41$ e Å⁻³

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.23409 (2)	0.55058 (3)	0.595181 (15)	0.04881 (10)
S1	0.40951 (4)	0.07887 (6)	0.25320 (3)	0.02349 (12)
F1	0.52925 (15)	0.6513 (2)	0.09244 (10)	0.0670 (5)
O1	0.68915 (11)	0.0164 (2)	0.42360 (8)	0.0308 (3)
O2	0.42383 (14)	-0.06341 (18)	0.20874 (9)	0.0363 (4)
O3	0.29924 (11)	0.1077 (2)	0.26702 (8)	0.0316 (3)
C1	0.51536 (16)	0.0748 (2)	0.34372 (11)	0.0240 (4)
C2	0.52174 (16)	0.1490 (3)	0.42090 (10)	0.0234 (4)
C3	0.44999 (16)	0.2428 (2)	0.45819 (10)	0.0236 (4)

C4	0.33609 (16)	0.2927 (3)	0.42204 (11)	0.0285 (4)
H4	0.3021	0.2637	0.3682	0.034*
C5	0.27304 (18)	0.3815 (3)	0.46198 (12)	0.0327 (5)
H5	0.1963	0.4132	0.4363	0.039*
C6	0.32294 (19)	0.4253 (3)	0.54121 (13)	0.0322 (5)
C7	0.43192 (19)	0.3815 (3)	0.57901 (12)	0.0329 (5)
H7	0.4638	0.4137	0.6326	0.040*
C8	0.49895 (17)	0.2882 (3)	0.53949 (11)	0.0277 (4)
C9	0.61271 (18)	0.2411 (3)	0.58063 (11)	0.0331 (5)
H9	0.6429	0.2747	0.6342	0.040*
C10	0.67917 (17)	0.1498 (3)	0.54546 (12)	0.0323 (5)
H10	0.7547	0.1165	0.5731	0.039*
C11	0.63028 (16)	0.1071 (3)	0.46594 (11)	0.0269 (4)
C12	0.61837 (17)	-0.0014 (3)	0.34912 (12)	0.0280 (4)
C13	0.6701 (2)	-0.0921 (3)	0.29391 (14)	0.0407 (6)
H13A	0.7321	-0.0267	0.2826	0.061*
H13B	0.6110	-0.1136	0.2443	0.061*
H13C	0.7014	-0.1964	0.3182	0.061*
C14	0.44463 (15)	0.2512 (3)	0.20340 (10)	0.0219 (4)
C15	0.49975 (17)	0.2294 (3)	0.14369 (11)	0.0327 (5)
H15	0.5180	0.1225	0.1292	0.039*
C16	0.5279 (2)	0.3660 (4)	0.10549 (13)	0.0423 (6)
H16	0.5651	0.3549	0.0641	0.051*
C17	0.5005 (2)	0.5172 (3)	0.12903 (14)	0.0400 (6)
C18	0.44494 (19)	0.5411 (3)	0.18684 (13)	0.0344 (5)
H18	0.4264	0.6484	0.2007	0.041*
C19	0.41645 (17)	0.4054 (3)	0.22448 (11)	0.0268 (4)
H19	0.3775	0.4180	0.2649	0.032*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.06439 (18)	0.04182 (17)	0.05384 (16)	-0.00541 (13)	0.03959 (13)	-0.00784 (12)
S1	0.0259 (2)	0.0209 (3)	0.0210 (2)	-0.00233 (19)	0.00152 (17)	-0.00073 (19)
F1	0.0806 (11)	0.0562 (11)	0.0654 (10)	-0.0212 (9)	0.0216 (9)	0.0257 (9)
O1	0.0236 (7)	0.0369 (9)	0.0295 (7)	0.0027 (6)	0.0025 (6)	0.0064 (7)
O2	0.0475 (9)	0.0228 (8)	0.0324 (7)	0.0016 (7)	0.0001 (7)	-0.0057 (7)
O3	0.0245 (7)	0.0399 (9)	0.0282 (7)	-0.0059 (7)	0.0031 (5)	0.0003 (7)
C1	0.0260 (9)	0.0230 (11)	0.0215 (8)	-0.0011 (8)	0.0035 (7)	0.0048 (8)
C2	0.0254 (9)	0.0204 (10)	0.0220 (8)	-0.0053 (8)	0.0019 (7)	0.0044 (8)
C3	0.0282 (9)	0.0200 (10)	0.0208 (8)	-0.0053 (8)	0.0034 (7)	0.0045 (8)
C4	0.0303 (10)	0.0307 (12)	0.0235 (9)	-0.0016 (9)	0.0054 (8)	0.0011 (9)
C5	0.0320 (10)	0.0350 (13)	0.0316 (10)	0.0018 (10)	0.0095 (8)	0.0037 (10)
C6	0.0453 (12)	0.0250 (12)	0.0328 (10)	-0.0040 (10)	0.0217 (9)	-0.0015 (9)
C7	0.0460 (12)	0.0310 (12)	0.0233 (9)	-0.0120 (10)	0.0116 (9)	-0.0008 (9)
C8	0.0342 (10)	0.0240 (11)	0.0235 (9)	-0.0085 (9)	0.0049 (8)	0.0018 (9)
C9	0.0374 (11)	0.0355 (13)	0.0210 (9)	-0.0114 (10)	-0.0017 (8)	0.0019 (9)
C10	0.0262 (10)	0.0376 (13)	0.0263 (9)	-0.0054 (9)	-0.0048 (8)	0.0076 (10)

C11	0.0258 (9)	0.0260 (11)	0.0269 (9)	-0.0033 (8)	0.0037 (8)	0.0058 (9)
C12	0.0286 (10)	0.0283 (11)	0.0252 (9)	-0.0005 (9)	0.0036 (8)	0.0062 (9)
C13	0.0378 (12)	0.0487 (16)	0.0375 (12)	0.0106 (11)	0.0131 (9)	0.0027 (11)
C14	0.0200 (8)	0.0243 (11)	0.0187 (8)	0.0015 (8)	0.0004 (6)	0.0014 (8)
C15	0.0360 (11)	0.0347 (13)	0.0292 (10)	0.0052 (10)	0.0116 (9)	-0.0006 (10)
C16	0.0404 (12)	0.0553 (18)	0.0359 (11)	0.0010 (12)	0.0183 (10)	0.0100 (12)
C17	0.0396 (12)	0.0381 (14)	0.0387 (12)	-0.0108 (11)	0.0040 (10)	0.0151 (11)
C18	0.0394 (12)	0.0222 (12)	0.0366 (11)	-0.0024 (10)	0.0012 (9)	0.0004 (10)
C19	0.0283 (10)	0.0259 (11)	0.0242 (9)	0.0014 (9)	0.0036 (7)	-0.0016 (9)

Geometric parameters (\AA , $^\circ$)

Br1—C6	1.900 (2)	C7—H7	0.9500
S1—O3	1.4303 (14)	C8—C9	1.421 (3)
S1—O2	1.4319 (16)	C9—C10	1.354 (3)
S1—C1	1.7487 (18)	C9—H9	0.9500
S1—C14	1.762 (2)	C10—C11	1.401 (3)
F1—C17	1.357 (3)	C10—H10	0.9500
O1—C12	1.361 (2)	C12—C13	1.477 (3)
O1—C11	1.369 (3)	C13—H13A	0.9800
C1—C12	1.368 (3)	C13—H13B	0.9800
C1—C2	1.461 (3)	C13—H13C	0.9800
C2—C11	1.378 (3)	C14—C19	1.378 (3)
C2—C3	1.433 (3)	C14—C15	1.387 (3)
C3—C4	1.410 (3)	C15—C16	1.386 (3)
C3—C8	1.434 (3)	C15—H15	0.9500
C4—C5	1.367 (3)	C16—C17	1.367 (4)
C4—H4	0.9500	C16—H16	0.9500
C5—C6	1.401 (3)	C17—C18	1.364 (3)
C5—H5	0.9500	C18—C19	1.376 (3)
C6—C7	1.352 (3)	C18—H18	0.9500
C7—C8	1.415 (3)	C19—H19	0.9500
O3—S1—O2	118.34 (10)	C9—C10—C11	116.25 (18)
O3—S1—C1	109.76 (9)	C9—C10—H10	121.9
O2—S1—C1	108.13 (9)	C11—C10—H10	121.9
O3—S1—C14	107.86 (9)	O1—C11—C2	112.00 (17)
O2—S1—C14	107.47 (9)	O1—C11—C10	121.84 (18)
C1—S1—C14	104.37 (9)	C2—C11—C10	126.2 (2)
C12—O1—C11	106.89 (15)	O1—C12—C1	110.17 (18)
C12—C1—C2	107.40 (16)	O1—C12—C13	114.29 (17)
C12—C1—S1	120.12 (15)	C1—C12—C13	135.52 (19)
C2—C1—S1	132.37 (15)	C12—C13—H13A	109.5
C11—C2—C3	117.91 (17)	C12—C13—H13B	109.5
C11—C2—C1	103.54 (17)	H13A—C13—H13B	109.5
C3—C2—C1	138.54 (17)	C12—C13—H13C	109.5
C4—C3—C2	125.66 (17)	H13A—C13—H13C	109.5
C4—C3—C8	117.73 (18)	H13B—C13—H13C	109.5

C2—C3—C8	116.61 (17)	C19—C14—C15	121.19 (19)
C5—C4—C3	122.05 (18)	C19—C14—S1	119.23 (14)
C5—C4—H4	119.0	C15—C14—S1	119.57 (17)
C3—C4—H4	119.0	C16—C15—C14	119.0 (2)
C4—C5—C6	119.2 (2)	C16—C15—H15	120.5
C4—C5—H5	120.4	C14—C15—H15	120.5
C6—C5—H5	120.4	C17—C16—C15	118.1 (2)
C7—C6—C5	121.5 (2)	C17—C16—H16	120.9
C7—C6—Br1	120.08 (16)	C15—C16—H16	120.9
C5—C6—Br1	118.42 (17)	F1—C17—C18	117.9 (2)
C6—C7—C8	120.63 (19)	F1—C17—C16	118.3 (2)
C6—C7—H7	119.7	C18—C17—C16	123.8 (2)
C8—C7—H7	119.7	C17—C18—C19	118.1 (2)
C7—C8—C9	119.80 (18)	C17—C18—H18	121.0
C7—C8—C3	118.90 (18)	C19—C18—H18	121.0
C9—C8—C3	121.30 (19)	C18—C19—C14	119.79 (19)
C10—C9—C8	121.73 (18)	C18—C19—H19	120.1
C10—C9—H9	119.1	C14—C19—H19	120.1
C8—C9—H9	119.1		
O3—S1—C1—C12	158.70 (17)	C12—O1—C11—C2	0.2 (2)
O2—S1—C1—C12	28.3 (2)	C12—O1—C11—C10	-179.3 (2)
C14—S1—C1—C12	-85.93 (18)	C3—C2—C11—O1	179.30 (17)
O3—S1—C1—C2	-25.7 (2)	C1—C2—C11—O1	0.2 (2)
O2—S1—C1—C2	-156.10 (19)	C3—C2—C11—C10	-1.1 (3)
C14—S1—C1—C2	89.7 (2)	C1—C2—C11—C10	179.8 (2)
C12—C1—C2—C11	-0.6 (2)	C9—C10—C11—O1	179.38 (19)
S1—C1—C2—C11	-176.64 (17)	C9—C10—C11—C2	-0.1 (3)
C12—C1—C2—C3	-179.4 (2)	C11—O1—C12—C1	-0.7 (2)
S1—C1—C2—C3	4.6 (4)	C11—O1—C12—C13	177.91 (19)
C11—C2—C3—C4	-178.3 (2)	C2—C1—C12—O1	0.8 (2)
C1—C2—C3—C4	0.4 (4)	S1—C1—C12—O1	177.41 (14)
C11—C2—C3—C8	1.3 (3)	C2—C1—C12—C13	-177.3 (3)
C1—C2—C3—C8	180.0 (2)	S1—C1—C12—C13	-0.7 (4)
C2—C3—C4—C5	179.7 (2)	O3—S1—C14—C19	38.53 (17)
C8—C3—C4—C5	0.1 (3)	O2—S1—C14—C19	167.16 (15)
C3—C4—C5—C6	0.3 (3)	C1—S1—C14—C19	-78.17 (16)
C4—C5—C6—C7	-0.1 (3)	O3—S1—C14—C15	-141.32 (15)
C4—C5—C6—Br1	179.40 (17)	O2—S1—C14—C15	-12.69 (18)
C5—C6—C7—C8	-0.6 (3)	C1—S1—C14—C15	101.98 (16)
Br1—C6—C7—C8	179.98 (16)	C19—C14—C15—C16	0.7 (3)
C6—C7—C8—C9	-179.0 (2)	S1—C14—C15—C16	-179.40 (16)
C6—C7—C8—C3	0.9 (3)	C14—C15—C16—C17	0.5 (3)
C4—C3—C8—C7	-0.7 (3)	C15—C16—C17—F1	179.2 (2)
C2—C3—C8—C7	179.70 (18)	C15—C16—C17—C18	-1.5 (4)
C4—C3—C8—C9	179.25 (19)	F1—C17—C18—C19	-179.53 (19)
C2—C3—C8—C9	-0.4 (3)	C16—C17—C18—C19	1.1 (3)
C7—C8—C9—C10	179.0 (2)	C17—C18—C19—C14	0.2 (3)

C3—C8—C9—C10	−0.9 (3)	C15—C14—C19—C18	−1.1 (3)
C8—C9—C10—C11	1.2 (3)	S1—C14—C19—C18	179.04 (15)

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C14—C19 4-fluorophenyl ring.

D—H···A	D—H	H···A	D···A	D—H···A
C5—H5···O2 ⁱ	0.95	2.57	3.312 (3)	135
C18—H18···O2 ⁱⁱ	0.95	2.36	3.266 (3)	160
C19—H19···O3 ⁱ	0.95	2.57	3.113 (2)	117
C10—H10···Cg1 ⁱⁱⁱ	0.95	2.69	3.609 (2)	152

Symmetry codes: (i) $-x+1/2, y+1/2, -z+1/2$; (ii) $x, y+1, z$; (iii) $x+1/2, -y+1/2, z+1/2$.