

Bis(trimethylammonium) tetrachlorido-diphenylstannate(IV)

Tidiane Diop,^a Libasse Diop,^{a*} K. C. Kieran Molloy^b and Gabrielle Kocioc-Köhn^b

^aLaboratoire de Chimie Minérale et Analytique, Département de Chimie, Faculté des Sciences et Techniques, Université Cheikh Anta Diop, Dakar, Senegal, and

^bDepartment of Chemistry, University of Bath, Claverton Down, Bath BA2 7AY, England

Correspondence e-mail: dlibasse@gmail.com


Received 14 December 2010; accepted 6 January 2011

Key indicators: single-crystal X-ray study; $T = 150\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.029; wR factor = 0.072; data-to-parameter ratio = 21.1.

The title compound, $[(\text{CH}_3)_3\text{NH}]_2[\text{Sn}(\text{C}_6\text{H}_5)_2\text{Cl}_4]$, consists of $[(\text{CH}_3)_3\text{NH}]^+$ cations and $[\text{SnPh}_2\text{Cl}_4]^{2-}$ anions in which the Sn atom, located on a centre of inversion, is bonded to four Cl atoms and two phenyl rings, giving an octahedral geometry with the phenyl rings in *trans* positions. In the crystal, the cations and the anions are connected by N—H···Cl hydrogen bonds and C—H···Cl interactions.

Related literature

For background to organotin(IV) chemistry, see: Evans & Karpel (1985); Kapoor *et al.* (2005); Zhang *et al.* (2006). For compounds containing the $[\text{SnPh}_2\text{Cl}_4]^{2-}$ ion in the *cis* or *trans* configuration, see: Ouyang *et al.* (1998); Hazell *et al.* (1998); Fernandez *et al.* (2002); Venkatraman *et al.* (2004); Garcia-Seijo *et al.* (2001); Casas *et al.* (1996); Teoh *et al.* (1992). For related crystal structures, see: Casas *et al.* (1996); Ouyang *et al.* (1998).



Experimental

Crystal data

$(\text{C}_3\text{H}_{10}\text{N})_2[\text{Sn}(\text{C}_6\text{H}_5)_2\text{Cl}_4]$

$M_r = 534.93$

Monoclinic, $P2_1/n$

$a = 9.0072(2)\text{ \AA}$

$b = 8.4125(2)\text{ \AA}$

$c = 14.9473(4)\text{ \AA}$

$\beta = 96.046(1)^\circ$

$V = 1126.30(5)\text{ \AA}^3$

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 1.61\text{ mm}^{-1}$

$T = 150\text{ K}$

$0.25 \times 0.25 \times 0.20\text{ mm}$

Data collection

Nonius KappaCCD diffractometer

Absorption correction: multi-scan
(SORTAV; Blessing, 1995)

$T_{\min} = 0.689$, $T_{\max} = 0.739$

12905 measured reflections

2571 independent reflections

2219 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$

$wR(F^2) = 0.072$

$S = 1.08$

2571 reflections

122 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 2.36\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -1.41\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
N1—H1···Cl1	0.85 (3)	2.42 (3)	3.152 (2)	144 (3)
C5—H5···Cl1 ⁱ	0.95	2.79	3.678 (3)	156

Symmetry code: (i) $x, y + 1, z$.

Data collection: COLLECT (Nonius, 1998); cell refinement: SCALEPACK (Otwinowski & Minor, 1997); data reduction: DENZO/SCALEPACK (Otwinowski & Minor, 1997); program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: SHEXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2238).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Molterini, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Casas, J., Castineiras, A., Couce, M. D., Martinez, G., Sordo, J. & Varela, J. M. (1996). *J. Organomet. Chem.* **517**, 165–172.
- Evans, J. C. & Karpel, S. (1985). *Organotin Compounds in Modern Technology. J. Organomet. Chem. Library*, Vol. 16. Amsterdam: Elsevier.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Fernandez, D., Garcia-Seijo, M. I., Kegl, T., Petocz, G., Kollar, L. & Garcia-Fernandez, M. E. (2002). *Inorg. Chem.* **41**, 4435–4443.
- Garcia-Seijo, M. I., Castineiras, A., Mahieu, B., Janosi, J., Berente, Z., Kollar, L. & Fernandez, G. (2001). *Polyhedron*, **20**, 855–868.
- Hazell, A., Khoo, L. E., Ouyang, J., Rausch, B. J. & Tavares, J. M. (1998). *Acta Cryst. C* **54**, 728–732.
- Kapoor, R. N., Guillory, P., Schulte, L., Cervantes-Lee, F., Haiduc, I., Parkanyi, L. & Pannell, K. H. (2005). *Appl. Organomet. Chem.* **19**, 510–517.
- Nonius (1998). COLLECT. Nonius BV, Delft, The Netherlands.

metal-organic compounds

- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C.W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Ouyang, J., Xu, Y. & Lian, E. K. (1998). *J. Organomet. Chem.* **561**, 143–152.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Teoh, S. G., Teo, S. B., Yeap, G. Y. & Declerq, J. P. (1992). *Polyhedron*, **11**, 2351–2356.
- Venkatraman, R., Ray, P. C. & Fronczek, F. R. (2004). *Acta Cryst. E* **60**, m1035–m1037.
- Zhang, W.-L., Ma, J.-F. & Jiang, H. (2006). *Acta Cryst. E* **62**, m460–m461.

supporting information

Acta Cryst. (2011). E67, m203–m204 [doi:10.1107/S1600536811000870]

Bis(trimethylammonium) tetrachloridodiphenylstannate(IV)

Tidiane Diop, Libasse Diop, K. C. Kieran Molloy and Gabrielle Kocioc-Köhn

S1. Comment

Our interest for organotin(IV) compounds is related to the various applications found for this family of compounds (Evans & Karpel, 1985; Kapoor *et al.*, 2005; Zhang *et al.*, 2006). Many compounds containing the $[\text{SnPh}_2\text{Cl}_4]^{2-}$ -ion in the *cis* or *trans* conformation have been reported (Ouyang *et al.*, 1998; Hazell *et al.*, 1998; Fernandez *et al.*, 2002; Venkatraman *et al.*, 2004; Garcia-Seijo *et al.*, 2001; Teoh *et al.*, 1992). In our search for new organotin(IV) compounds we have initiated here the study of the interactions between $(\text{CH}_3)_3\text{N} \cdot \text{HCl}$ and SnPh_2Cl_2 , which has yielded the title compound.

In the $[\text{Ph}_2\text{SnCl}_4]^{2-}$ -anion, the tin atom is located on a centre of inversion and is bonded to four Cl atoms and two phenyl groups giving an octahedral geometry with the phenyl groups in *trans*- positions (Fig. 1). Consequently, the angle between the two *trans* groups is exactly 180° while the phenyl rings are almost perpendicular to the equitorial SnCl_4 -plane [$\text{C}1-\text{Sn}1-\text{C}11 = 89.39(6)^\circ$, $\text{C}1-\text{Sn}1-\text{C}12 = 90.86(7)^\circ$]. The two $\text{Sn}-\text{C}$ (phenyl) bond distances are $2.149(3)$ Å. The $\text{Sn}-\text{Cl}$ bond distances [$2.5722(6)$ and $2.5796(6)$ Å] are similar to those reported for [Hthiamine] $[\text{SnPh}_2\text{Cl}_4]$. H_2O (Casas *et al.*, 1996), *i.e.* $2.573(2)$ and $2.571(2)$ Å. However, in 8-methoxyquinolinium SnPh_2Cl_4 (Ouyang *et al.*, 1998) these two bond lengths are slightly different [$2.5727(8)$ and $2.6099(8)$ Å].


In the crystal the anion and the cations are linked by $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds (Fig. 1) and $\text{C}-\text{H}\cdots\text{Cl}$ intermolecular interactions (Table 1).

S2. Experimental

The title compound was obtained as a white crystalline solid by reacting trimethylammonium chloride with diphenyltin dichloride in chloroform (2/1 ratio; *M.p.*: 443 K). After slow evaporation of the solvent colourless crystals, suitable for X-ray diffraction analysis, were obtained.

S3. Refinement

The NH H-atom was located in a difference Fourier map and was freely refined. The C-bound H-atoms were included in calculated positions and treated as riding: $\text{C}-\text{H} = 0.95$ and 0.98 Å for CH and CH_3 H-atoms, respectively, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C})$, where $k = 1.2$ for CH H-atoms, and $k = 1.5$ for CH_3 H-atoms.

**Figure 1**

The molecular structure of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level [the C-bound H-atoms have been omitted for clarity; symmetry code: $(') = -x + 1, -y + 1, -z + 1$].

Bis(trimethylammonium) tetrachloridodiphenylstannate(IV)

Crystal data



$M_r = 534.93$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 9.0072 (2)$ Å

$b = 8.4125 (2)$ Å

$c = 14.9473 (4)$ Å

$\beta = 96.046 (1)^\circ$

$V = 1126.30 (5)$ Å³

$Z = 2$

$F(000) = 540$

$D_x = 1.577 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 7261 reflections

$\theta = 2.9\text{--}27.5^\circ$

$\mu = 1.61 \text{ mm}^{-1}$

$T = 150$ K

Block, colourless

$0.25 \times 0.25 \times 0.20$ mm

Data collection

Nonius KappaCCD
diffractometer

Radiation source: fine-focus sealed tube
Graphite monochromator

142 2.0 degree images with ν scans

Absorption correction: multi-scan

(*SORTAV*; Blessing, 1995)

$T_{\min} = 0.689$, $T_{\max} = 0.739$

12905 measured reflections
 2571 independent reflections
 2219 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.3^\circ$
 $h = -11 \rightarrow 11$
 $k = -10 \rightarrow 10$
 $l = -19 \rightarrow 19$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.029$
 $wR(F^2) = 0.072$
 $S = 1.08$
 2571 reflections
 122 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0339P)^2 + 0.7419P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 2.36 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -1.41 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. multi-scan from symmetry-related measurements Sortav (Blessing 1995)

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.50000	0.50000	0.50000	0.0152 (1)
Cl1	0.47692 (7)	0.31690 (7)	0.36079 (4)	0.0227 (2)
Cl2	0.25333 (7)	0.63136 (8)	0.43640 (4)	0.0256 (2)
C1	0.6252 (3)	0.6661 (3)	0.42855 (15)	0.0158 (7)
C2	0.7359 (3)	0.6103 (3)	0.37812 (16)	0.0192 (7)
C3	0.8137 (3)	0.7152 (3)	0.32797 (17)	0.0233 (8)
C4	0.7807 (3)	0.8763 (3)	0.32840 (17)	0.0251 (8)
C5	0.6730 (3)	0.9329 (3)	0.37967 (18)	0.0245 (8)
C6	0.5941 (3)	0.8278 (3)	0.42938 (16)	0.0202 (7)
N1	0.1432 (2)	0.2116 (3)	0.37059 (14)	0.0237 (7)
C7	0.1802 (4)	0.0396 (4)	0.3674 (3)	0.0423 (10)
C8	0.0931 (4)	0.2743 (4)	0.2800 (2)	0.0424 (10)
C9	0.0323 (3)	0.2447 (4)	0.4350 (2)	0.0328 (9)
H2	0.75850	0.50000	0.37790	0.0230*
H3	0.88900	0.67660	0.29360	0.0280*
H4	0.83220	0.94790	0.29340	0.0300*
H5	0.65260	1.04370	0.38110	0.0290*
H6	0.51900	0.86680	0.46380	0.0240*

H1	0.224 (3)	0.258 (3)	0.391 (2)	0.022 (7)*
H7A	0.09310	-0.01930	0.34000	0.0630*
H7B	0.20770	0.00040	0.42870	0.0630*
H7C	0.26400	0.02430	0.33160	0.0630*
H8A	0.16890	0.25150	0.23930	0.0640*
H8B	0.07840	0.38950	0.28350	0.0640*
H8C	-0.00120	0.22340	0.25710	0.0640*
H9A	0.01660	0.35970	0.43870	0.0490*
H9B	0.06950	0.20380	0.49450	0.0490*
H9C	-0.06240	0.19260	0.41420	0.0490*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.0147 (1)	0.0171 (1)	0.0144 (1)	-0.0006 (1)	0.0041 (1)	0.0010 (1)
Cl1	0.0237 (3)	0.0258 (3)	0.0196 (3)	-0.0035 (2)	0.0072 (2)	-0.0058 (2)
Cl2	0.0181 (3)	0.0304 (3)	0.0286 (3)	0.0049 (2)	0.0043 (2)	0.0089 (3)
C1	0.0158 (11)	0.0185 (12)	0.0132 (11)	-0.0032 (9)	0.0025 (8)	0.0009 (8)
C2	0.0196 (12)	0.0212 (12)	0.0169 (12)	-0.0013 (9)	0.0030 (9)	0.0000 (9)
C3	0.0199 (12)	0.0335 (15)	0.0172 (12)	-0.0030 (10)	0.0056 (9)	0.0014 (10)
C4	0.0240 (13)	0.0290 (14)	0.0215 (13)	-0.0092 (11)	-0.0009 (10)	0.0070 (10)
C5	0.0287 (14)	0.0166 (12)	0.0275 (13)	-0.0034 (10)	-0.0007 (11)	0.0033 (10)
C6	0.0211 (12)	0.0191 (12)	0.0204 (12)	0.0005 (9)	0.0028 (9)	-0.0018 (9)
N1	0.0182 (11)	0.0313 (12)	0.0210 (11)	-0.0051 (9)	-0.0002 (9)	-0.0037 (9)
C7	0.0308 (16)	0.0350 (16)	0.060 (2)	-0.0007 (13)	-0.0005 (15)	-0.0132 (15)
C8	0.0371 (17)	0.068 (2)	0.0211 (15)	-0.0077 (15)	-0.0020 (13)	0.0081 (14)
C9	0.0277 (15)	0.0436 (17)	0.0282 (15)	-0.0044 (12)	0.0078 (12)	-0.0067 (12)

Geometric parameters (\AA , ^\circ)

Sn1—Cl1	2.5796 (6)	C5—C6	1.397 (4)
Sn1—Cl2	2.5722 (6)	C2—H2	0.9500
Sn1—C1	2.149 (3)	C3—H3	0.9500
Sn1—Cl1 ⁱ	2.5796 (6)	C4—H4	0.9500
Sn1—Cl2 ⁱ	2.5722 (6)	C5—H5	0.9500
Sn1—C1 ⁱ	2.149 (3)	C6—H6	0.9500
N1—C9	1.485 (3)	C7—H7A	0.9800
N1—C7	1.487 (4)	C7—H7B	0.9800
N1—C8	1.479 (4)	C7—H7C	0.9800
N1—H1	0.85 (3)	C8—H8A	0.9800
C1—C6	1.389 (4)	C8—H8B	0.9800
C1—C2	1.393 (4)	C8—H8C	0.9800
C2—C3	1.394 (4)	C9—H9A	0.9800
C3—C4	1.388 (4)	C9—H9B	0.9800
C4—C5	1.383 (4)	C9—H9C	0.9800
Cl1—Sn1—Cl2	88.01 (2)	C1—C2—H2	120.00
Cl1—Sn1—C1	89.39 (6)	C3—C2—H2	120.00

Cl1—Sn1—Cl1 ⁱ	180.00	C2—C3—H3	120.00
Cl1—Sn1—Cl2 ⁱ	91.99 (2)	C4—C3—H3	120.00
Cl1—Sn1—C1 ⁱ	90.61 (6)	C5—C4—H4	120.00
Cl2—Sn1—C1	90.86 (7)	C3—C4—H4	120.00
Cl1 ⁱ —Sn1—Cl2	91.99 (2)	C4—C5—H5	120.00
Cl2—Sn1—Cl2 ⁱ	180.00	C6—C5—H5	120.00
Cl2—Sn1—C1 ⁱ	89.14 (7)	C5—C6—H6	120.00
Cl1 ⁱ —Sn1—C1	90.61 (6)	C1—C6—H6	120.00
Cl2 ⁱ —Sn1—C1	89.14 (7)	N1—C7—H7A	109.00
C1—Sn1—C1 ⁱ	180.00	N1—C7—H7B	109.00
Cl1 ⁱ —Sn1—Cl2 ⁱ	88.01 (2)	N1—C7—H7C	110.00
Cl1 ⁱ —Sn1—C1 ⁱ	89.39 (6)	H7A—C7—H7B	109.00
Cl2 ⁱ —Sn1—C1 ⁱ	90.86 (7)	H7A—C7—H7C	110.00
C7—N1—C8	111.4 (3)	H7B—C7—H7C	109.00
C7—N1—C9	111.8 (2)	N1—C8—H8A	109.00
C8—N1—C9	111.4 (2)	N1—C8—H8B	109.00
C8—N1—H1	109.3 (19)	N1—C8—H8C	109.00
C9—N1—H1	106.8 (19)	H8A—C8—H8B	109.00
C7—N1—H1	105.7 (17)	H8A—C8—H8C	109.00
C2—C1—C6	119.4 (2)	H8B—C8—H8C	109.00
Sn1—C1—C2	119.52 (18)	N1—C9—H9A	109.00
Sn1—C1—C6	121.02 (19)	N1—C9—H9B	109.00
C1—C2—C3	120.4 (2)	N1—C9—H9C	109.00
C2—C3—C4	119.7 (2)	H9A—C9—H9B	109.00
C3—C4—C5	120.1 (2)	H9A—C9—H9C	109.00
C4—C5—C6	120.2 (2)	H9B—C9—H9C	110.00
C1—C6—C5	120.1 (2)		
Cl1—Sn1—C1—C2	−43.53 (19)	Sn1—C1—C2—C3	177.05 (19)
Cl1—Sn1—C1—C6	134.3 (2)	C6—C1—C2—C3	−0.8 (4)
Cl2—Sn1—C1—C2	−131.53 (19)	Sn1—C1—C6—C5	−177.54 (19)
Cl2—Sn1—C1—C6	46.3 (2)	C2—C1—C6—C5	0.3 (4)
Cl1 ⁱ —Sn1—C1—C2	136.47 (19)	C1—C2—C3—C4	0.1 (4)
Cl1 ⁱ —Sn1—C1—C6	−45.7 (2)	C2—C3—C4—C5	1.2 (4)
Cl2 ⁱ —Sn1—C1—C2	48.47 (19)	C3—C4—C5—C6	−1.7 (4)
Cl2 ⁱ —Sn1—C1—C6	−133.7 (2)	C4—C5—C6—C1	1.0 (4)

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1 \cdots Cl1	0.85 (3)	2.42 (3)	3.152 (2)	144 (3)
C5—H5 \cdots Cl1 ⁱⁱ	0.95	2.79	3.678 (3)	156

Symmetry code: (ii) $x, y+1, z$.