

(Z)-5-Benzylidene-3-butyl-4-phenyl-1,3-oxazolidin-2-one

Jin-Wu Zhao* and Jing-Xiu Xu

School of Pharmacy, Guang Dong Medical College, Dong Guan 523808, People's Republic of China
Correspondence e-mail: jinwuzhao@gmail.com


Received 1 March 2011; accepted 7 March 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.042; wR factor = 0.117; data-to-parameter ratio = 14.3.

In the title compound, $C_{20}H_{21}NO_2$, the benzyl group and the oxazolidin-2-one unit are each essentially planar, with maximum deviations of 0.026 (2) and 0.031 (2) \AA , respectively. The dihedral angle between the phenyl ring and the oxazolidin-2-one unit is $69.25(2)^\circ$. In the crystal, molecules are linked by weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For general background to 2-oxazolidinone derivatives and for heterocyclic systems of antibacterial interest, see: Mukhtar & Wright (2005); Ager *et al.* (1996); Renslo *et al.* (2006). For bond-length data, see: Allen *et al.* (1987). For the chemical structure of the title compound established from NMR data, see: Yoo & Li (2008).



Experimental

Crystal data

$C_{20}H_{21}NO_2$
 $M_r = 307.38$
Monoclinic, $P2_1/n$

$a = 10.029(2)\text{ \AA}$
 $b = 9.1941(18)\text{ \AA}$
 $c = 18.389(4)\text{ \AA}$

$\beta = 100.51(3)^\circ$
 $V = 1667.1(6)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.08\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.31 \times 0.25 \times 0.18\text{ mm}$

Data collection

Rigaku/MSC Mercury CCD diffractometer
Absorption correction: multi-scan (*REQAB*; Jacobson, 1998)
 $T_{\min} = 0.989$, $T_{\max} = 0.997$

12969 measured reflections
2994 independent reflections
1803 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.117$
 $S = 1.03$
2994 reflections

209 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.17\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.18\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$ and $Cg2$ are the centroids of the C1–C6 and C11–C16 rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$C4-\text{H4}\cdots O2^i$	0.93	2.58	3.343 (3)	141
$C3-\text{H3}\cdots Cg2^{ii}$	0.93	2.93	3.674 (2)	138
$C12-\text{H12}\cdots Cg1^{iii}$	0.93	2.91	3.706 (3)	145
$C20-\text{H20B}\cdots Cg2^{iv}$	0.96	2.92	3.812 (1)	156

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (iii) $-x + 1, -y, -z$; (iv) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPII* (Johnson, 1976); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge Guang Dong Medical College for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BG2393).

References

- Ager, D. J., Prakash, I. & Schaad, D. R. (1996). *Chem. Rev.* **96**, 835–876.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Jacobson, R. (1998). *REQAB*. Molecular Structure Corporation, The Woodlands, Texas, USA.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Mukhtar, T. A. & Wright, G. D. (2005). *Chem. Rev.* **105**, 529–452.
- Renslo, A. R., Luehr, G. W. & Gordeev, M. F. (2006). *Bioorg. Med. Chem.* **14**, 4227–4240.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2002). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Yoo, W. J. & Li, C. J. (2008). *Adv. Synth. Catal.* **350**, 1503–1506.

supporting information

Acta Cryst. (2011). E67, o882 [doi:10.1107/S1600536811008762]

(Z)-5-Benzylidene-3-butyl-4-phenyl-1,3-oxazolidin-2-one

Jin-Wu Zhao and Jing-Xiu Xu

S1. Comment

The title compound, (I), $C_{19}H_{17}N_3O_2$, is a 2-oxazolidinone derivative. The 2-oxazolidinone ring is an important heterocyclic structural unit. It possesses significant antibacterial activities and plays an important role as an intermediate for the synthesis of more complex active organic compounds and further functionalized heterocyclic systems of antibacterial interest (Mukhtar *et al.*, 2005; Ager *et al.*, 1996; Renslo *et al.*, 2006). The structure of title compound has been established from the NMR data (Yoo *et al.*, 2008). However, the crystal structure of title compound has not been reported. In view of this, the crystal structure determination of the title compound was carried out and the results are presented here.

As depicted in Fig. 1, the benzyl group A (C1–C7) and the heterocyclic ring of 2-oxazolidinone B (N1/C8–C10/O1/O2) are almost planar with maximum deviations of 0.024 (2) Å for C7 and 0.031 (2) Å for O2, respectively, and determine a dihedral angle of 2.56 (2) °. The phenyl ring C (C11–C16) is of also planar (max. deviation 0.007 (2) Å for C15).

The butyl moiety (C17–C20) adopts a slight twist conformation with C20 displaced by 0.123 (3) Å from the plane defined by the atoms C17–C19 (D). The dihedral angles between C/A, C/B, C/D, D/A and D/B are 69.05 (2) °, 69.30 (3) °, 75.17 (1) °, 85.79 (4) ° and 83.29 (2) °, respectively. The bond lengths and bond angles are within normal range (Allen *et al.*, 1987). The molecules are linked into a three-dimensional supramolecular network through intermolecular C—H···O hydrogen bonding interactions and C—H···π stacking interactions (Table 1, Fig. 2). The H-to-centroid distances of $H3\cdots Cg2^i = 2.93$ (2), $H12\cdots Cg1^{ii} = 2.91$ (3) and $H20B\cdots Cg2^{iii} = 2.91$ (4) [$Cg1$ and $Cg2$ are the centroids of the C1, C2, C3, C4, C5, C6 ring, and C11, C12, C13, C14, C15, C16 ring, respectively. Symmetry codes: (i) $-1/2 + x, 1/2 - y, 1/2 + z$; (ii) $3/2 - x, 1/2 + y, 1/2 - z$; (iii) $-1/2 + x, 1/2 - y, -1/2 + z$]. In addition, intramolecular C—H···O and C—H···N hydrogen bonds are also observed.

S2. Experimental

A 15 ml polytetrafluoroethylene (PTFE) reaction vessel was charged with copper(I)iodide (0.6 mmol, 0.114 g), butylamine (4 mmol, 0.293 g), benzaldehyde (4 mmol, 0.425 g) and ethynylbenzene (3 mmol, 0.204 g). Then the vessel was fixed into a stainless steel autoclave with a pressure-regulating system. The autoclave was sealed. Liquid CO₂ was introduced from a cylinder and the reaction mixture was magnetically stirred at 373 K under 8 MPa for 12 h. The vessel was cooled with an ice bath and the pressure was released slowly to atmospheric pressure after the reaction completed. The reaction mixture was flushed with EtOAc (30 ml) and the ethyl acetate fractions were combined. The resulting solvent was placed through a plug of silica gel, and then evaporated. The residue was purified by silica gel (200–300 mesh) column by elution with MSO:EtOAc (8:1) to give 20 fractions (200 ml per fraction). The title compound (539.6 mg) was isolated from the fractions 3–17 (yield 71.2%). Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of the title compound in petroleum ether at room temperature.

S3. Refinement

All H atoms were located on the difference maps, and were treated as riding atoms with C—H distances of 0.93, 0.96, 0.97 and 0.98 Å, for aryl, methyl, methine and tertiary alkyl, respectively, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}$ (methyl C-atoms) and $1.2U_{\text{eq}}$ (non-methyl C-atoms). The highest peak is located 1.54 Å from C1 and the deepest hole is located 0.86 Å from H13.





Figure 1

The molecular structure of the tile compound, showing 50% probability displacement ellipsoids and the atom-numbering scheme.

**Figure 2**

Packing diagram of the title compound; C—H···O, C—H···N and C—H··· π interactions are shown as dashed lines. The H-atoms not involved in H-bonds have been excluded for clarity.

(Z)-5-Benzylidene-3-butyl-4-phenyl-1,3-oxazolidin-2-one

Crystal data

$C_{20}H_{21}NO_2$
 $M_r = 307.38$
Monoclinic, $P2_1/n$
Hall symbol: -P 2yn
 $a = 10.029 (2) \text{ \AA}$
 $b = 9.1941 (18) \text{ \AA}$
 $c = 18.389 (4) \text{ \AA}$
 $\beta = 100.51 (3)^\circ$
 $V = 1667.1 (6) \text{ \AA}^3$
 $Z = 4$

Data collection

Rigaku/MSC Mercury CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
Absorption correction: multi-scan
(REQAB; Jacobson, 1998)
 $T_{\min} = 0.989$, $T_{\max} = 0.997$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.117$
 $S = 1.03$
2994 reflections
209 parameters

$F(000) = 656$
 $D_x = 1.225 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Cell parameters from 5837 reflections
 $\theta = 2.8\text{--}27.9^\circ$
 $\mu = 0.08 \text{ mm}^{-1}$
 $T = 293 \text{ K}$
Block, colorless
 $0.31 \times 0.25 \times 0.18 \text{ mm}$

12969 measured reflections
2994 independent reflections
1803 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$
 $\theta_{\max} = 25.2^\circ$, $\theta_{\min} = 3.0^\circ$
 $h = -12 \rightarrow 12$
 $k = -11 \rightarrow 11$
 $l = -22 \rightarrow 21$

0 restraints
Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0472P)^2 + 0.3223P]$
 where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.17 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.59314 (19)	0.3280 (2)	0.36032 (11)	0.0590 (5)
H1	0.5441	0.3603	0.3154	0.071*
C2	0.5426 (2)	0.3511 (3)	0.42412 (12)	0.0675 (6)
H2	0.4609	0.4003	0.4218	0.081*
C3	0.6115 (2)	0.3023 (3)	0.49109 (12)	0.0680 (6)
H3	0.5764	0.3174	0.5339	0.082*
C4	0.7330 (2)	0.2308 (2)	0.49425 (11)	0.0636 (6)
H4	0.7798	0.1966	0.5393	0.076*
C5	0.78512 (19)	0.2099 (2)	0.43106 (10)	0.0532 (5)
H5	0.8682	0.1630	0.4342	0.064*
C6	0.71618 (17)	0.2574 (2)	0.36199 (10)	0.0477 (5)
C7	0.77658 (18)	0.2274 (2)	0.29672 (10)	0.0517 (5)
H7	0.8565	0.1737	0.3059	0.062*
C8	0.73519 (17)	0.2650 (2)	0.22699 (10)	0.0477 (5)
C9	0.6003 (2)	0.3656 (2)	0.12780 (11)	0.0560 (5)
C10	0.79952 (18)	0.2321 (2)	0.16033 (10)	0.0509 (5)
H10	0.7967	0.1270	0.1512	0.061*
C11	0.94436 (17)	0.2858 (2)	0.16712 (9)	0.0471 (5)
C12	0.9744 (2)	0.4219 (2)	0.14300 (11)	0.0569 (5)
H12	0.9047	0.4837	0.1217	0.068*
C13	1.1078 (2)	0.4670 (3)	0.15040 (12)	0.0669 (6)
H13	1.1271	0.5584	0.1333	0.080*
C14	1.2119 (2)	0.3782 (3)	0.18269 (12)	0.0688 (6)
H14	1.3013	0.4090	0.1874	0.083*
C15	1.1831 (2)	0.2442 (3)	0.20790 (12)	0.0678 (6)
H15	1.2532	0.1842	0.2306	0.081*
C16	1.05059 (19)	0.1974 (2)	0.19989 (11)	0.0585 (5)
H16	1.0322	0.1054	0.2167	0.070*
C17	0.7052 (2)	0.2948 (3)	0.02324 (10)	0.0651 (6)
H17A	0.7983	0.3011	0.0159	0.078*
H17B	0.6563	0.3771	-0.0016	0.078*
C18	0.6430 (2)	0.1560 (2)	-0.01191 (10)	0.0643 (6)

H18A	0.5527	0.1447	-0.0008	0.077*
H18B	0.6971	0.0738	0.0094	0.077*
C19	0.6343 (2)	0.1550 (3)	-0.09517 (10)	0.0707 (6)
H19A	0.7254	0.1578	-0.1060	0.085*
H19B	0.5875	0.2421	-0.1158	0.085*
C20	0.5609 (3)	0.0224 (3)	-0.13268 (12)	0.0880 (8)
H20A	0.6069	-0.0643	-0.1128	0.132*
H20B	0.5600	0.0270	-0.1849	0.132*
H20C	0.4694	0.0210	-0.1240	0.132*
N1	0.70329 (15)	0.30487 (19)	0.10237 (8)	0.0570 (5)
O1	0.61860 (12)	0.34597 (15)	0.20363 (7)	0.0572 (4)
O2	0.50276 (13)	0.43149 (18)	0.09551 (8)	0.0707 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0542 (12)	0.0656 (14)	0.0581 (12)	0.0036 (10)	0.0127 (10)	0.0080 (10)
C2	0.0576 (12)	0.0786 (16)	0.0709 (14)	0.0087 (12)	0.0242 (11)	0.0050 (12)
C3	0.0719 (14)	0.0809 (17)	0.0569 (13)	-0.0049 (13)	0.0272 (11)	-0.0012 (11)
C4	0.0655 (13)	0.0775 (16)	0.0483 (12)	-0.0063 (12)	0.0121 (10)	0.0030 (10)
C5	0.0486 (10)	0.0606 (13)	0.0501 (11)	-0.0024 (9)	0.0079 (9)	0.0028 (9)
C6	0.0439 (10)	0.0522 (12)	0.0478 (11)	-0.0035 (9)	0.0103 (8)	0.0021 (9)
C7	0.0475 (10)	0.0575 (13)	0.0496 (11)	0.0031 (9)	0.0079 (9)	0.0033 (9)
C8	0.0403 (9)	0.0536 (12)	0.0484 (11)	0.0000 (9)	0.0061 (8)	0.0021 (9)
C9	0.0478 (11)	0.0695 (14)	0.0498 (11)	-0.0064 (11)	0.0066 (9)	0.0065 (10)
C10	0.0509 (10)	0.0564 (12)	0.0440 (10)	0.0004 (9)	0.0050 (8)	-0.0012 (9)
C11	0.0451 (10)	0.0569 (13)	0.0397 (10)	0.0054 (9)	0.0090 (8)	-0.0048 (9)
C12	0.0545 (12)	0.0586 (14)	0.0572 (12)	0.0060 (10)	0.0087 (9)	0.0024 (10)
C13	0.0655 (13)	0.0702 (15)	0.0667 (14)	-0.0091 (12)	0.0163 (11)	-0.0021 (11)
C14	0.0495 (12)	0.0943 (19)	0.0639 (13)	-0.0051 (13)	0.0134 (10)	-0.0137 (13)
C15	0.0531 (12)	0.0852 (18)	0.0638 (14)	0.0157 (12)	0.0074 (10)	-0.0040 (12)
C16	0.0573 (12)	0.0622 (14)	0.0551 (12)	0.0092 (11)	0.0074 (10)	0.0002 (10)
C17	0.0596 (12)	0.0931 (17)	0.0412 (11)	-0.0056 (12)	0.0060 (9)	0.0018 (11)
C18	0.0695 (13)	0.0788 (16)	0.0429 (11)	0.0114 (12)	0.0061 (9)	0.0005 (10)
C19	0.0640 (13)	0.1041 (19)	0.0441 (11)	0.0091 (13)	0.0105 (10)	-0.0017 (12)
C20	0.122 (2)	0.091 (2)	0.0481 (13)	0.0186 (16)	0.0078 (13)	-0.0100 (12)
N1	0.0505 (9)	0.0798 (13)	0.0386 (9)	0.0037 (9)	0.0026 (7)	0.0022 (8)
O1	0.0494 (8)	0.0730 (10)	0.0493 (8)	0.0095 (7)	0.0088 (6)	0.0084 (7)
O2	0.0487 (8)	0.0929 (12)	0.0676 (9)	0.0062 (8)	0.0029 (7)	0.0216 (8)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.376 (3)	C11—C16	1.387 (3)
C1—C6	1.390 (3)	C12—C13	1.383 (3)
C1—H1	0.9300	C12—H12	0.9300
C2—C3	1.373 (3)	C13—C14	1.372 (3)
C2—H2	0.9300	C13—H13	0.9300
C3—C4	1.376 (3)	C14—C15	1.365 (3)

C3—H3	0.9300	C14—H14	0.9300
C4—C5	1.372 (3)	C15—C16	1.379 (3)
C4—H4	0.9300	C15—H15	0.9300
C5—C6	1.401 (2)	C16—H16	0.9300
C5—H5	0.9300	C17—N1	1.461 (2)
C6—C7	1.466 (3)	C17—C18	1.513 (3)
C7—C8	1.320 (2)	C17—H17A	0.9700
C7—H7	0.9300	C17—H17B	0.9700
C8—O1	1.387 (2)	C18—C19	1.518 (3)
C8—C10	1.516 (3)	C18—H18A	0.9700
C9—O2	1.210 (2)	C18—H18B	0.9700
C9—N1	1.332 (2)	C19—C20	1.523 (3)
C9—O1	1.385 (2)	C19—H19A	0.9700
C10—N1	1.463 (2)	C19—H19B	0.9700
C10—C11	1.518 (2)	C20—H20A	0.9600
C10—H10	0.9800	C20—H20B	0.9600
C11—C12	1.380 (3)	C20—H20C	0.9600
C2—C1—C6	121.09 (19)	C14—C13—H13	119.7
C2—C1—H1	119.5	C12—C13—H13	119.7
C6—C1—H1	119.5	C15—C14—C13	119.5 (2)
C1—C2—C3	120.7 (2)	C15—C14—H14	120.2
C1—C2—H2	119.6	C13—C14—H14	120.2
C3—C2—H2	119.6	C14—C15—C16	120.3 (2)
C4—C3—C2	119.4 (2)	C14—C15—H15	119.9
C4—C3—H3	120.3	C16—C15—H15	119.9
C2—C3—H3	120.3	C15—C16—C11	120.8 (2)
C5—C4—C3	120.2 (2)	C15—C16—H16	119.6
C5—C4—H4	119.9	C11—C16—H16	119.6
C3—C4—H4	119.9	N1—C17—C18	113.57 (17)
C4—C5—C6	121.53 (19)	N1—C17—H17A	108.9
C4—C5—H5	119.2	C18—C17—H17A	108.9
C6—C5—H5	119.2	N1—C17—H17B	108.9
C1—C6—C5	117.08 (18)	C18—C17—H17B	108.9
C1—C6—C7	124.57 (17)	H17A—C17—H17B	107.7
C5—C6—C7	118.34 (17)	C17—C18—C19	112.31 (18)
C8—C7—C6	130.01 (18)	C17—C18—H18A	109.1
C8—C7—H7	115.0	C19—C18—H18A	109.1
C6—C7—H7	115.0	C17—C18—H18B	109.1
C7—C8—O1	122.65 (17)	C19—C18—H18B	109.1
C7—C8—C10	128.95 (17)	H18A—C18—H18B	107.9
O1—C8—C10	108.40 (15)	C18—C19—C20	113.3 (2)
O2—C9—N1	130.30 (19)	C18—C19—H19A	108.9
O2—C9—O1	120.42 (19)	C20—C19—H19A	108.9
N1—C9—O1	109.27 (17)	C18—C19—H19B	108.9
N1—C10—C8	100.16 (15)	C20—C19—H19B	108.9
N1—C10—C11	113.93 (16)	H19A—C19—H19B	107.7
C8—C10—C11	114.25 (15)	C19—C20—H20A	109.5

N1—C10—H10	109.4	C19—C20—H20B	109.5
C8—C10—H10	109.4	H20A—C20—H20B	109.5
C11—C10—H10	109.4	C19—C20—H20C	109.5
C12—C11—C16	118.39 (18)	H20A—C20—H20C	109.5
C12—C11—C10	122.06 (17)	H20B—C20—H20C	109.5
C16—C11—C10	119.54 (19)	C9—N1—C10	112.76 (15)
C13—C12—C11	120.3 (2)	C9—N1—C17	121.86 (16)
C13—C12—H12	119.8	C10—N1—C17	124.73 (16)
C11—C12—H12	119.8	C9—O1—C8	109.36 (15)
C14—C13—C12	120.6 (2)		

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 are the centroids of the C1—C6 and C11—C16 rings, respectively.

D—H···A	D—H	H···A	D···A	D—H···A
C4—H4···O2 ⁱ	0.93	2.58	3.343 (3)	141
C3—H3···Cg2 ⁱⁱ	0.93	2.93	3.674 (2)	138
C12—H12···Cg1 ⁱⁱⁱ	0.93	2.91	3.706 (3)	145
C20—H20B···Cg2 ^{iv}	0.96	2.92	3.812 (1)	156

Symmetry codes: (i) $x+1/2, -y+1/2, z+1/2$; (ii) $x-1/2, -y+1/2, z+1/2$; (iii) $-x+1, -y, -z$; (iv) $x-1/2, -y+1/2, z-1/2$.