

2,4-Bis(4-propoxypyphenyl)-3-azabicyclo-[3.3.1]nonan-9-one

P. Parthiban,^a V. Ramkumar^b and Yeon Tae Jeong^{a*}

^aDepartment of Image Science and Engineering, Pukyong National University, Busan 608 739, Republic of Korea, and ^bDepartment of Chemistry, IIT Madras, Chennai, TamilNadu, India

Correspondence e-mail: ytjeong@pknu.ac.kr


Received 19 February 2011; accepted 28 February 2011

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.044; wR factor = 0.104; data-to-parameter ratio = 12.7.

In the title compound, $\text{C}_{26}\text{H}_{33}\text{NO}_3$, a crystallographic mirror plane bisects the molecule (two C atoms, one O atom and one N atom lie on the mirror plane). The molecule exists in a twin-chair conformation with equatorial orientations of the 4-propoxypyphenyl groups. The dihedral angle between the 4-propoxypyphenyl groups is $31.58(3)^\circ$.

Related literature

For background to 3-azabicyclononanes, see: Jeyaraman & Avila (1981); Barker *et al.* (2005); Parthiban *et al.* (2009a, 2010b,c). For related structures, see: Parthiban *et al.* (2009b,c, 2010a); Smith-Verdier *et al.* (1983); Padegimas & Kovacic (1972). For ring puckering and asymmetry parameters, see: Cremer & Pople (1975); Nardelli (1983).



Experimental

Crystal data

$\text{C}_{26}\text{H}_{33}\text{NO}_3$
 $M_r = 407.53$
Orthorhombic, $Pnma$
 $a = 7.3846(4)\text{ \AA}$

$b = 29.3963(19)\text{ \AA}$
 $c = 10.2739(7)\text{ \AA}$
 $V = 2230.3(2)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.08\text{ mm}^{-1}$

$T = 298\text{ K}$
 $0.25 \times 0.22 \times 0.20\text{ mm}$

Data collection

Bruker APEXII CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2004)
 $T_{\min} = 0.981$, $T_{\max} = 0.985$

7260 measured reflections
1860 independent reflections
1121 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.104$
 $S = 1.00$
1860 reflections
146 parameters

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\max} = 0.14\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.21\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT-Plus* (Bruker, 2004); data reduction: *SAINT-Plus* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge the Department of Chemistry, IIT Madras, for the X-ray data collection. This research was supported by the Industrial Technology Development program, which was conducted by the Ministry of Knowledge Economy of the Korean Government.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5803).

References

- Barker, D., Lin, D. H. S., Carland, J. E., Chu, C. P. Y., Chebib, M., Brimble, M. A., Savage, G. P. & McLeod, M. D. (2005). *Bioorg. Med. Chem.* **13**, 4565–4575.
- Bruker (2004). *APEX2*, *XPREP*, *SAINT-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Jeyaraman, R. & Avila, S. (1981). *Chem. Rev.* **81**, 149–174.
- Nardelli, M. (1983). *Acta Cryst. C39*, 1141–1142.
- Padegimas, S. J. & Kovacic, P. (1972). *J. Org. Chem.* **37**, 2672–2676.
- Parthiban, P., Aridoss, G., Rathika, P., Ramkumar, V. & Kabilan, S. (2009a). *Bioorg. Med. Chem. Lett.* **19**, 6981–6985.
- Parthiban, P., Ramkumar, V., Amirthaganesan, S. & Jeong, Y. T. (2009b). *Acta Cryst. E65*, o1356.
- Parthiban, P., Ramkumar, V., Jeong, Y. T. (2010a). *Acta Cryst. E66*, o48–o49.
- Parthiban, P., Ramkumar, V., Kim, M. S., Son, S. M. & Jeong, Y. T. (2009c). *Acta Cryst. E65*, o1383.
- Parthiban, P., Rathika, P., Park, K. S. & Jeong, Y. T. (2010b). *Monatsh. Chem.* **141**, 79–93.
- Parthiban, P., Rathika, P., Ramkumar, V., Son, S. M. & Jeong, Y. T. (2010c). *Bioorg. Med. Chem. Lett.* **20**, 1642–1647.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Smith-Verdier, P., Florencio, F. & García-Blanco, S. (1983). *Acta Cryst. C39*, 101–103.

supporting information

Acta Cryst. (2011). E67, o790 [doi:10.1107/S1600536811007483]

2,4-Bis(4-propoxyphenyl)-3-azabicyclo[3.3.1]nonan-9-one

P. Parthiban, V. Ramkumar and Yeon Tae Jeong

S1. Comment

3-Azabicyclononanes are important class of heterocycles due to their broad spectrum of biological activities such as antibacterial, antimycobacterial, antifungal, anticancer, antitussive, anti-inflammatory, sedative, antipyretic and calcium antagonistic activities (Jeyaraman & Avila, 1981; Barker *et al.*, 2005; Parthiban *et al.*, 2009*a*, 2010*b,c*). Since the stereochemistry plays an important role in biological actions, it is important to establish the stereochemistry of the synthesized biologically potent molecules. Owing to the diverse possibilities in conformation of the 3-azabicycles, *viz.*, chair-chair (Parthiban *et al.*, 2009*a*), chair-boat (Smith-Verdier *et al.*, 1983) and boat-boat (Padegimas & Kovacic, 1972). This crystal study has been carried out to expose the conformation of the title bicyclic compound.


The analysis of torsion angles, asymmetry parameters and puckering parameters calculated for the title compound shows that the piperidine ring adopts near ideal chair conformation with a total puckering amplitude, Q_T of 0.615 (2) Å and the phase angle θ is 0.00 (1)°. (Cremer & Pople, 1975). The smallest displacement asymmetry parameters q_2 and q_3 are 0.00 and 0.615 (2)°, respectively (Nardelli, 1983). However, the cyclohexane ring deviates from the ideal chair conformation according to Cremer and Pople by Q_T = 0.562 (2) and θ = 16.8 (2)° (Cremer & Pople, 1975) as well as Nardelli by q_2 = 0.162 (2) and q_3 = 0.538 (2)° (Nardelli, 1983). Hence, the title compound $C_{28}H_{33}N O_3$, exists in a twin-chair conformation with equatorial orientation of 4-propoxyphenyl groups on the heterocycle and are orientated at an angle of 31.58 (3)° to each other. The torsion angle of C3—C2—C1—C6 are 64.09 (3)°. The crystal crystal packing is stabilized by weak van der Waals interaction.

S2. Experimental

To the warm solution of 0.075 mol (5.78 g) ammonium acetate in 50 ml of absolute ethanol, 0.1 mol (16.42 g/15.80 ml) of *para*-n-propoxybenzaldehyde and 0.05 mol (4.90 g/5.18 ml) of cyclohexanone were added. The mixture was gently warmed on a hot plate at 303–308 K (30–35°C) with moderate stirring till the complete consumption of the starting materials, monitored by TLC. At the end, the crude azabicyclic ketone was separated by filtration and washed with 1:5 cold ethanol-ether mixture. Colourless blocks of the title compound were obtained by recrystallization from ethanol.

S3. Refinement

Nitrogen H atoms were located in a difference Fourier map and refined isotropically. Other hydrogen atoms were fixed geometrically and allowed to ride on the parent carbon atoms, with aromatic C—H = 0.93 Å, aliphatic C—H = 0.98 Å and methylene C—H = 0.97 Å. The displacement parameters were set for phenyl, methylene and aliphatic H atoms at $U_{iso}(H)$ = 1.2 $U_{eq}(C)$.

**Figure 1**

Anisotropic displacement representation of the molecule with atoms represented with 30% probability ellipsoids.
Symmetry code: (i) $x, 1/2-y, z$.

2,4-Bis(4-propoxyphenyl)-3-azabicyclo[3.3.1]nonan-9-one

Crystal data

$C_{26}H_{33}NO_3$
 $M_r = 407.53$
Orthorhombic, $Pnma$
Hall symbol: -P 2ac 2n
 $a = 7.3846 (4)$ Å
 $b = 29.3963 (19)$ Å
 $c = 10.2739 (7)$ Å
 $V = 2230.3 (2)$ Å³
 $Z = 4$

$F(000) = 880$
 $D_x = 1.214 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 1042 reflections
 $\theta = 2.3-20.8^\circ$
 $\mu = 0.08 \text{ mm}^{-1}$
 $T = 298$ K
Block, colourless
 $0.25 \times 0.22 \times 0.20$ mm

Data collection

Bruker APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2004)
 $T_{\min} = 0.981$, $T_{\max} = 0.985$

7260 measured reflections
1860 independent reflections
1121 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$
 $\theta_{\max} = 26.2^\circ$, $\theta_{\min} = 2.1^\circ$
 $h = -8 \rightarrow 8$
 $k = -31 \rightarrow 32$
 $l = -12 \rightarrow 10$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.104$
 $S = 1.00$
1860 reflections
146 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0404P)^2 + 0.4368P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.14 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6512 (3)	0.29134 (6)	0.6070 (2)	0.0384 (5)
H1	0.6934	0.2893	0.6972	0.046*
C2	0.8207 (3)	0.29225 (7)	0.5180 (2)	0.0426 (6)
H2	0.8944	0.3188	0.5408	0.051*
C3	0.7815 (3)	0.29351 (7)	0.3708 (2)	0.0474 (6)
H3A	0.7004	0.3187	0.3529	0.057*
H3B	0.8940	0.2993	0.3249	0.057*
C4	0.6973 (4)	0.2500	0.3174 (3)	0.0482 (8)
H4A	0.7093	0.2500	0.2234	0.058*
H4B	0.5691	0.2500	0.3377	0.058*
C5	0.9286 (4)	0.2500	0.5451 (3)	0.0437 (8)
C6	0.5396 (3)	0.33407 (7)	0.5949 (2)	0.0364 (5)
C7	0.3900 (3)	0.33809 (7)	0.5140 (2)	0.0418 (6)
H7	0.3506	0.3130	0.4666	0.050*
C8	0.2990 (3)	0.37892 (7)	0.5032 (2)	0.0426 (6)
H8	0.1988	0.3811	0.4488	0.051*
C9	0.3553 (3)	0.41666 (7)	0.5724 (2)	0.0389 (6)
C10	0.5009 (3)	0.41290 (7)	0.6550 (2)	0.0480 (6)
H10	0.5384	0.4378	0.7040	0.058*
C11	0.5911 (3)	0.37188 (7)	0.6647 (2)	0.0462 (6)
H11	0.6901	0.3697	0.7204	0.055*
C12	0.3046 (3)	0.49521 (7)	0.6265 (2)	0.0484 (6)
H12A	0.4284	0.5039	0.6070	0.058*
H12B	0.2959	0.4889	0.7190	0.058*
C13	0.1778 (3)	0.53290 (7)	0.5904 (2)	0.0578 (7)
H13A	0.0545	0.5237	0.6096	0.069*
H13B	0.1863	0.5386	0.4976	0.069*
C14	0.2201 (4)	0.57630 (8)	0.6634 (3)	0.0768 (9)
H14A	0.2116	0.5708	0.7554	0.115*
H14B	0.1350	0.5995	0.6390	0.115*
H14C	0.3405	0.5861	0.6423	0.115*
N1	0.5477 (4)	0.2500	0.5790 (3)	0.0405 (7)
O1	1.0831 (3)	0.2500	0.5851 (2)	0.0633 (7)
O2	0.25630 (19)	0.45579 (5)	0.55359 (16)	0.0531 (5)
H1N	0.458 (4)	0.2500	0.623 (3)	0.029 (9)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0415 (12)	0.0358 (12)	0.0380 (13)	-0.0030 (10)	-0.0027 (10)	-0.0016 (10)
C2	0.0377 (13)	0.0352 (12)	0.0548 (16)	-0.0049 (10)	0.0003 (11)	-0.0056 (11)
C3	0.0497 (14)	0.0433 (13)	0.0492 (15)	0.0036 (11)	0.0091 (12)	0.0038 (12)
C4	0.055 (2)	0.049 (2)	0.0407 (19)	0.000	0.0047 (16)	0.000
C5	0.0354 (19)	0.049 (2)	0.047 (2)	0.000	0.0018 (16)	0.000
C6	0.0379 (12)	0.0346 (12)	0.0368 (13)	-0.0022 (10)	0.0005 (10)	-0.0019 (10)
C7	0.0445 (13)	0.0362 (13)	0.0445 (14)	-0.0045 (11)	-0.0023 (11)	-0.0051 (11)
C8	0.0439 (13)	0.0424 (14)	0.0415 (14)	-0.0020 (11)	-0.0081 (11)	-0.0027 (11)
C9	0.0413 (13)	0.0318 (13)	0.0436 (14)	-0.0017 (10)	0.0042 (11)	0.0011 (10)
C10	0.0508 (15)	0.0388 (14)	0.0543 (16)	-0.0033 (11)	-0.0069 (13)	-0.0129 (12)
C11	0.0435 (14)	0.0436 (14)	0.0515 (15)	0.0004 (11)	-0.0117 (12)	-0.0081 (12)
C12	0.0519 (14)	0.0357 (13)	0.0575 (15)	-0.0046 (11)	0.0027 (12)	-0.0054 (12)
C13	0.0594 (15)	0.0400 (14)	0.0740 (18)	0.0013 (12)	0.0019 (14)	-0.0047 (13)
C14	0.0743 (19)	0.0471 (15)	0.109 (3)	0.0032 (14)	-0.0007 (17)	-0.0183 (16)
N1	0.0371 (16)	0.0360 (16)	0.0483 (18)	0.000	0.0066 (15)	0.000
O1	0.0369 (14)	0.0686 (16)	0.0845 (18)	0.000	-0.0115 (13)	0.000
O2	0.0604 (10)	0.0345 (9)	0.0644 (11)	0.0053 (7)	-0.0124 (8)	-0.0074 (8)

Geometric parameters (\AA , $^\circ$)

C1—N1	1.464 (2)	C8—C9	1.381 (3)
C1—C6	1.507 (3)	C8—H8	0.9300
C1—C2	1.551 (3)	C9—C10	1.375 (3)
C1—H1	0.9800	C9—O2	1.376 (2)
C2—C5	1.502 (3)	C10—C11	1.381 (3)
C2—C3	1.540 (3)	C10—H10	0.9300
C2—H2	0.9800	C11—H11	0.9300
C3—C4	1.524 (3)	C12—O2	1.425 (2)
C3—H3A	0.9700	C12—C13	1.497 (3)
C3—H3B	0.9700	C12—H12A	0.9700
C4—C3 ⁱ	1.524 (3)	C12—H12B	0.9700
C4—H4A	0.9700	C13—C14	1.512 (3)
C4—H4B	0.9700	C13—H13A	0.9700
C5—O1	1.213 (3)	C13—H13B	0.9700
C5—C2 ⁱ	1.502 (3)	C14—H14A	0.9600
C6—C11	1.377 (3)	C14—H14B	0.9600
C6—C7	1.387 (3)	C14—H14C	0.9600
C7—C8	1.380 (3)	N1—C1 ⁱ	1.464 (2)
C7—H7	0.9300	N1—H1N	0.80 (3)
N1—C1—C6	112.97 (17)	C7—C8—H8	119.6
N1—C1—C2	108.69 (19)	C9—C8—H8	119.6
C6—C1—C2	112.25 (17)	C10—C9—O2	124.69 (19)
N1—C1—H1	107.6	C10—C9—C8	119.24 (19)
C6—C1—H1	107.6	O2—C9—C8	116.06 (19)

C2—C1—H1	107.6	C9—C10—C11	119.5 (2)
C5—C2—C3	107.6 (2)	C9—C10—H10	120.3
C5—C2—C1	107.73 (19)	C11—C10—H10	120.3
C3—C2—C1	115.30 (17)	C6—C11—C10	122.3 (2)
C5—C2—H2	108.7	C6—C11—H11	118.9
C3—C2—H2	108.7	C10—C11—H11	118.9
C1—C2—H2	108.7	O2—C12—C13	108.35 (18)
C4—C3—C2	114.2 (2)	O2—C12—H12A	110.0
C4—C3—H3A	108.7	C13—C12—H12A	110.0
C2—C3—H3A	108.7	O2—C12—H12B	110.0
C4—C3—H3B	108.7	C13—C12—H12B	110.0
C2—C3—H3B	108.7	H12A—C12—H12B	108.4
H3A—C3—H3B	107.6	C12—C13—C14	111.9 (2)
C3—C4—C3 ⁱ	114.1 (3)	C12—C13—H13A	109.2
C3—C4—H4A	108.7	C14—C13—H13A	109.2
C3 ⁱ —C4—H4A	108.7	C12—C13—H13B	109.2
C3—C4—H4B	108.7	C14—C13—H13B	109.2
C3 ⁱ —C4—H4B	108.7	H13A—C13—H13B	107.9
H4A—C4—H4B	107.6	C13—C14—H14A	109.5
O1—C5—C2 ⁱ	124.19 (13)	C13—C14—H14B	109.5
O1—C5—C2	124.19 (13)	H14A—C14—H14B	109.5
C2 ⁱ —C5—C2	111.6 (3)	C13—C14—H14C	109.5
C11—C6—C7	117.62 (19)	H14A—C14—H14C	109.5
C11—C6—C1	118.61 (18)	H14B—C14—H14C	109.5
C7—C6—C1	123.75 (19)	C1—N1—C1 ⁱ	112.2 (2)
C8—C7—C6	120.7 (2)	C1—N1—H1N	108.6 (10)
C8—C7—H7	119.7	C1 ⁱ —N1—H1N	108.6 (10)
C6—C7—H7	119.7	C9—O2—C12	118.21 (16)
C7—C8—C9	120.7 (2)		
N1—C1—C2—C5	-58.5 (2)	C1—C6—C7—C8	-177.0 (2)
C6—C1—C2—C5	175.77 (19)	C6—C7—C8—C9	0.2 (3)
N1—C1—C2—C3	61.6 (2)	C7—C8—C9—C10	-1.6 (3)
C6—C1—C2—C3	-64.1 (2)	C7—C8—C9—O2	179.35 (19)
C5—C2—C3—C4	52.0 (3)	O2—C9—C10—C11	-179.3 (2)
C1—C2—C3—C4	-68.2 (3)	C8—C9—C10—C11	1.7 (3)
C2—C3—C4—C3 ⁱ	-42.1 (3)	C7—C6—C11—C10	-0.9 (3)
C3—C2—C5—O1	115.5 (3)	C1—C6—C11—C10	177.2 (2)
C1—C2—C5—O1	-119.6 (3)	C9—C10—C11—C6	-0.5 (3)
C3—C2—C5—C2 ⁱ	-65.1 (3)	O2—C12—C13—C14	-179.8 (2)
C1—C2—C5—C2 ⁱ	59.8 (3)	C6—C1—N1—C1 ⁱ	-172.82 (15)
N1—C1—C6—C11	154.5 (2)	C2—C1—N1—C1 ⁱ	61.9 (3)
C2—C1—C6—C11	-82.1 (2)	C10—C9—O2—C12	-1.4 (3)
N1—C1—C6—C7	-27.5 (3)	C8—C9—O2—C12	177.64 (18)
C2—C1—C6—C7	95.8 (2)	C13—C12—O2—C9	-179.52 (18)
C11—C6—C7—C8	1.0 (3)		

Symmetry code: (i) $x, -y+1/2, z$.