

**6-Amino-3,4-dimethyl-4-phenyl-2H,4H-pyranopyrazole-5-carbonitrile**

**Nishith Saurav Topno,<sup>a</sup> Kandhasamy Kumaravel,<sup>b</sup> M. Kannan,<sup>a</sup> Gnanasambandam Vasuki<sup>b</sup> and R. Krishna<sup>a\*</sup>**

<sup>a</sup>Centre for Bioinformatics, Pondicherry University, Puducherry 605 014, India, and <sup>b</sup>Department of Chemistry, Pondicherry University, Puducherry 605 014, India

Correspondence e-mail: krishstrucbio@gmail.com


Received 16 February 2011; accepted 11 March 2011

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.057;  $wR$  factor = 0.173; data-to-parameter ratio = 27.3.

In the title compound,  $\text{C}_{15}\text{H}_{14}\text{N}_4\text{O}$ , the pyrazole ring is aligned at  $88.23(4)^\circ$  with respect to the aromatic ring and at  $3.75(4)^\circ$  with respect to the pyran ring. In the crystal,  $\text{N}-\text{H}\cdots\text{N}$  hydrogen bonds link adjacent molecules into a linear chain motif.  $\text{C}-\text{H}\cdots\text{N}$  interactions are also observed.

**Related literature**

For the synthesis, see: Vasuki & Kumaravel (2008). For the use of related compounds in organic synthesis, see: Liang *et al.* (2009). For related structures, see: Kannan *et al.* (2010).

**Experimental***Crystal data*

$\text{C}_{15}\text{H}_{14}\text{N}_4\text{O}$   
 $M_r = 266.3$   
Triclinic,  $P\bar{1}$   
 $a = 6.682(5)\text{ \AA}$   
 $b = 9.347(5)\text{ \AA}$

$c = 11.078(5)\text{ \AA}$   
 $\alpha = 99.213(5)^\circ$   
 $\beta = 102.740(5)^\circ$   
 $\gamma = 97.767(5)^\circ$   
 $V = 655.6(7)\text{ \AA}^3$

$Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.09\text{ mm}^{-1}$   
 $T = 293\text{ K}$   
 $0.22 \times 0.20 \times 0.16\text{ mm}$

*Data collection*

Bruker APEXII Kappa CCD detector diffractometer  
18424 measured reflections  
4996 independent reflections  
3169 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.036$

*Refinement*

$R[F^2 > 2\sigma(F^2)] = 0.057$   
 $wR(F^2) = 0.173$   
 $S = 0.99$   
4996 reflections  
183 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.30\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.29\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}3-\text{H}3\text{A}\cdots\text{N}1^{\text{i}}$	0.86	2.27	3.129 (2)	173
$\text{N}3-\text{H}3\text{B}\cdots\text{N}4^{\text{ii}}$	0.86	2.26	3.087 (2)	160
$\text{C}10-\text{H}10\cdots\text{N}4^{\text{iii}}$	0.93	2.53	3.455 (3)	172
$\text{C}14-\text{H}14\text{A}\cdots\text{N}1^{\text{iv}}$	0.96	2.59	3.522 (3)	163

Symmetry codes: (i)  $-x, -y + 1, -z + 1$ ; (ii)  $-x + 1, -y + 1, -z + 2$ ; (iii)  $-x + 2, -y + 2, -z + 2$ ; (iv)  $-x + 1, -y + 1, -z + 1$ .

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *PLATON* (Spek, 2009).

The authors acknowledge the Centre of Excellence in Bioinformatics, Pondicherry University, for providing the facilities for carrying out this work. RK and MK thank the University Grant Commission for providing a Major Research Project and Rajiv Gandhi National Fellowship, respectively. GV thanks the Department of Science and Technology, New Delhi, Government of India (grant No. SR/S5/GC-22/2007) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5120).

**References**

- Bruker (2004). *APEX2, SAINT* and *XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Kannan, M., Kumaravel, K., Vasuki, G. & Krishna, R. (2010). *Acta Cryst. E66*, o1242.
- Liang, F., Cheng, X., Liu, J. & Liu, Q. (2009). *Chem. Commun.* pp. 3636–3538.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Vasuki, G. & Kumaravel, K. (2008). *Tetrahedron Lett.* **49**, 5636–5638.

# supporting information

*Acta Cryst.* (2011). E67, o956 [doi:10.1107/S1600536811009354]

## 6-Amino-3,4-dimethyl-4-phenyl-2*H*,4*H*-pyrano[2,3-*c*]pyrazole-5-carbonitrile

**Nishith Saurav Topno, Kandhasamy Kumaravel, M. Kannan, Gnanasambandam Vasuki and R. Krishna**

### S1. Comment

The pyrano pyrazole derivatives are widely used as organic intermediates (Liang *et al.*, 2009) and are well known for their biological contribution as ChK1 inhibitors. In view of the growing importance of the pyrano pyrazole derivatives, we have synthesized the title compound by Rapid four-component reactions in water (Vasuki & Kumaravel, 2008) and the single-crystal structure analysis was undertaken.


In the title compound, the attached benzyl ring makes the dihedral angle of 88.23 (4) ° and orient in (+)-*syn*-clinal conformation with pyrazole ring, whereas the fused pyrazole ring makes 3.75 (4) ° dihedral angle and orient by an (+)-*syn*-periplanar conformation with respect to the pyran ring (Fig. 1). The methyl group is attached to C6 atom of pyran with an angle of 109.29 (11) °. Two N—H···N and C—H···N intermolecular hydrogen bond interactions (Fig. 2) are observed for maintaining the crystal packing (Fig. 3), in which the N3—H···N4 intermolecular interaction are observed to form  $R_2^2$  (12) ring motifs (Fig. 4). The weak N—H···π intermolecular interaction is also observed for the stabilization of the crystal packing, with a bond distance of 3.382 (3) Å (Fig. 5).

### S2. Experimental


The title compound was prepared by the successive addition of acetophenone 2 (0.240 g, 2 mmol), malononitrile (0.132 g, 2 mmol) and piperidine (5 mol%) to a stirred aqueous mixture of hydrazine hydrate 96% 1 (0.107 g, 2 mmol) and ethyl acetoacetate 2 (0.520 g, 2 mmol) at room temperature under an open atmosphere with vigorous stirring for 5–10 min. The precipitated solid was then filtered, followed by washing with water and then with a mixture of ethyl acetate/hexane (20:80). The product obtained was pure by TLC and  ${}^1\text{H}$  NMR spectroscopy. Nevertheless, the products were further purified by recrystallization from ethanol. Analysis calculated for 6-amino-3,4-dimethyl-4-phenyl-2*H*,4*H*-pyrano[2,3-*c*]pyrazole-5-carbonitrile showed that it has  $\text{C}_{15}\text{H}_{14}\text{N}_4\text{O}$ .

### S3. Refinement


All hydrogen atoms were placed in calculated positions, with N—H=0.86 and C—H=0.93–0.97 and included in the final cycles of refinement using a riding model with  $U(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ .

**Figure 1**


The molecular structure of (I), showing the atom-numbering scheme and displacement ellipsoids drawn at the 50% probability level.

**Figure 2**


The crystal packing of (I), showing intermolecular hydrogen bonding interactions as dashed lines.

**Figure 3**

The crystal packing of Compound (I) viewed down the  $XO$ -axis, showing intermolecular hydrogen bonding interactions as dashed lines.

**Figure 4**

A view of  $R_2^2(12)$  ring motifs formed by N—H···N interaction between two molecules. The ring forming atoms are shown in ball and stick model and the Hydrogen bond are shown in green dashed lines.

**Figure 5**

The molecular interaction showing the weak N—H···pi interaction in Compound (I). Cg is a centroid of C<sub>8</sub>—C<sub>13</sub> ring.

### 6-Amino-3,4-dimethyl-4-phenyl-2*H*,4*H*-pyrano[2,3-*c*]pyrazole- 5-carbonitrile

#### Crystal data

C<sub>15</sub>H<sub>14</sub>N<sub>4</sub>O

$M_r = 266.3$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a = 6.682 (5) \text{ \AA}$

$b = 9.347 (5) \text{ \AA}$

$c = 11.078 (5) \text{ \AA}$

$\alpha = 99.213 (5)^\circ$

$\beta = 102.740 (5)^\circ$

$\gamma = 97.767 (5)^\circ$

$V = 655.6 (7) \text{ \AA}^3$   
 $Z = 2$   
 $F(000) = 280$   
 $D_x = 1.349 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71069 \text{ \AA}$   
Cell parameters from 4350 reflections

$\theta = 1.9\text{--}33.3^\circ$   
 $\mu = 0.09 \text{ mm}^{-1}$   
 $T = 293 \text{ K}$   
Prism, colourless  
 $0.22 \times 0.20 \times 0.16 \text{ mm}$

#### Data collection

Bruker APEXII Kappa CCD-detector  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
Detector resolution: 0 pixels  $\text{mm}^{-1}$   
 $\omega$  and  $\varphi$  scans  
18424 measured reflections

4996 independent reflections  
3169 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.036$   
 $\theta_{\text{max}} = 33.3^\circ, \theta_{\text{min}} = 1.9^\circ$   
 $h = -10 \rightarrow 9$   
 $k = -14 \rightarrow 14$   
 $l = -17 \rightarrow 16$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.057$   
 $wR(F^2) = 0.173$   
 $S = 0.99$   
4996 reflections  
183 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0847P)^2 + 0.1309P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.014$   
 $\Delta\rho_{\text{max}} = 0.30 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.29 \text{ e \AA}^{-3}$

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.22065 (13)	0.57205 (11)	0.58851 (9)	0.0380 (2)
C2	0.52515 (18)	0.74174 (14)	0.57658 (11)	0.0306 (3)
N1	0.23543 (18)	0.67259 (14)	0.41251 (11)	0.0406 (3)
C6	0.64969 (18)	0.74989 (14)	0.70869 (12)	0.0304 (2)
C5	0.52503 (19)	0.63581 (14)	0.76101 (12)	0.0315 (3)
C7	0.6205 (2)	0.60943 (15)	0.88018 (13)	0.0377 (3)
C4	0.32984 (19)	0.55767 (14)	0.70373 (12)	0.0315 (3)
C3	0.32497 (19)	0.66098 (15)	0.52808 (12)	0.0320 (3)
C8	0.6664 (2)	0.90474 (14)	0.78630 (12)	0.0332 (3)
N3	0.21879 (18)	0.45814 (14)	0.74817 (12)	0.0439 (3)
H3A	0.0968	0.4142	0.7044	0.053*
H3B	0.2692	0.4379	0.8206	0.053*

N2	0.38757 (19)	0.76374 (15)	0.38430 (11)	0.0432 (3)
H2	0.3736	0.7912	0.3129	0.052*
N4	0.7004 (2)	0.59058 (17)	0.97741 (14)	0.0571 (4)
C1	0.5621 (2)	0.80702 (16)	0.47848 (13)	0.0376 (3)
C14	0.8640 (2)	0.70915 (18)	0.70847 (16)	0.0438 (3)
H14A	0.8467	0.6116	0.6599	0.066*
H14B	0.9397	0.7124	0.7936	0.066*
H14C	0.9401	0.7779	0.6718	0.066*
C13	0.4882 (2)	0.95084 (17)	0.81184 (15)	0.0442 (3)
H13	0.3617	0.8859	0.7841	0.053*
C9	0.8513 (3)	1.00406 (18)	0.82993 (17)	0.0517 (4)
H9	0.9738	0.9768	0.8150	0.062*
C10	0.8558 (3)	1.1452 (2)	0.89631 (18)	0.0639 (5)
H10	0.9817	1.2109	0.9251	0.077*
C15	0.7477 (3)	0.9053 (2)	0.46603 (16)	0.0536 (4)
H15A	0.8448	0.8469	0.4411	0.080*
H15B	0.8130	0.9701	0.5456	0.080*
H15C	0.7051	0.9625	0.4033	0.080*
C12	0.4943 (3)	1.0905 (2)	0.87732 (17)	0.0568 (4)
H12	0.3727	1.1186	0.8930	0.068*
C11	0.6797 (4)	1.18818 (19)	0.91953 (16)	0.0595 (5)
H11	0.6843	1.2826	0.9635	0.071*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0256 (4)	0.0500 (6)	0.0318 (5)	-0.0082 (4)	-0.0019 (3)	0.0142 (4)
C2	0.0260 (5)	0.0314 (6)	0.0302 (6)	-0.0015 (4)	0.0032 (4)	0.0049 (4)
N1	0.0346 (6)	0.0480 (7)	0.0319 (6)	-0.0072 (5)	-0.0007 (4)	0.0116 (5)
C6	0.0233 (5)	0.0309 (6)	0.0322 (6)	-0.0011 (4)	0.0012 (4)	0.0056 (4)
C5	0.0272 (5)	0.0296 (6)	0.0320 (6)	-0.0019 (4)	-0.0014 (4)	0.0073 (5)
C7	0.0308 (6)	0.0350 (7)	0.0397 (7)	-0.0061 (5)	-0.0026 (5)	0.0109 (5)
C4	0.0266 (5)	0.0338 (6)	0.0303 (6)	0.0004 (4)	0.0013 (4)	0.0072 (5)
C3	0.0273 (5)	0.0362 (6)	0.0290 (6)	-0.0015 (4)	0.0029 (4)	0.0073 (5)
C8	0.0330 (6)	0.0318 (6)	0.0286 (6)	-0.0033 (5)	-0.0005 (5)	0.0066 (5)
N3	0.0331 (6)	0.0509 (7)	0.0396 (6)	-0.0111 (5)	-0.0038 (5)	0.0191 (5)
N2	0.0415 (6)	0.0517 (7)	0.0308 (6)	-0.0082 (5)	0.0030 (5)	0.0144 (5)
N4	0.0472 (7)	0.0635 (9)	0.0477 (8)	-0.0143 (6)	-0.0114 (6)	0.0260 (7)
C1	0.0327 (6)	0.0407 (7)	0.0352 (7)	-0.0031 (5)	0.0057 (5)	0.0075 (5)
C14	0.0271 (6)	0.0477 (8)	0.0529 (9)	0.0045 (5)	0.0052 (6)	0.0076 (7)
C13	0.0464 (8)	0.0385 (8)	0.0450 (8)	0.0025 (6)	0.0122 (6)	0.0034 (6)
C9	0.0384 (7)	0.0422 (8)	0.0600 (10)	-0.0085 (6)	-0.0025 (7)	0.0016 (7)
C10	0.0682 (12)	0.0430 (9)	0.0577 (11)	-0.0165 (8)	-0.0087 (9)	-0.0007 (8)
C15	0.0450 (8)	0.0609 (10)	0.0498 (9)	-0.0142 (7)	0.0090 (7)	0.0193 (8)
C12	0.0746 (12)	0.0457 (9)	0.0514 (10)	0.0136 (8)	0.0213 (9)	0.0032 (7)
C11	0.0922 (14)	0.0371 (8)	0.0405 (8)	0.0039 (9)	0.0083 (9)	-0.0001 (6)

Geometric parameters ( $\text{\AA}$ ,  $\text{^{\circ}}$ )

O1—C4	1.3602 (16)	N2—C1	1.3465 (19)
O1—C3	1.3611 (16)	N2—H2	0.8600
C2—C1	1.3799 (19)	C1—C15	1.486 (2)
C2—C3	1.3885 (18)	C14—H14A	0.9600
C2—C6	1.5001 (18)	C14—H14B	0.9600
N1—C3	1.3164 (17)	C14—H14C	0.9600
N1—N2	1.3595 (17)	C13—C12	1.379 (2)
C6—C5	1.5289 (18)	C13—H13	0.9300
C6—C14	1.531 (2)	C9—C10	1.397 (3)
C6—C8	1.5369 (19)	C9—H9	0.9300
C5—C4	1.3661 (18)	C10—C11	1.357 (3)
C5—C7	1.4091 (19)	C10—H10	0.9300
C7—N4	1.1440 (19)	C15—H15A	0.9600
C4—N3	1.3340 (17)	C15—H15B	0.9600
C8—C9	1.380 (2)	C15—H15C	0.9600
C8—C13	1.390 (2)	C12—C11	1.374 (3)
N3—H3A	0.8600	C12—H12	0.9300
N3—H3B	0.8600	C11—H11	0.9300
C4—O1—C3	115.59 (10)	N2—C1—C2	106.17 (12)
C1—C2—C3	103.33 (11)	N2—C1—C15	122.22 (14)
C1—C2—C6	133.30 (11)	C2—C1—C15	131.61 (13)
C3—C2—C6	123.33 (11)	C6—C14—H14A	109.5
C3—N1—N2	101.74 (11)	C6—C14—H14B	109.5
C2—C6—C5	105.36 (10)	H14A—C14—H14B	109.5
C2—C6—C14	110.78 (11)	C6—C14—H14C	109.5
C5—C6—C14	109.28 (12)	H14A—C14—H14C	109.5
C2—C6—C8	109.09 (10)	H14B—C14—H14C	109.5
C5—C6—C8	109.98 (11)	C12—C13—C8	121.64 (15)
C14—C6—C8	112.12 (10)	C12—C13—H13	119.2
C4—C5—C7	116.92 (12)	C8—C13—H13	119.2
C4—C5—C6	126.31 (11)	C8—C9—C10	120.50 (17)
C7—C5—C6	116.77 (11)	C8—C9—H9	119.8
N4—C7—C5	178.69 (15)	C10—C9—H9	119.8
N3—C4—O1	110.00 (11)	C11—C10—C9	121.11 (17)
N3—C4—C5	127.07 (12)	C11—C10—H10	119.4
O1—C4—C5	122.93 (12)	C9—C10—H10	119.4
N1—C3—O1	119.33 (11)	C1—C15—H15A	109.5
N1—C3—C2	114.94 (12)	C1—C15—H15B	109.5
O1—C3—C2	125.71 (12)	H15A—C15—H15B	109.5
C9—C8—C13	117.40 (14)	C1—C15—H15C	109.5
C9—C8—C6	123.01 (13)	H15A—C15—H15C	109.5
C13—C8—C6	119.57 (11)	H15B—C15—H15C	109.5
C4—N3—H3A	120.0	C11—C12—C13	120.16 (18)
C4—N3—H3B	120.0	C11—C12—H12	119.9
H3A—N3—H3B	120.0	C13—C12—H12	119.9

C1—N2—N1	113.80 (12)	C10—C11—C12	119.19 (17)
C1—N2—H2	123.1	C10—C11—H11	120.4
N1—N2—H2	123.1	C12—C11—H11	120.4
C1—C2—C6—C5	−173.79 (14)	C6—C2—C3—N1	176.69 (12)
C3—C2—C6—C5	8.91 (17)	C1—C2—C3—O1	177.46 (13)
C1—C2—C6—C14	−55.7 (2)	C6—C2—C3—O1	−4.6 (2)
C3—C2—C6—C14	126.98 (14)	C2—C6—C8—C9	−111.80 (15)
C1—C2—C6—C8	68.17 (18)	C5—C6—C8—C9	133.10 (14)
C3—C2—C6—C8	−109.13 (14)	C14—C6—C8—C9	11.29 (19)
C2—C6—C5—C4	−7.33 (18)	C2—C6—C8—C13	66.30 (15)
C14—C6—C5—C4	−126.40 (15)	C5—C6—C8—C13	−48.80 (16)
C8—C6—C5—C4	110.12 (15)	C14—C6—C8—C13	−170.61 (13)
C2—C6—C5—C7	173.06 (12)	C3—N1—N2—C1	−0.42 (17)
C14—C6—C5—C7	53.99 (16)	N1—N2—C1—C2	−0.34 (18)
C8—C6—C5—C7	−69.50 (15)	N1—N2—C1—C15	179.23 (15)
C4—C5—C7—N4	−158 (8)	C3—C2—C1—N2	0.91 (15)
C6—C5—C7—N4	21 (8)	C6—C2—C1—N2	−176.77 (14)
C3—O1—C4—N3	−174.36 (12)	C3—C2—C1—C15	−178.61 (17)
C3—O1—C4—C5	5.24 (19)	C6—C2—C1—C15	3.7 (3)
C7—C5—C4—N3	−0.2 (2)	C9—C8—C13—C12	0.5 (2)
C6—C5—C4—N3	−179.80 (13)	C6—C8—C13—C12	−177.74 (14)
C7—C5—C4—O1	−179.72 (12)	C13—C8—C9—C10	−0.4 (2)
C6—C5—C4—O1	0.7 (2)	C6—C8—C9—C10	177.75 (15)
N2—N1—C3—O1	−177.77 (12)	C8—C9—C10—C11	0.0 (3)
N2—N1—C3—C2	1.06 (17)	C8—C13—C12—C11	−0.2 (3)
C4—O1—C3—N1	175.29 (12)	C9—C10—C11—C12	0.3 (3)
C4—O1—C3—C2	−3.4 (2)	C13—C12—C11—C10	−0.2 (3)
C1—C2—C3—N1	−1.29 (17)		

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N3—H3A···N1 <sup>i</sup>	0.86	2.27	3.129 (2)	173
N3—H3B···N4 <sup>ii</sup>	0.86	2.26	3.087 (2)	160
C10—H10···N4 <sup>iii</sup>	0.93	2.53	3.455 (3)	172
C14—H14A···N1 <sup>iv</sup>	0.96	2.59	3.522 (3)	163

Symmetry codes: (i)  $-x, -y+1, -z+1$ ; (ii)  $-x+1, -y+1, -z+2$ ; (iii)  $-x+2, -y+2, -z+2$ ; (iv)  $-x+1, -y+1, -z+1$ .