

Tetraqua(nitrato- κ^2O,O')bis(pyridinium-4-carboxylate- κO)europium(III) dinitrate

Zhi-Guo Zhong,^{a*} Jin-Fan Song,^b Jing Li^c and Zhao-Hui Meng^d

^aCenter of Analysis and Testing, Nanyang Normal University, Nanyang 473061, People's Republic of China, ^bSchool of Physics and Electronic Engineering, Nanyang Normal University, Nanyang 473061, People's Republic of China, ^cSchool of Mathematics and Statistics, Nanyang Normal University, Nanyang 473061, People's Republic of China, and ^dCollege of Chemistry and Pharmacy Engineering, Nanyang Normal University, Nanyang 473061, People's Republic of China
Correspondence e-mail: zhongzhiguo1978@126.com


Received 7 April 2011; accepted 20 April 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.015; wR factor = 0.038; data-to-parameter ratio = 10.4.

The asymmetric unit of the title compound, $[\text{Eu}(\text{NO}_3)(\text{C}_6\text{H}_5\text{NO}_2)_2(\text{H}_2\text{O})_4](\text{NO}_3)_2$, consists of one-half of the C_2 symmetric coordination cation and one nitrate anion. The eight-coordinated Eu^{III} atom is in a distorted dodecahedral coordination environment. The coordination cations and nitrate anions are connected via $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds into a three-dimensional network.

Related literature

For photophysical properties of lanthanide(III) coordination compounds, see, for example: Jüstel *et al.* (1998); Xu *et al.* (2010). For potential applications of lanthanide(III) coordination compounds as light-conversion molecular devices, see, for example: Lehn (1990).



Experimental

Crystal data

$[\text{Eu}(\text{NO}_3)(\text{C}_6\text{H}_5\text{NO}_2)_2(\text{H}_2\text{O})_4]\cdot(\text{NO}_3)_2$	$\beta = 118.728 (4)^\circ$
$M_r = 656.27$	$V = 2136.6 (11)\text{ \AA}^3$
Monoclinic, $C2/c$	$Z = 4$
$a = 14.612 (4)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 12.498 (4)\text{ \AA}$	$\mu = 3.03\text{ mm}^{-1}$
$c = 13.342 (4)\text{ \AA}$	$T = 293\text{ K}$
	$0.35 \times 0.32 \times 0.28\text{ mm}$

Data collection

Bruker APEXII CCD diffractometer	5224 measured reflections
Absorption correction: multi-scan <i>SADABS</i> (Bruker, 1997)	1881 independent reflections
$T_{\min} = 0.417$, $T_{\max} = 0.484$	1847 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.017$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.015$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.038$	$\Delta\rho_{\text{max}} = 0.58\text{ e \AA}^{-3}$
$S = 1.02$	$\Delta\rho_{\text{min}} = -0.62\text{ e \AA}^{-3}$
1881 reflections	
180 parameters	
8 restraints	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1W—H1WB···O7 ⁱ	0.81 (2)	1.99 (2)	2.790 (2)	175 (3)
O1W—H1WA···O1 ⁱⁱ	0.83 (2)	1.87 (2)	2.653 (2)	155 (2)
O2W—H2WB···O1 ⁱⁱⁱ	0.83 (2)	1.84 (2)	2.661 (2)	173 (3)
O2W—H2WA···O5	0.82 (2)	2.23 (2)	2.958 (3)	148 (3)
N1—H6···O7 ^{iv}	0.94 (3)	1.88 (3)	2.814 (3)	179 (3)

Symmetry codes: (i) $x - \frac{1}{2}, y - \frac{1}{2}, z$; (ii) $-x, y, -z + \frac{1}{2}$; (iii) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$; (iv) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINT* (Bruker, 1997); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2366).

References

- Bruker (1997). *SMART, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Jüstel, T., Nikol, H. & Ronda, C. (1998). *Angew. Chem. Int. Ed.* **37**, 3084–3103.
- Lehn, J. M. (1990). *Angew. Chem. Int. Ed.* **29**, 1304–1319.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Xu, H., Wei, Y., Zhao, B. M. & Huang, W. (2010). *J. Rare Earths*, **28**, 666–670.

supporting information

Acta Cryst. (2011). E67, m637 [doi:10.1107/S1600536811014772]

Tetraaqua(nitrato- κ^2O,O')bis(pyridinium-4-carboxylate- κO)europium(III) dinitrate

Zhi-Guo Zhong, Jin-Fan Song, Jing Li and Zhao-Hui Meng

S1. Comment


The lanthanide (III) coordination compounds have attracted considerable attention due to their interesting photophysical properties and their potential application as light-conversion molecular devices. Herein, we report a new rare-earth metal–organic compound $[\text{Eu}(\text{C}_6\text{H}_5\text{NO}_2)_2(\text{H}_2\text{O})_4(\text{NO}_3)](\text{NO}_3)_2$. The structural unit of the title compound consists of one coordination cation which has a crystallographic twofold axis symmetry, $[\text{Eu}(\text{C}_6\text{H}_5\text{NO}_2)_2(\text{H}_2\text{O})_4(\text{NO}_3)]^{2+}$, and two nitrate anions (Fig. 1). In the coordination cation the Eu(III) center is coordinated by eight O atoms: two from $\text{C}_6\text{H}_5\text{NO}_2$ ligands, two from NO_3^- anion and four from water molecules. The coordination geometry around the Eu(III) center can be described as dodecahedral with O—Eu—O bond angles ranging from 51.07 (10) to 152.96 (9) $^\circ$ and the Eu—O bond lengths ranging from 2.3614 (16) to 2.5000 (19) Å. The electrostatic interactions and hydrogen bonds result in the formation of three-dimensional network (Fig. 2). Obviously, electrostatic interactions and hydrogen bonds play a crucial role in the chemical stability of the title compound.

S2. Experimental


All chemicals were of reagent grade quality obtained from commercial sources and used without further purification. 0.24 g of isonicotinic acid (1 mmol) and 0.45 g $\text{Eu}(\text{NO}_3)_3 \cdot 6\text{H}_2\text{O}$ (1 mmol) were dissolved in 30 ml of distilled water. Then pH of the mixture was carefully adjusted to 5.0 with 1M HCl solution. After stirring for half an hour, the solution was filtered and left for slowly evaporation at room temperature to obtain colorless crystals suitable for X-ray structure determination.

S3. Refinement

The H atoms bonded to C were positioned geometrically and refined using a riding model, with C—H = 0.93 Å and with $U_{\text{iso}}(\text{H}) = 1.2$ times $U_{\text{eq}}(\text{C})$. The H atoms bonded to O atoms were located from Fourier difference maps and refined with distance restraints of O1W—H1WA = 0.82, O1W—H1WB = 0.82, O2W—H2WA = 0.82, O2W—H2WB = 0.82, H1WA…H1WB = 1.36, H2WA…H2WB = 1.36 and H1WA…Eu1 = 2.85 Å. The H atom bonded to N atom was located from Fourier difference maps and freely refined. In addition, the O4 and N2 atoms were refined with SHELXL97 restraint 'DELUdelu 0.01'.

**Figure 1**

View of the title compound with displacement ellipsoids drawn at the 30% probability level . Symmetry code for the atoms with the label A: $-x, y, -z + 1/2$.

**Figure 2**

Crystal packing viewed along the *c* axis. Hydrogen bonds are shown with dashed lines.

Tetraqua(nitrato- κ^2O,O')bis(pyridinium-4-carboxylate- κO)europium(III) dinitrate

Crystal data



$M_r = 656.27$

Monoclinic, $C2/c$

Hall symbol: -C 2yc

$a = 14.612 (4)$ Å

$b = 12.498 (4)$ Å

$c = 13.342 (4)$ Å

$\beta = 118.728 (4)^\circ$

$V = 2136.6 (11)$ Å³

$Z = 4$

$F(000) = 1296$

$D_x = 2.040 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5781 reflections

$\theta = 2.3\text{--}28.3^\circ$

$\mu = 3.03 \text{ mm}^{-1}$

$T = 293$ K

Block, colourless

$0.35 \times 0.32 \times 0.28$ mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

SADABS (Bruker, 1997)

$T_{\min} = 0.417$, $T_{\max} = 0.484$

5224 measured reflections

1881 independent reflections

1847 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.3^\circ$

$h = -13 \rightarrow 17$

$k = -14 \rightarrow 12$

$l = -15 \rightarrow 15$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.015$$

$$wR(F^2) = 0.038$$

$$S = 1.02$$

1881 reflections

180 parameters

8 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0211P)^2 + 2.5852P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.58 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.62 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Eu1	0.0000	0.184699 (10)	0.2500	0.02398 (6)
O1	0.27428 (13)	0.30594 (12)	0.37370 (16)	0.0421 (4)
O1W	-0.17399 (12)	0.12138 (14)	0.15960 (17)	0.0453 (4)
H1WB	-0.199 (2)	0.0622 (15)	0.144 (2)	0.044 (8)*
H1WA	-0.2202 (13)	0.1684 (17)	0.134 (3)	0.063 (10)*
O2	0.10558 (12)	0.33068 (12)	0.25718 (14)	0.0350 (4)
O2W	-0.04309 (13)	0.22924 (14)	0.05884 (14)	0.0390 (4)
H2WB	-0.0982 (17)	0.213 (2)	0.002 (2)	0.051 (8)*
H2WA	-0.021 (2)	0.2840 (19)	0.045 (3)	0.061 (10)*
O3	0.00037 (14)	0.00420 (15)	0.33097 (17)	0.0495 (4)
O4	0.0000	-0.1458 (2)	0.2500	0.1018 (15)
O5	0.08162 (14)	0.35758 (14)	-0.01819 (16)	0.0462 (4)
O6	0.11039 (14)	0.52783 (13)	-0.00657 (16)	0.0471 (4)
O7	0.23063 (12)	0.42067 (12)	0.10876 (14)	0.0386 (4)
N1	0.23834 (19)	0.69898 (16)	0.3531 (2)	0.0390 (5)
N2	0.0000	-0.0487 (2)	0.2500	0.0506 (9)
N3	0.13887 (14)	0.43631 (14)	0.02637 (16)	0.0312 (4)
C1	0.3165 (2)	0.63532 (19)	0.4224 (2)	0.0403 (5)
H1	0.3784	0.6647	0.4788	0.048*
C2	0.30538 (18)	0.52613 (18)	0.4104 (2)	0.0353 (5)
H2	0.3604	0.4813	0.4567	0.042*
C3	0.21208 (16)	0.48366 (16)	0.32907 (17)	0.0262 (4)
C4	0.13218 (17)	0.55193 (17)	0.25749 (19)	0.0317 (5)
H4	0.0691	0.5246	0.2012	0.038*
C5	0.1479 (2)	0.65997 (19)	0.2713 (2)	0.0387 (5)
H5	0.0954	0.7066	0.2235	0.046*
C6	0.19639 (16)	0.36402 (16)	0.31971 (17)	0.0270 (4)
H6	0.248 (2)	0.773 (3)	0.367 (2)	0.050 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Eu1	0.01991 (9)	0.01759 (9)	0.02573 (9)	0.000	0.00400 (6)	0.000
O1	0.0282 (8)	0.0270 (8)	0.0463 (10)	0.0018 (6)	-0.0019 (8)	0.0010 (7)
O1W	0.0250 (8)	0.0221 (9)	0.0734 (13)	-0.0035 (7)	0.0112 (8)	-0.0035 (8)
O2	0.0258 (8)	0.0258 (8)	0.0379 (9)	-0.0057 (6)	0.0028 (7)	0.0032 (6)
O2W	0.0332 (9)	0.0382 (10)	0.0296 (8)	-0.0127 (8)	0.0023 (7)	0.0047 (7)
O3	0.0463 (10)	0.0388 (10)	0.0516 (11)	-0.0004 (8)	0.0140 (9)	0.0148 (8)
O4	0.088 (3)	0.0177 (14)	0.202 (5)	0.000	0.071 (3)	0.000
O5	0.0419 (10)	0.0339 (9)	0.0486 (10)	-0.0067 (8)	0.0104 (8)	-0.0042 (8)
O6	0.0487 (10)	0.0281 (9)	0.0581 (11)	0.0106 (8)	0.0207 (9)	0.0114 (8)
O7	0.0331 (8)	0.0291 (8)	0.0428 (9)	0.0014 (6)	0.0098 (7)	0.0034 (7)
N1	0.0544 (13)	0.0233 (10)	0.0466 (12)	-0.0076 (9)	0.0302 (11)	-0.0047 (8)
N2	0.0305 (15)	0.0215 (14)	0.083 (3)	0.000	0.0135 (16)	0.000
N3	0.0333 (10)	0.0281 (10)	0.0337 (10)	0.0021 (8)	0.0172 (8)	0.0016 (8)
C1	0.0452 (14)	0.0326 (13)	0.0373 (13)	-0.0140 (11)	0.0151 (11)	-0.0061 (10)
C2	0.0327 (11)	0.0286 (12)	0.0336 (12)	-0.0056 (9)	0.0070 (10)	0.0001 (9)
C3	0.0277 (10)	0.0254 (10)	0.0253 (10)	-0.0023 (8)	0.0127 (9)	-0.0003 (8)
C4	0.0275 (10)	0.0288 (11)	0.0359 (12)	0.0009 (9)	0.0127 (9)	0.0017 (9)
C5	0.0438 (14)	0.0271 (11)	0.0479 (14)	0.0064 (10)	0.0242 (12)	0.0051 (10)
C6	0.0255 (10)	0.0237 (11)	0.0239 (10)	-0.0034 (8)	0.0056 (8)	-0.0015 (8)

Geometric parameters (\AA , ^\circ)

Eu1—O2	2.3614 (16)	O5—N3	1.242 (3)
Eu1—O2 ⁱ	2.3614 (16)	O6—N3	1.224 (2)
Eu1—O1W	2.3657 (17)	O7—N3	1.276 (2)
Eu1—O1W ⁱ	2.3657 (17)	N1—C1	1.333 (4)
Eu1—O2W ⁱ	2.3806 (18)	N1—C5	1.337 (4)
Eu1—O2W	2.3806 (18)	N1—H6	0.94 (3)
Eu1—O3 ⁱ	2.5000 (19)	N2—O3 ⁱ	1.264 (2)
Eu1—O3	2.5000 (19)	C1—C2	1.375 (3)
Eu1—N2	2.917 (3)	C1—H1	0.9300
O1—C6	1.246 (3)	C2—C3	1.377 (3)
O1W—H1WB	0.805 (17)	C2—H2	0.9300
O1W—H1WA	0.834 (16)	C3—C4	1.390 (3)
O2—C6	1.251 (3)	C3—C6	1.509 (3)
O2W—H2WB	0.826 (17)	C4—C5	1.367 (3)
O2W—H2WA	0.818 (18)	C4—H4	0.9300
O3—N2	1.264 (2)	C5—H5	0.9300
O4—N2	1.214 (4)		
O2—Eu1—O2 ⁱ	78.82 (8)	Eu1—O1W—H1WA	115.7 (15)
O2—Eu1—O1W	143.15 (6)	H1WB—O1W—H1WA	111 (2)
O2 ⁱ —Eu1—O1W	73.54 (6)	C6—O2—Eu1	138.33 (14)
O2—Eu1—O1W ⁱ	73.54 (6)	Eu1—O2W—H2WB	125 (2)
O2 ⁱ —Eu1—O1W ⁱ	143.15 (6)	Eu1—O2W—H2WA	121 (2)

O1W—Eu1—O1W ⁱ	140.91 (8)	H2WB—O2W—H2WA	109 (3)
O2—Eu1—O2W ⁱ	86.83 (6)	N2—O3—Eu1	95.99 (15)
O2 ⁱ —Eu1—O2W ⁱ	72.18 (6)	C1—N1—C5	122.0 (2)
O1W—Eu1—O2W ⁱ	106.89 (7)	C1—N1—H6	116.8 (18)
O1W ⁱ —Eu1—O2W ⁱ	82.30 (7)	C5—N1—H6	121.2 (18)
O2—Eu1—O2W	72.18 (6)	O4—N2—O3	121.53 (14)
O2 ⁱ —Eu1—O2W	86.83 (6)	O4—N2—O3 ⁱ	121.53 (14)
O1W—Eu1—O2W	82.30 (7)	O3—N2—O3 ⁱ	116.9 (3)
O1W ⁱ —Eu1—O2W	106.89 (7)	O4—N2—Eu1	180.0
O2W ⁱ —Eu1—O2W	152.96 (9)	O3—N2—Eu1	58.47 (14)
O2—Eu1—O3 ⁱ	125.48 (7)	O3 ⁱ —N2—Eu1	58.47 (14)
O2 ⁱ —Eu1—O3 ⁱ	144.49 (6)	O6—N3—O5	122.3 (2)
O1W—Eu1—O3 ⁱ	72.50 (6)	O6—N3—O7	119.18 (19)
O1W ⁱ —Eu1—O3 ⁱ	72.37 (6)	O5—N3—O7	118.56 (18)
O2W ⁱ —Eu1—O3 ⁱ	128.20 (6)	N1—C1—C2	119.9 (2)
O2W—Eu1—O3 ⁱ	78.67 (7)	N1—C1—H1	120.1
O2—Eu1—O3	144.49 (6)	C2—C1—H1	120.1
O2 ⁱ —Eu1—O3	125.48 (7)	C1—C2—C3	119.4 (2)
O1W—Eu1—O3	72.37 (6)	C1—C2—H2	120.3
O1W ⁱ —Eu1—O3	72.50 (6)	C3—C2—H2	120.3
O2W ⁱ —Eu1—O3	78.67 (7)	C2—C3—C4	119.4 (2)
O2W—Eu1—O3	128.20 (6)	C2—C3—C6	120.06 (19)
O3 ⁱ —Eu1—O3	51.07 (10)	C4—C3—C6	120.50 (19)
O2—Eu1—N2	140.59 (4)	C5—C4—C3	118.8 (2)
O2 ⁱ —Eu1—N2	140.59 (4)	C5—C4—H4	120.6
O1W—Eu1—N2	70.46 (4)	C3—C4—H4	120.6
O1W ⁱ —Eu1—N2	70.46 (4)	N1—C5—C4	120.4 (2)
O2W ⁱ —Eu1—N2	103.52 (4)	N1—C5—H5	119.8
O2W—Eu1—N2	103.52 (4)	C4—C5—H5	119.8
O3 ⁱ —Eu1—N2	25.54 (5)	O1—C6—O2	124.92 (19)
O3—Eu1—N2	25.54 (5)	O1—C6—C3	117.98 (18)
Eu1—O1W—H1WB	133 (2)	O2—C6—C3	117.09 (18)

Symmetry code: (i) $-x, y, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

D—H···A	D—H	H···A	D···A	D—H···A
O1W—H1WB···O7 ⁱⁱ	0.81 (2)	1.99 (2)	2.790 (2)	175 (3)
O1W—H1WA···O1 ⁱ	0.83 (2)	1.87 (2)	2.653 (2)	155 (2)
O2W—H2WB···O1 ⁱⁱⁱ	0.83 (2)	1.84 (2)	2.661 (2)	173 (3)
O2W—H2WA···O5	0.82 (2)	2.23 (2)	2.958 (3)	148 (3)
N1—H6···O7 ^{iv}	0.94 (3)	1.88 (3)	2.814 (3)	179 (3)

Symmetry codes: (i) $-x, y, -z+1/2$; (ii) $x-1/2, y-1/2, z$; (iii) $x-1/2, -y+1/2, z-1/2$; (iv) $-x+1/2, y+1/2, -z+1/2$.