

8-(2,2,2-Trifluoroethoxy)quinolinium perchlorate–8-(2,2,2-trifluoroethoxy)-quinoline (1/1)

Jun Wu, Lu-Sheng Chen, Qi-Kui Liu and Dian-Shun Guo*

Department of Chemistry, Shandong Normal University, Jinan 250014, People's Republic of China
Correspondence e-mail: chdsguo@sdu.edu.cn


Received 31 March 2011; accepted 8 April 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; disorder in main residue; R factor = 0.050; wR factor = 0.137; data-to-parameter ratio = 12.1.

The title compound, $\text{C}_{11}\text{H}_9\text{F}_3\text{NO}^+\cdot\text{ClO}_4^-\cdot\text{C}_{11}\text{H}_8\text{F}_3\text{NO}$ or $[(\text{C}_{11}\text{H}_8\text{F}_3\text{NO})\text{H}(\text{C}_{11}\text{H}_8\text{F}_3\text{NO})]\text{ClO}_4$, contains two 8-(2,2,2-trifluoroethoxy)quinoline molecules, one of which combines a proton from perchloric acid to form the corresponding quinolinium cation. The quinolinium and quinoline rings form a cationic unit *via* an intermolecular $\text{N}-\text{H}\cdots\text{N}$ hydrogen bond. The heterocyclic units are almost perpendicular to each other [interplanar angle $86.97(6)\text{ }^\circ$]. In the crystal, each perchlorate anion bridges two adjacent cationic units and creates a chain by a combination of $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds. Two inversion-related chains associate into a molecular column by $\pi-\pi$ stacking interactions between the quinolinium rings. The perpendicular and centroid–centroid distances between adjacent quinolinium rings are $3.501(3)$ and $3.634(9)\text{ \AA}$, respectively. The molecular column is linked to its neighbors, creating a two-dimensional network *via* the weak $\pi-\pi$ stacking between the quinoline rings [perpendicular and centroid–centroid separations $3.340(4)$ and $4.408(4)\text{ \AA}$, respectively]. Finally, a three-dimensional framework is formed by a combination of intermolecular $\text{C}-\text{F}\cdots\pi$ contacts. One $-\text{CF}_3$ group is disordered over two positions of equal occupancy.

Related literature

For background to quinoline derivatives, see: Moret *et al.* (2006); Kalita *et al.* (2009). For related structures, see: Ouyang & Khoo *et al.* (1998); Karmakar *et al.* (2009); Al-Mandhary & Steel (2003); Zhang *et al.* (2006); Zheng *et al.* (2006). For $\pi-\pi$ stacking, see: Kalita & Baruah (2010); Chen *et al.* (2005); Liang *et al.* (2002). For $\text{C}-\text{F}\cdots\pi$ contacts, see: Prasanna & Row (2000); Saraogi *et al.* (2003); Choudhury & Row (2004).



Experimental

Crystal data

$\text{C}_{11}\text{H}_9\text{F}_3\text{NO}^+\cdot\text{ClO}_4^-\cdot\text{C}_{11}\text{H}_8\text{F}_3\text{NO}$	$\gamma = 74.536(3)\text{ }^\circ$
$M_r = 554.83$	$V = 1178.0(5)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 9.462(2)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 11.229(3)\text{ \AA}$	$\mu = 0.25\text{ mm}^{-1}$
$c = 11.832(3)\text{ \AA}$	$T = 293\text{ K}$
$\alpha = 82.910(3)\text{ }^\circ$	$0.50 \times 0.32 \times 0.25\text{ mm}$
$\beta = 77.048(3)\text{ }^\circ$	

Data collection

Bruker SMART CCD area-detector diffractometer	6301 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 1999)	4364 independent reflections
$T_{\min} = 0.884$, $T_{\max} = 0.940$	3400 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$	361 parameters
$wR(F^2) = 0.137$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\text{max}} = 0.25\text{ e \AA}^{-3}$
4364 reflections	$\Delta\rho_{\text{min}} = -0.27\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$, $Cg2$ and $Cg3$ are the midpoints of the N1–C5, C5–C9 and C17–C18 bonds, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1–H1A…N2	0.86	1.87	2.684(3)	158
C22–F5… $Cg1$	1.33	3.10	3.796(3)	111
C22–F6… $Cg2$	1.33	3.17	3.804(4)	109
C13–H13…O3 ⁱ	0.93	2.60	3.393(6)	144
C21–H21B…O4 ⁱⁱ	0.97	2.48	3.437(7)	169
C22–F5… $Cg3$ ⁱⁱⁱ	1.33	3.24	3.860(8)	108

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 1, -y + 1, -z + 2$; (iii) $-x + 2, -y + 1, -z + 2$.

Data collection: *SMART* (Bruker, 1999); cell refinement: *SAINT* (Bruker, 1999); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Financial support from the National Natural Science Foundation of China (grant No. 20572064) and the Natural Science Foundation of Shandong Province (grant No. ZR2010BM022) is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2278).

References

- Al-Mandhary, M. R. A. & Steel, P. J. (2003). *Inorg. Chim. Acta*, **351**, 7–11.
- Bruker (1999). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, C.-L., Zhang, J.-A., Li, X.-P., Chen, Z.-N., Kang, B.-S. & Su, C.-Y. (2005). *Inorg. Chim. Acta*, **358**, 4527–4533.
- Choudhury, A. R. & Row, T. N. G. (2004). *Cryst. Growth Des.* **4**, 47–52.
- Kalita, D. & Baruah, J. B. (2010). *CrystEngComm*, **12**, 1562–1567.
- Kalita, D., Sarma, R. & Baruah, J. B. (2009). *CrystEngComm*, **11**, 803–810.
- Karmakar, A., Kalita, D. & Baruah, J. B. (2009). *J. Mol. Struct.* **935**, 47–52.
- Liang, F., Xie, Z., Wang, L., Jing, X. & Wang, F. (2002). *Tetrahedron Lett.* **43**, 3427–3430.
- Moret, V., Dereudre-Bosquet, N., Clayette, P., Laras, Y., Pietrancosta, N., Rolland, A., Weck, C., Marc, S. & Kraus, J.-L. (2006). *Bioorg. Med. Chem. Lett.* **16**, 5988–5992.
- Ouyang, J., Xu, Y. & Khoo, L. E. (1998). *J. Organomet. Chem.* **561**, 143–152.
- Prasanna, M. D. & Row, T. N. G. (2000). *Cryst. Eng.* **3**, 135–154.
- Saraogi, I., Vijay, V. G., Das, S., Sekar, K. & Row, T. N. G. (2003). *Cryst. Eng.* **6**, 69–77.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Zhang, S.-S., Xu, L.-L., Wen, H.-L., Li, X.-M. & Wen, Y.-H. (2006). *Acta Cryst. E* **62**, o3071–o3072.
- Zheng, Z.-B., Li, J.-K., Wu, R.-T. & Sun, Y.-F. (2006). *Acta Cryst. E* **62**, o4611–o4612.

supporting information

Acta Cryst. (2011). E67, o1131–o1132 [doi:10.1107/S1600536811013250]

8-(2,2,2-Trifluoroethoxy)quinolinium perchlorate–8-(2,2,2-trifluoroethoxy)-quinoline (1/1)

Jun Wu, Lu-Sheng Chen, Qi-Kui Liu and Dian-Shun Guo

S1. Comment

Molecules containing quinoline moieties have attracted much interest due to their significant bioactivities (Moret *et al.*, 2006) and potential applications for constructing supramolecular systems by various hydrogen bonds and π – π interactions (Kalita *et al.*, 2009). Numerous crystal structures of 8-substituted quinolines, most of which are 8-alkyloxyquinoline derivatives have been reported (Karmakar *et al.*, 2009; Al-Mandhary *et al.*, 2003; Zhang *et al.*, 2006; Zheng *et al.*, 2006), while the crystal structures of such compounds exhibiting fluorinated alkyl groups have rarely been described. We synthesized a new trifluoroethoxyquinoline derivative and attempted to prepare its Ni(II)-complex. Surprisingly, a complex without the Ni(II) ion, namely 8-(2,2,2-trifluoroethoxy)-quinolinium-[8-(2,2,2-trifluoroethoxy)-quinoline] perchlorate, was obtained. Here, we report its crystal structure.

In the crystal structure of the title complex the asymmetric unit contains one perchlorate anion and one organic cation consisting of two 8-(2,2,2-trifluoroethoxy)-quinolines and a proton (Fig. 1). In the cationic unit, one quinoline ring is protonated forming a quinolinium and is connected to the other one *via* a N—H \cdots N hydrogen bond (N—H \cdots N angle 155.33 (5) $^\circ$, N1 \cdots N2 2.684 (3) Å, Fig. 2, Table 1). This distance is shorter than 2.729 Å reported for a similar quinoline derivative (Ouyang *et al.*, 1998). Such a short distance may be ascribed to the presence of two weak C—F \cdots π contacts (Prasanna & Row, 2000; Saraogi *et al.*, 2003; Choudhury & Row, 2004) between the non-disordered trifluoromethyl group and the quinoline ring. Separations of F5 \cdots Cg1 and F6 \cdots Cg2 (Cg1 and Cg2 are the centroids of N1—C5 and C5—C9 bonds, respectively) are 3.099 (3) and 3.166 (3) Å, respectively. The quinolinium and quinoline rings are almost perpendicular to each other, with a dihedral angle of 86.97 (6) $^\circ$. The other trifluoromethyl group is disordered over two orientations, with refined site-occupancy factors of 0.5.

In the packing of the title complex, there exist intermolecular C—H \cdots O hydrogen bonds, C—F \cdots π contacts (Table 1) and π – π stackings (Kalita & Baruah, 2010; Chen *et al.*, 2005; Liang *et al.*, 2002). The cationic units are alternately bridged by perchlorate anions with intermolecular C—H \cdots O hydrogen bonds and form an infinite one-dimensional chain along the *c* axis (Fig. 3). Hydrogen bonds arise from atoms C13—H13 and C21—H21B in the molecule at (*x*, *y*, *z*) acting as hydrogen bond donors towards atoms O3 at ($-x + 1$, $-y + 1$, $-z + 1$) and O4 ($-x + 1$, $-y + 1$, $-z + 2$), respectively. Two adjacent chains running along the alternate orientation are further combined to a molecular column by π – π stacking interactions between the quinolinium rings, with perpendicular and centroid-centroid distances of 3.501 (3) and 3.634 (9) Å, respectively, between neighboring phenyl rings of the quinolinium units. Finally, a complicated three-dimensional framework is formed by a combination of C—F \cdots π contacts, with a F5 \cdots Cg3 (Cg3 is the centroid of C17—C18 bond) distance of 3.240 (7) Å (Table 1), and weak π – π stackings between adjacent phenyl rings of the quinoline moieties, with perpendicular and centroid-centroid separations of 3.340 (4) and 4.408 (4) Å, respectively.


S2. Experimental

A suspension of 8-hydroxyquinoline (0.200 g, 1.378 mmol), anhydrous KOH (0.093 g, 1.657 mmol) and 2,2,2-trifluoroethyl-4-methylbenzenesulfonate (0.385 g, 1.514 mmol) in dry NMP (5 ml) was stirred for 5 h at 393 K and then cooled to room temperature. The resulting mixture was neutralized with 5% aqueous HCl and extracted with CH₂Cl₂. The organic layer was separated and washed with saturated sodium hydrogen carbonate and brine, and dried over anhydrous MgSO₄. Removal of the solvent under reduced pressure gave 8-(2,2,2-trifluoroethoxy)-quinoline as a yellow solid (yield 73%), which was purified by flash column chromatography (EtOAc/petroleum ether = 1:3, *R*_F = 1/2).


Single crystals of the title complex suitable for X-ray analysis were obtained from a solution of Ni(ClO₄)₂ (0.013 g, 0.050 mmol) in MeOH (1 ml) which was layered onto a solution of 8-(2,2,2-trifluoroethoxy)-quinoline (0.012 g, 0.053 mmol) in CH₂Cl₂ (1 ml) at 298 K.

S3. Refinement


All non-hydrogen atoms were refined with anisotropic displacement parameters. Hydrogen atoms attached to anisotropically refined atoms were placed in geometrically idealized positions and included as riding atoms with C—H = 0.93 Å and *U*_{iso}(H) = 1.2*U*_{eq}(C) (aromatic); C—H = 0.97 Å and *U*_{iso}(H) = 1.2*U*_{eq}(C) (methylene); N—H = 0.86 Å and *U*_{iso}(H) = 1.2*U*_{eq}(N). In the title molecule, one trifluoromethyl group was disordered over two orientations, with refined site occupation factors of 0.5: 0.5.

**Figure 1**

Molecular structure of the title complex, with displacement ellipsoids drawn at the 30% probability level.

**Figure 2**

Hydrogen-bonded cationic unit, with weak C—F··· π contacts. For the sake of clarity, the disordered moieties and the H atoms not involved in hydrogen bonding have been omitted.

**Figure 3**

Molecular column formed by hydrogen bonds and π — π stackings, viewed along the a axis. The disordered moieties and H atoms not involved in hydrogen bonding have been omitted for clarity. [Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 1, -y + 1, -z + 2$].

8-(2,2,2-Trifluoroethoxy)quinolinium perchlorate–8-(2,2,2-trifluoroethoxy)quinoline (1/1)

Crystal data



$M_r = 554.83$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.462 (2)$ Å

$b = 11.229 (3)$ Å

$c = 11.832 (3)$ Å

$\alpha = 82.910 (3)^\circ$

$\beta = 77.048 (3)^\circ$

$\gamma = 74.536 (3)^\circ$

$V = 1178.0 (5)$ Å³

$Z = 2$

$F(000) = 564$

$D_x = 1.564$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 2404 reflections
 $\theta = 2.5\text{--}25.1^\circ$
 $\mu = 0.25 \text{ mm}^{-1}$

$T = 293 \text{ K}$
 Block, colourless
 $0.50 \times 0.32 \times 0.25 \text{ mm}$

Data collection

Bruker SMART CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ϕ and ω scans
 Absorption correction: multi-scan
 (*SADABS*; Bruker, 1999)
 $T_{\min} = 0.884$, $T_{\max} = 0.940$

6301 measured reflections
 4364 independent reflections
 3400 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$
 $\theta_{\max} = 25.6^\circ$, $\theta_{\min} = 1.8^\circ$
 $h = -11 \rightarrow 8$
 $k = -13 \rightarrow 9$
 $l = -14 \rightarrow 14$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.137$
 $S = 1.02$
 4364 reflections
 361 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.068P)^2 + 0.293P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.25 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.27 \text{ e \AA}^{-3}$

Special details

Experimental. ^1H NMR (300 MHz, CDCl \sim 3): δ 8.97 (dd, 1H, $J = 4.04$ Hz, 1.52 Hz), 8.17 (d, 1H, $J = 7.52$ Hz), 7.54–7.44 (m, 3H), 7.25 (d, H, $J = 7.65$ Hz), 4.78 (q, 2H, $J = 8.33$ Hz).

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	1.1468 (3)	0.6885 (3)	0.6473 (2)	0.0583 (6)	
H1	1.1569	0.6034	0.6558	0.070*	
C2	1.2727 (3)	0.7332 (3)	0.6049 (2)	0.0694 (8)	
H2	1.3662	0.6791	0.5845	0.083*	
C3	1.2575 (3)	0.8571 (3)	0.5937 (2)	0.0682 (8)	
H3	1.3416	0.8880	0.5653	0.082*	
C4	1.1164 (3)	0.9399 (2)	0.6243 (2)	0.0548 (6)	
C5	0.9930 (3)	0.8878 (2)	0.66440 (18)	0.0448 (5)	
C6	1.0927 (4)	1.0696 (3)	0.6165 (2)	0.0677 (8)	
H6	1.1731	1.1056	0.5909	0.081*	
C7	0.9530 (4)	1.1412 (3)	0.6463 (2)	0.0738 (8)	

H7	0.9385	1.2268	0.6415	0.089*
C8	0.8288 (3)	1.0905 (2)	0.6843 (2)	0.0643 (7)
H8	0.7334	1.1424	0.7027	0.077*
C9	0.8474 (3)	0.9661 (2)	0.6943 (2)	0.0505 (6)
C10	0.5911 (3)	0.9754 (3)	0.7691 (3)	0.0844 (10)
H10A	0.5518	1.0228	0.7037	0.101*
H10B	0.5902	1.0325	0.8245	0.101*
C12	0.7912 (3)	0.5853 (2)	0.6481 (2)	0.0637 (7)
H12	0.8219	0.6259	0.5770	0.076*
C13	0.7078 (4)	0.4983 (3)	0.6515 (3)	0.0731 (8)
H13	0.6806	0.4839	0.5845	0.088*
C14	0.6678 (3)	0.4362 (2)	0.7528 (3)	0.0687 (8)
H14	0.6140	0.3772	0.7557	0.082*
C15	0.7062 (3)	0.4590 (2)	0.8546 (2)	0.0533 (6)
C16	0.7845 (2)	0.55191 (19)	0.8454 (2)	0.0449 (5)
C17	0.6663 (3)	0.3978 (2)	0.9643 (3)	0.0664 (7)
H17	0.6164	0.3355	0.9710	0.080*
C18	0.7003 (3)	0.4296 (2)	1.0592 (3)	0.0670 (7)
H18	0.6739	0.3884	1.1308	0.080*
C19	0.7750 (3)	0.5238 (2)	1.0520 (2)	0.0574 (6)
H19	0.7962	0.5450	1.1188	0.069*
C20	0.8163 (2)	0.5841 (2)	0.9478 (2)	0.0465 (5)
C21	0.9028 (3)	0.7245 (2)	1.0323 (2)	0.0550 (6)
H21A	0.9774	0.6653	1.0687	0.066*
H21B	0.8083	0.7397	1.0876	0.066*
C22	0.9496 (3)	0.8415 (3)	0.9960 (2)	0.0621 (7)
C11	0.35067 (7)	0.30823 (6)	0.64594 (5)	0.0564 (2)
F4	0.9743 (3)	0.88475 (17)	1.08657 (15)	0.0965 (6)
F5	1.07304 (19)	0.82890 (16)	0.91311 (15)	0.0783 (5)
F6	0.8460 (2)	0.92888 (15)	0.95375 (16)	0.0850 (5)
N1	1.0134 (2)	0.76298 (17)	0.67574 (16)	0.0463 (4)
H1A	0.9374	0.7321	0.7021	0.056*
N2	0.8280 (2)	0.61195 (17)	0.74111 (17)	0.0482 (5)
O1	0.73864 (18)	0.90405 (15)	0.73088 (16)	0.0613 (5)
O2	0.88758 (18)	0.67771 (15)	0.93048 (13)	0.0526 (4)
O3	0.4147 (3)	0.4098 (2)	0.6013 (2)	0.0932 (7)
O4	0.4074 (2)	0.25456 (19)	0.74758 (18)	0.0835 (6)
O5	0.3897 (3)	0.2197 (2)	0.5613 (2)	0.0979 (8)
O6	0.1932 (2)	0.3522 (2)	0.6752 (2)	0.0888 (7)
C11	0.4994 (4)	0.8884 (4)	0.8241 (4)	0.0902 (10) 0.50
F1	0.5026 (16)	0.8110 (15)	0.7456 (17)	0.177 (7) 0.50
F2	0.3618 (17)	0.9521 (18)	0.8772 (11)	0.131 (4) 0.50
F3	0.5473 (15)	0.8205 (11)	0.9134 (14)	0.118 (5) 0.50
C11W	0.4994 (4)	0.8884 (4)	0.8241 (4)	0.0902 (10) 0.50
F1W	0.5117 (10)	0.7933 (8)	0.7704 (12)	0.105 (4) 0.50
F2W	0.3555 (16)	0.9444 (18)	0.8260 (11)	0.109 (3) 0.50
F3W	0.5318 (19)	0.8454 (16)	0.9246 (13)	0.142 (6) 0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0535 (15)	0.0587 (15)	0.0562 (15)	-0.0040 (12)	-0.0086 (12)	-0.0055 (12)
C2	0.0472 (15)	0.085 (2)	0.0672 (18)	-0.0088 (14)	-0.0025 (13)	-0.0029 (15)
C3	0.0540 (16)	0.094 (2)	0.0623 (17)	-0.0336 (15)	-0.0074 (13)	-0.0012 (15)
C4	0.0612 (16)	0.0688 (16)	0.0412 (13)	-0.0279 (13)	-0.0108 (11)	-0.0022 (11)
C5	0.0539 (13)	0.0484 (13)	0.0347 (11)	-0.0179 (11)	-0.0090 (10)	-0.0007 (9)
C6	0.087 (2)	0.0718 (18)	0.0563 (16)	-0.0452 (17)	-0.0110 (15)	0.0005 (13)
C7	0.116 (3)	0.0494 (15)	0.0600 (17)	-0.0323 (17)	-0.0124 (17)	-0.0016 (13)
C8	0.0779 (18)	0.0474 (14)	0.0599 (16)	-0.0087 (13)	-0.0078 (14)	-0.0004 (12)
C9	0.0564 (14)	0.0476 (13)	0.0445 (13)	-0.0114 (11)	-0.0081 (11)	0.0008 (10)
C10	0.0535 (17)	0.0661 (18)	0.111 (3)	0.0037 (14)	0.0015 (16)	0.0071 (17)
C12	0.0788 (19)	0.0615 (16)	0.0556 (16)	-0.0193 (14)	-0.0209 (14)	-0.0046 (12)
C13	0.085 (2)	0.0695 (18)	0.080 (2)	-0.0252 (16)	-0.0348 (17)	-0.0145 (16)
C14	0.0658 (18)	0.0522 (15)	0.099 (2)	-0.0214 (13)	-0.0283 (16)	-0.0109 (15)
C15	0.0451 (13)	0.0400 (12)	0.0764 (17)	-0.0112 (10)	-0.0144 (12)	-0.0041 (11)
C16	0.0404 (12)	0.0361 (11)	0.0566 (14)	-0.0069 (9)	-0.0101 (10)	-0.0027 (10)
C17	0.0574 (16)	0.0501 (15)	0.093 (2)	-0.0240 (12)	-0.0120 (15)	0.0066 (14)
C18	0.0654 (17)	0.0598 (16)	0.0700 (18)	-0.0220 (14)	-0.0037 (14)	0.0135 (13)
C19	0.0573 (15)	0.0566 (15)	0.0567 (15)	-0.0169 (12)	-0.0069 (12)	0.0003 (12)
C20	0.0437 (12)	0.0412 (12)	0.0534 (14)	-0.0109 (10)	-0.0062 (10)	-0.0035 (10)
C21	0.0618 (15)	0.0595 (15)	0.0467 (13)	-0.0199 (12)	-0.0097 (11)	-0.0066 (11)
C22	0.0790 (19)	0.0619 (16)	0.0541 (15)	-0.0259 (15)	-0.0184 (14)	-0.0094 (12)
C11	0.0547 (4)	0.0556 (4)	0.0571 (4)	-0.0178 (3)	-0.0036 (3)	-0.0017 (3)
F4	0.1570 (19)	0.0919 (12)	0.0696 (11)	-0.0634 (13)	-0.0396 (12)	-0.0119 (9)
F5	0.0817 (11)	0.0814 (11)	0.0788 (11)	-0.0408 (9)	-0.0069 (9)	-0.0032 (9)
F6	0.1038 (13)	0.0591 (10)	0.0926 (13)	-0.0115 (9)	-0.0315 (11)	-0.0039 (9)
N1	0.0444 (11)	0.0473 (11)	0.0459 (11)	-0.0131 (9)	-0.0054 (8)	-0.0011 (8)
N2	0.0519 (11)	0.0442 (10)	0.0500 (11)	-0.0117 (9)	-0.0124 (9)	-0.0051 (8)
O1	0.0449 (9)	0.0506 (10)	0.0792 (12)	-0.0075 (8)	-0.0004 (8)	-0.0011 (9)
O2	0.0648 (10)	0.0551 (9)	0.0453 (9)	-0.0284 (8)	-0.0091 (8)	-0.0050 (7)
O3	0.1065 (17)	0.0857 (15)	0.1007 (17)	-0.0546 (14)	-0.0250 (13)	0.0195 (12)
O4	0.0875 (15)	0.0769 (13)	0.0738 (13)	-0.0045 (11)	-0.0184 (11)	0.0110 (11)
O5	0.0924 (16)	0.1109 (18)	0.0948 (16)	-0.0381 (14)	0.0101 (13)	-0.0478 (14)
O6	0.0554 (12)	0.0968 (16)	0.1055 (17)	-0.0104 (11)	-0.0081 (11)	-0.0077 (13)
C11	0.0475 (18)	0.089 (3)	0.120 (4)	-0.0082 (17)	0.0047 (19)	-0.010 (3)
F1	0.109 (7)	0.217 (14)	0.235 (12)	-0.025 (7)	-0.069 (7)	-0.097 (10)
F2	0.054 (5)	0.124 (6)	0.179 (12)	0.000 (4)	0.034 (7)	-0.030 (9)
F3	0.084 (5)	0.082 (3)	0.155 (12)	-0.026 (3)	0.021 (5)	0.038 (5)
C11W	0.0475 (18)	0.089 (3)	0.120 (4)	-0.0082 (17)	0.0047 (19)	-0.010 (3)
F1W	0.056 (4)	0.077 (4)	0.181 (9)	-0.031 (3)	0.001 (4)	-0.023 (5)
F2W	0.045 (3)	0.128 (5)	0.143 (8)	-0.010 (3)	0.007 (5)	-0.041 (7)
F3W	0.142 (11)	0.194 (13)	0.093 (7)	-0.070 (9)	-0.007 (6)	0.002 (7)

Geometric parameters (\AA , $\text{^{\circ}}$)

C1—N1	1.312 (3)	C14—H14	0.9300
C1—C2	1.383 (4)	C15—C16	1.413 (3)
C1—H1	0.9300	C15—C17	1.414 (4)
C2—C3	1.352 (4)	C16—N2	1.364 (3)
C2—H2	0.9300	C16—C20	1.421 (3)
C3—C4	1.408 (4)	C17—C18	1.346 (4)
C3—H3	0.9300	C17—H17	0.9300
C4—C5	1.406 (3)	C18—C19	1.404 (4)
C4—C6	1.408 (4)	C18—H18	0.9300
C5—N1	1.357 (3)	C19—C20	1.361 (3)
C5—C9	1.418 (3)	C19—H19	0.9300
C6—C7	1.346 (4)	C20—O2	1.365 (3)
C6—H6	0.9300	C21—O2	1.422 (3)
C7—C8	1.401 (4)	C21—C22	1.480 (4)
C7—H7	0.9300	C21—H21A	0.9700
C8—C9	1.354 (3)	C21—H21B	0.9700
C8—H8	0.9300	C22—F4	1.319 (3)
C9—O1	1.357 (3)	C22—F6	1.326 (3)
C10—O1	1.416 (3)	C22—F5	1.334 (3)
C10—C11	1.474 (5)	C11—O6	1.415 (2)
C10—H10A	0.9700	C11—O5	1.418 (2)
C10—H10B	0.9700	C11—O3	1.423 (2)
C12—N2	1.316 (3)	C11—O4	1.427 (2)
C12—C13	1.402 (4)	N1—H1A	0.8600
C12—H12	0.9300	C11—F3	1.317 (15)
C13—C14	1.341 (4)	C11—F1	1.338 (17)
C13—H13	0.9300	C11—F2	1.359 (15)
C14—C15	1.403 (4)		
N1—C1—C2	121.7 (3)	N2—C16—C15	121.5 (2)
N1—C1—H1	119.1	N2—C16—C20	119.7 (2)
C2—C1—H1	119.1	C15—C16—C20	118.8 (2)
C3—C2—C1	118.9 (3)	C18—C17—C15	120.2 (2)
C3—C2—H2	120.6	C18—C17—H17	119.9
C1—C2—H2	120.6	C15—C17—H17	119.9
C2—C3—C4	121.0 (3)	C17—C18—C19	121.3 (3)
C2—C3—H3	119.5	C17—C18—H18	119.4
C4—C3—H3	119.5	C19—C18—H18	119.4
C5—C4—C6	119.0 (3)	C20—C19—C18	120.3 (3)
C5—C4—C3	116.9 (2)	C20—C19—H19	119.8
C6—C4—C3	124.1 (3)	C18—C19—H19	119.8
N1—C5—C4	120.2 (2)	C19—C20—O2	125.5 (2)
N1—C5—C9	120.1 (2)	C19—C20—C16	120.1 (2)
C4—C5—C9	119.8 (2)	O2—C20—C16	114.40 (19)
C7—C6—C4	119.7 (3)	O2—C21—C22	107.24 (19)
C7—C6—H6	120.2	O2—C21—H21A	110.3

C4—C6—H6	120.2	C22—C21—H21A	110.3
C6—C7—C8	121.9 (3)	O2—C21—H21B	110.3
C6—C7—H7	119.0	C22—C21—H21B	110.3
C8—C7—H7	119.0	H21A—C21—H21B	108.5
C9—C8—C7	120.1 (3)	F4—C22—F6	107.2 (2)
C9—C8—H8	119.9	F4—C22—F5	108.3 (2)
C7—C8—H8	119.9	F6—C22—F5	105.6 (2)
C8—C9—O1	126.7 (2)	F4—C22—C21	109.5 (2)
C8—C9—C5	119.5 (2)	F6—C22—C21	112.6 (2)
O1—C9—C5	113.7 (2)	F5—C22—C21	113.3 (2)
O1—C10—C11	107.3 (2)	O6—C11—O5	109.93 (14)
O1—C10—H10A	110.3	O6—C11—O3	109.00 (14)
C11—C10—H10A	110.3	O5—C11—O3	109.86 (15)
O1—C10—H10B	110.3	O6—C11—O4	109.81 (14)
C11—C10—H10B	110.3	O5—C11—O4	110.33 (15)
H10A—C10—H10B	108.5	O3—C11—O4	107.87 (14)
N2—C12—C13	122.8 (3)	C1—N1—C5	121.2 (2)
N2—C12—H12	118.6	C1—N1—H1A	119.4
C13—C12—H12	118.6	C5—N1—H1A	119.4
C14—C13—C12	118.9 (3)	C12—N2—C16	118.9 (2)
C14—C13—H13	120.5	C9—O1—C10	117.2 (2)
C12—C13—H13	120.5	C20—O2—C21	116.17 (18)
C13—C14—C15	120.8 (3)	F3—C11—F1	107.5 (10)
C13—C14—H14	119.6	F3—C11—F2	100.6 (10)
C15—C14—H14	119.6	F1—C11—F2	116.5 (9)
C14—C15—C16	117.0 (2)	F3—C11—C10	113.4 (7)
C14—C15—C17	123.7 (2)	F1—C11—C10	108.9 (8)
C16—C15—C17	119.3 (2)	F2—C11—C10	109.9 (9)
N1—C1—C2—C3	0.6 (4)	C16—C15—C17—C18	1.2 (4)
C1—C2—C3—C4	0.0 (4)	C15—C17—C18—C19	0.3 (4)
C2—C3—C4—C5	-1.3 (4)	C17—C18—C19—C20	-0.8 (4)
C2—C3—C4—C6	179.0 (3)	C18—C19—C20—O2	179.2 (2)
C6—C4—C5—N1	-178.2 (2)	C18—C19—C20—C16	-0.2 (4)
C3—C4—C5—N1	2.1 (3)	N2—C16—C20—C19	-179.2 (2)
C6—C4—C5—C9	1.1 (3)	C15—C16—C20—C19	1.7 (3)
C3—C4—C5—C9	-178.7 (2)	N2—C16—C20—O2	1.3 (3)
C5—C4—C6—C7	-0.7 (4)	C15—C16—C20—O2	-177.81 (19)
C3—C4—C6—C7	179.0 (3)	O2—C21—C22—F4	-176.1 (2)
C4—C6—C7—C8	-0.6 (4)	O2—C21—C22—F6	64.7 (3)
C6—C7—C8—C9	1.4 (4)	O2—C21—C22—F5	-55.0 (3)
C7—C8—C9—O1	179.3 (2)	C2—C1—N1—C5	0.2 (4)
C7—C8—C9—C5	-0.9 (4)	C4—C5—N1—C1	-1.6 (3)
N1—C5—C9—C8	179.0 (2)	C9—C5—N1—C1	179.1 (2)
C4—C5—C9—C8	-0.3 (3)	C13—C12—N2—C16	0.6 (4)
N1—C5—C9—O1	-1.2 (3)	C15—C16—N2—C12	2.4 (3)
C4—C5—C9—O1	179.6 (2)	C20—C16—N2—C12	-176.7 (2)
N2—C12—C13—C14	-2.4 (4)	C8—C9—O1—C10	-3.9 (4)

C12—C13—C14—C15	1.1 (4)	C5—C9—O1—C10	176.3 (2)
C13—C14—C15—C16	1.6 (4)	C11—C10—O1—C9	−169.7 (3)
C13—C14—C15—C17	179.3 (3)	C19—C20—O2—C21	−7.9 (3)
C14—C15—C16—N2	−3.5 (3)	C16—C20—O2—C21	171.53 (19)
C17—C15—C16—N2	178.8 (2)	C22—C21—O2—C20	−167.1 (2)
C14—C15—C16—C20	175.6 (2)	O1—C10—C11—F3	59.6 (8)
C17—C15—C16—C20	−2.2 (3)	O1—C10—C11—F1	−60.0 (9)
C14—C15—C17—C18	−176.4 (3)	O1—C10—C11—F2	171.3 (6)

Hydrogen-bond geometry (Å, °)

Cg1, Cg2 and Cg3 are the midpoints of the N1—C5, C5—C9 and C17—C18 bonds, respectively.

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1A···N2	0.86	1.87	2.684 (3)	158
C22—F5···Cg1	1.33	3.10	3.796 (3)	111
C22—F6···Cg2	1.33	3.17	3.804 (4)	109
C13—H13···O3 ⁱ	0.93	2.60	3.393 (6)	144
C21—H21B···O4 ⁱⁱ	0.97	2.48	3.437 (7)	169
C22—F5···Cg3 ⁱⁱⁱ	1.33	3.24	3.860 (8)	108

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+1, -y+1, -z+2$; (iii) $-x+2, -y+1, -z+2$.