

(E)-1-(2,4-Dinitrophenyl)-2-(2-fluoro-benzylidene)hydrazine

Jerry P. Jasinski,^{a*} Adam N. Braley,^a C. S. Chidan Kumar,^b H. S. Yathirajan^b and A. N. Mayekar^c

^aDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, ^bDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, and ^cSeQuent Scientific Ltd, Baikampady, New Mangalore 575 011, India

Correspondence e-mail: jjasinski@keene.edu


Received 13 April 2011; accepted 16 April 2011

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.045; wR factor = 0.143; data-to-parameter ratio = 17.4.

In the title compound, $\text{C}_{13}\text{H}_9\text{FN}_4\text{O}_4$, the dihedral angle between the mean planes of the two benzene rings of the nearly planar molecule is $6.6(9)^\circ$. The dihedral angles between the mean planes of the benzene ring and its two attached nitro groups are $6.7(7)$ and $7.2(9)^\circ$. Crystal packing is stabilized by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, weak $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{F}$ intermolecular interactions and centroid–centroid π -ring stacking interactions.

Related literature

For Schiff base properties, see: Liang (2007). For nonlinear optical and crystalline properties, see: Baughman *et al.* (2004). For DNA-damaging and mutagenic agents, see: Okabe *et al.* (1993). For related structures, see: Bolte & Dill (1998); Shan *et al.* (2002); Fan *et al.* (2004); Motherwell & Ramsay, (2007); Shi *et al.* (2008); Ji *et al.* (2010); Kia *et al.* (2009); Jasinski *et al.* (2010).



Experimental

Crystal data



$M_r = 304.24$

Triclinic, $P\bar{1}$

$a = 7.0961(8)\text{ \AA}$

$b = 8.2714(9)\text{ \AA}$

$c = 11.7230(8)\text{ \AA}$

$\alpha = 88.614(7)^\circ$

$\beta = 80.544(8)^\circ$

$\gamma = 71.368(10)^\circ$

$V = 642.86(11)\text{ \AA}^3$

$Z = 2$

$\text{Mo K}\alpha$ radiation

$\mu = 0.13\text{ mm}^{-1}$

$T = 173\text{ K}$

$0.20 \times 0.18 \times 0.15\text{ mm}$

Data collection

Oxford Diffraction Xcalibur Eos
Gemini diffractometer
Absorption correction: multi-scan
(*CrysAlis RED*; Oxford
Diffraction, 2007)
 $T_{\min} = 0.967$, $T_{\max} = 1.000$

6346 measured reflections
3466 independent reflections
2802 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.143$
 $S = 1.09$
3466 reflections

199 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.31\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.17\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2A \cdots O1	0.88	2.02	2.6317 (15)	126
N2—H2A \cdots O1 ⁱ	0.88	2.51	3.3424 (15)	158
C2—H2B \cdots F1 ⁱⁱ	0.95	2.45	3.3386 (17)	156
C3—H3A \cdots O4 ⁱⁱⁱ	0.95	2.48	3.3177 (19)	148
C5—H5A \cdots O3 ^{iv}	0.95	2.43	3.2694 (17)	148

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 3, -y, -z + 1$; (iii) $x + 2, y - 1, z$; (iv) $-x + 1, -y, -z$.

Table 2
 $Cg\cdots Cg$ π -ring stacking interactions.

$Cg1$ and $Cg2$ are the centroids of rings C1–C6 and C8–C13, respectively.

$CgI\cdots CgJ$	$Cg\cdots Cg$ (\AA)	$Cg I_{\text{Perp}}$ (\AA)	$Cg J_{\text{Perp}}$ (\AA)
$Cg1\cdots Cg2^i$	3.6916 (10)	-3.4632 (6)	3.3267 (5)
$Cg2\cdots Cg1^{ii}$	3.6916 (10)	3.3267 (5)	-3.4632 (6)

Symmetry codes: (i) $1 + x, y, z$; (ii) $-1 + x, y, z$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2007); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

CSCK and HSY thank the University of Mysore for research facilities. JPJ acknowledges the NSF–MRI program (grant No. CHE1039027) for funds to purchase the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5151).

References

- Baughman, R. G., Martin, K. L., Singh, R. K. & Stoffer, J. O. (2004). *Acta Cryst. C* **60**, o103–o106.
- Bolte, M. & Dill, M. (1998). *Acta Cryst. C* **54**, IUC9800065.
- Fan, Z., Shan, S., Hu, W.-X. & Xu, D.-J. (2004). *Acta Cryst. E* **60**, o1102–o1104.
- Jasinski, J. P., Guild, C. J., Chidan Kumar, C. S., Yathirajan, H. S. & Mayekar, A. N. (2010). *Acta Cryst. E* **66**, o2832–o2833.
- Ji, N.-N., Shi, Z.-Q., Zhao, R.-G., Zheng, Z.-B. & Li, Z.-F. (2010). *Bull. Korean Chem. Soc.* **31**, 881–886.
- Kia, R., Fun, H.-K., Etemadi, B. & Kargar, H. (2009). *Acta Cryst. E* **65**, o833–o834.

- Liang, Z.-P. (2007). *Acta Cryst.* E**63**, o2943.
Motherwell, W. D. S. & Ramsay, J. (2007). *Acta Cryst.* E**63**, o4043.
Okabe, N., Nakamura, T. & Fukuda, H. (1993). *Acta Cryst.* C**49**, 1678–1680.
Oxford Diffraction (2007). *CrysAlis PRO* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
Shan, S., Xu, D.-J., Wu, J.-Y. & Chiang, M. Y. (2002). *Acta Cryst.* E**58**, o1333–o1335.
Sheldrick, G. M. (2008). *Acta Cryst.* A**64**, 112–122.
Shi, Z.-Q., Ji, N.-N. & Li, X.-Y. (2008). *Acta Cryst.* E**64**, o2135.

supporting information

Acta Cryst. (2011). E67, o1200–o1201 [doi:10.1107/S1600536811014383]

(E)-1-(2,4-Dinitrophenyl)-2-(2-fluorobenzylidene)hydrazine

Jerry P. Jasinski, Adam N. Braley, C. S. Chidan Kumar, H. S. Yathirajan and A. N. Mayekar

S1. Comment

Schiff bases and their complexes are widely used in the fields of biology, catalysis etc. (Liang, 2007). Especially, the dinitrophenyl hydrazones exhibit good nonlinear optical (NLO) and crystalline properties (Baughman *et al.*, 2004) and are found to have versatile coordinating abilities towards different metal ions. In addition, some 2,4-dinitrophenyl hydrazone derivatives have been shown to be potentially DNA-damaging and mutagenic agents (Okabe *et al.*, 1993). As a result of their significant molecular nonlinearities many x-ray structural studies of 2,4-dinitrophenylhydrazones have been reported. Among them, the most closely related structures are (E)-p-methoxy-acetophenone 2,4-dinitrophenylhydrazone (Bolte & Dill, 1998), acetophenone (2,4-dinitrophenyl)hydrazone (Shan *et al.*, 2002), 3-chloroacetophenone 2,4-dinitrophenyl- hydrazone (Fan *et al.*, 2004), 2,4-dihydroxyacetophenone 2,4-dinitrophenylhydrazone (Baughman *et al.*, 2004), syn-acetophenone (2,4-dinitrophenyl) hydrazone (Motherwell & Ramsay, 2007), 1-(2-chlorobenzylidene)-2-(2,4-dinitrophenyl)hydrazine (Shi *et al.*, 2008), N-(2,4-dinitrophenyl)-N'-(1-p-tolylethylidene) hydrazine (Kia *et al.*, 2009), N-(2,4-dinitrophenyl)-N'-(1-phenylethylidene)hydrazine (Ji *et al.*, 2010) and (1E)-1-(3-bromophenyl)ethanone 2,4-dinitrophenyl- hydrazone (Jasinski *et al.*, 2010). In view of the importance of 2,4-dinitrophenylhydrazones, this paper reports the crystal structure of the title compound, $C_{13}H_9FN_4O_4$, (I).


In the title compound the dihedral angle between the mean planes of the two benzene rings of a nearly planar molecule is 6.69° , (Fig. 2). The dihedral angle between the mean planes of the benzene ring and its two bonded nitro groups are 6.7 (7) $^\circ$ and 7.2 (9) $^\circ$, respectively. Crystal packing is stabilized by N—H \cdots O hydrogen bonds (Fig. 3), weak C—H \cdots O intermolecular interactions and Cg—Cg π -ring stacking interactions (Table 2).

S2. Experimental


A mixture of 2,4-dinitrophenylhydrazine (1.98 g) and 2-fluorobenzaldehyde (1.24 g) was dissolved in methanol and refluxed for about 6h. The precipitate formed was filtered, dried and recrystallized in ethlyacetate. X-ray quality crystals of the title compound (I), were obtained after three days by the slow evaporation of a 1:1 mixture of dimethylformamide and pyridine at room temperature. (mp: 502 - 505 K).

S3. Refinement

The parameters of all the H atoms have been constrained within the riding atom approximation. C—H bond lengths were constrained to 0.95 \AA for aryl atoms, $U_{\text{iso}}(\text{H}) = 1.18\text{--}1.20U_{\text{eq}}(\text{C}_{\text{aryl}})$. N—H bond lengths were constrained to 0.88 \AA , $U_{\text{iso}}(\text{H}) = 1.20U_{\text{eq}}(\text{N})$.

**Figure 1**

Molecular structure of the title compound showing the atom labeling scheme and 50% probability displacement ellipsoids.

**Figure 2**

Packing diagram of the title compound viewed down the b axis. Dashed lines indicate $\text{N}—\text{H} \cdots \text{O}$ hydrogen bonds and weak $\text{N}—\text{H} \cdots \text{O}$ intermolecular interactions.

(E)-1-(2,4-Dinitrophenyl)-2-(2-fluorobenzylidene)hydrazine

Crystal data

$\text{C}_{13}\text{H}_9\text{FN}_4\text{O}_4$
 $M_r = 304.24$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 7.0961 (8)$ Å
 $b = 8.2714 (9)$ Å
 $c = 11.7230 (8)$ Å
 $\alpha = 88.614 (7)^\circ$
 $\beta = 80.544 (8)^\circ$

$\gamma = 71.368 (10)^\circ$
 $V = 642.86 (11)$ Å³
 $Z = 2$
 $F(000) = 312$
 $D_x = 1.572$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 3528 reflections
 $\theta = 3.1\text{--}32.2^\circ$
 $\mu = 0.13$ mm⁻¹

$T = 173\text{ K}$
Block, orange-red

$0.20 \times 0.18 \times 0.15\text{ mm}$

Data collection

Oxford Diffraction Xcalibur Eos Gemini
diffractometer
Radiation source: Enhance (Mo) X-ray Source
Graphite monochromator
Detector resolution: 16.1500 pixels mm^{-1}
 φ and ω scans
Absorption correction: multi-scan
(*CrysAlis RED*; Oxford Diffraction, 2007)
 $T_{\min} = 0.967$, $T_{\max} = 1.000$

6346 measured reflections
3466 independent reflections
2802 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$
 $\theta_{\max} = 29.1^\circ$, $\theta_{\min} = 3.1^\circ$
 $h = -9 \rightarrow 8$
 $k = -11 \rightarrow 11$
 $l = -16 \rightarrow 15$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.143$
 $S = 1.09$
3466 reflections
199 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0733P)^2 + 0.0845P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.31\text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.17\text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	1.25967 (13)	0.08268 (12)	0.46347 (7)	0.0542 (3)
O1	0.33061 (16)	0.49706 (16)	0.46244 (9)	0.0576 (3)
O2	0.04251 (15)	0.62057 (14)	0.41195 (9)	0.0539 (3)
O3	0.0526 (2)	0.21724 (19)	-0.01824 (12)	0.0791 (4)
O4	-0.15820 (15)	0.42413 (15)	0.08977 (10)	0.0561 (3)
N1	0.79790 (15)	0.14775 (14)	0.28682 (9)	0.0377 (2)
N2	0.61603 (16)	0.26801 (15)	0.32453 (9)	0.0391 (3)
H2A	0.5959	0.3291	0.3884	0.047*
N3	0.00969 (19)	0.32023 (16)	0.06233 (11)	0.0443 (3)
N4	0.20937 (16)	0.51447 (14)	0.39505 (10)	0.0393 (3)
C1	1.2885 (2)	-0.02327 (17)	0.37123 (11)	0.0381 (3)
C2	1.4763 (2)	-0.1424 (2)	0.33969 (14)	0.0499 (3)
H2B	1.5823	-0.1511	0.3819	0.060*
C3	1.5057 (2)	-0.2487 (2)	0.24485 (15)	0.0534 (4)

H3A	1.6336	-0.3321	0.2214	0.064*
C4	1.3510 (2)	-0.23481 (18)	0.18385 (13)	0.0482 (3)
H4A	1.3727	-0.3086	0.1187	0.058*
C5	1.1651 (2)	-0.11396 (17)	0.21735 (11)	0.0404 (3)
H5A	1.0595	-0.1052	0.1747	0.049*
C6	1.12930 (18)	-0.00419 (15)	0.31287 (10)	0.0332 (3)
C7	0.93392 (18)	0.12582 (16)	0.34949 (11)	0.0358 (3)
H7A	0.9091	0.1924	0.4185	0.043*
C8	0.46749 (17)	0.29040 (15)	0.26058 (10)	0.0329 (3)
C9	0.26980 (18)	0.40469 (15)	0.29198 (10)	0.0324 (3)
C10	0.12047 (18)	0.41764 (14)	0.22616 (10)	0.0336 (3)
H10A	-0.0115	0.4954	0.2487	0.040*
C11	0.16630 (19)	0.31661 (15)	0.12820 (11)	0.0345 (3)
C12	0.3609 (2)	0.20726 (17)	0.09136 (11)	0.0396 (3)
H12A	0.3909	0.1412	0.0216	0.048*
C13	0.50799 (19)	0.19515 (17)	0.15534 (11)	0.0390 (3)
H13A	0.6408	0.1213	0.1290	0.047*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0485 (5)	0.0679 (6)	0.0446 (5)	-0.0099 (4)	-0.0184 (4)	-0.0156 (4)
O1	0.0428 (6)	0.0767 (7)	0.0512 (6)	-0.0123 (5)	-0.0111 (5)	-0.0292 (5)
O2	0.0424 (6)	0.0553 (6)	0.0517 (6)	0.0007 (5)	-0.0030 (4)	-0.0189 (5)
O3	0.0686 (8)	0.0871 (9)	0.0790 (9)	-0.0060 (7)	-0.0370 (7)	-0.0388 (7)
O4	0.0380 (5)	0.0613 (7)	0.0664 (7)	-0.0057 (5)	-0.0209 (5)	-0.0032 (5)
N1	0.0290 (5)	0.0430 (6)	0.0384 (6)	-0.0080 (4)	-0.0046 (4)	-0.0036 (4)
N2	0.0292 (5)	0.0480 (6)	0.0373 (5)	-0.0077 (4)	-0.0053 (4)	-0.0108 (4)
N3	0.0440 (6)	0.0463 (6)	0.0455 (6)	-0.0131 (5)	-0.0174 (5)	-0.0022 (5)
N4	0.0348 (5)	0.0427 (6)	0.0392 (6)	-0.0121 (4)	-0.0013 (4)	-0.0110 (4)
C1	0.0385 (6)	0.0426 (6)	0.0338 (6)	-0.0116 (5)	-0.0102 (5)	-0.0005 (5)
C2	0.0391 (7)	0.0540 (8)	0.0529 (8)	-0.0048 (6)	-0.0174 (6)	0.0011 (6)
C3	0.0407 (7)	0.0461 (7)	0.0613 (9)	0.0020 (6)	-0.0060 (6)	-0.0041 (7)
C4	0.0529 (8)	0.0415 (7)	0.0455 (8)	-0.0103 (6)	-0.0031 (6)	-0.0091 (6)
C5	0.0408 (7)	0.0430 (7)	0.0393 (7)	-0.0136 (5)	-0.0107 (5)	-0.0040 (5)
C6	0.0317 (6)	0.0357 (6)	0.0329 (6)	-0.0113 (5)	-0.0061 (4)	0.0007 (4)
C7	0.0333 (6)	0.0418 (6)	0.0331 (6)	-0.0129 (5)	-0.0054 (5)	-0.0042 (5)
C8	0.0286 (5)	0.0363 (6)	0.0339 (6)	-0.0108 (5)	-0.0044 (4)	-0.0034 (4)
C9	0.0313 (6)	0.0335 (5)	0.0318 (6)	-0.0103 (4)	-0.0027 (4)	-0.0058 (4)
C10	0.0298 (6)	0.0313 (5)	0.0382 (6)	-0.0076 (4)	-0.0056 (4)	-0.0021 (4)
C11	0.0354 (6)	0.0335 (6)	0.0363 (6)	-0.0104 (5)	-0.0115 (5)	-0.0006 (5)
C12	0.0394 (7)	0.0403 (6)	0.0361 (6)	-0.0071 (5)	-0.0077 (5)	-0.0092 (5)
C13	0.0316 (6)	0.0416 (6)	0.0385 (6)	-0.0044 (5)	-0.0045 (5)	-0.0090 (5)

Geometric parameters (\AA , $^\circ$)

F1—C1	1.3555 (15)	C3—H3A	0.9500
O1—N4	1.2343 (15)	C4—C5	1.3785 (19)

O2—N4	1.2161 (15)	C4—H4A	0.9500
O3—N3	1.2204 (16)	C5—C6	1.3967 (17)
O4—N3	1.2225 (15)	C5—H5A	0.9500
N1—C7	1.2724 (16)	C6—C7	1.4618 (17)
N1—N2	1.3653 (15)	C7—H7A	0.9500
N2—C8	1.3548 (16)	C8—C9	1.4131 (17)
N2—H2A	0.8800	C8—C13	1.4181 (16)
N3—C11	1.4463 (16)	C9—C10	1.3863 (16)
N4—C9	1.4500 (15)	C10—C11	1.3685 (17)
C1—C2	1.3783 (19)	C10—H10A	0.9500
C1—C6	1.3800 (17)	C11—C12	1.3916 (18)
C2—C3	1.381 (2)	C12—C13	1.3597 (18)
C2—H2B	0.9500	C12—H12A	0.9500
C3—C4	1.379 (2)	C13—H13A	0.9500
C7—N1—N2	117.04 (11)	C1—C6—C5	116.70 (11)
C8—N2—N1	117.99 (10)	C1—C6—C7	121.17 (11)
C8—N2—H2A	121.0	C5—C6—C7	122.12 (11)
N1—N2—H2A	121.0	N1—C7—C6	118.51 (11)
O3—N3—O4	123.15 (12)	N1—C7—H7A	120.7
O3—N3—C11	117.69 (12)	C6—C7—H7A	120.7
O4—N3—C11	119.16 (11)	N2—C8—C9	123.98 (11)
O2—N4—O1	122.32 (11)	N2—C8—C13	119.48 (11)
O2—N4—C9	119.24 (11)	C9—C8—C13	116.55 (11)
O1—N4—C9	118.44 (11)	C10—C9—C8	121.82 (10)
F1—C1—C2	118.20 (12)	C10—C9—N4	115.72 (11)
F1—C1—C6	118.24 (11)	C8—C9—N4	122.45 (11)
C2—C1—C6	123.55 (12)	C11—C10—C9	118.91 (11)
C1—C2—C3	118.00 (13)	C11—C10—H10A	120.5
C1—C2—H2B	121.0	C9—C10—H10A	120.5
C3—C2—H2B	121.0	C10—C11—C12	121.27 (11)
C4—C3—C2	120.60 (14)	C10—C11—N3	119.78 (11)
C4—C3—H3A	119.7	C12—C11—N3	118.93 (11)
C2—C3—H3A	119.7	C13—C12—C11	119.89 (11)
C5—C4—C3	120.00 (13)	C13—C12—H12A	120.1
C5—C4—H4A	120.0	C11—C12—H12A	120.1
C3—C4—H4A	120.0	C12—C13—C8	121.44 (11)
C4—C5—C6	121.14 (12)	C12—C13—H13A	119.3
C4—C5—H5A	119.4	C8—C13—H13A	119.3
C6—C5—H5A	119.4	 	
C7—N1—N2—C8	-178.38 (11)	N2—C8—C9—N4	2.32 (19)
F1—C1—C2—C3	-179.24 (13)	C13—C8—C9—N4	-177.74 (11)
C6—C1—C2—C3	-0.3 (2)	O2—N4—C9—C10	-6.98 (17)
C1—C2—C3—C4	0.2 (2)	O1—N4—C9—C10	173.40 (12)
C2—C3—C4—C5	0.0 (2)	O2—N4—C9—C8	173.56 (12)
C3—C4—C5—C6	-0.1 (2)	O1—N4—C9—C8	-6.07 (19)
F1—C1—C6—C5	179.17 (11)	C8—C9—C10—C11	0.07 (18)

C2—C1—C6—C5	0.3 (2)	N4—C9—C10—C11	-179.40 (11)
F1—C1—C6—C7	-0.39 (18)	C9—C10—C11—C12	-2.71 (19)
C2—C1—C6—C7	-179.28 (13)	C9—C10—C11—N3	176.08 (11)
C4—C5—C6—C1	0.0 (2)	O3—N3—C11—C10	-173.46 (14)
C4—C5—C6—C7	179.51 (12)	O4—N3—C11—C10	6.06 (19)
N2—N1—C7—C6	178.86 (10)	O3—N3—C11—C12	5.4 (2)
C1—C6—C7—N1	173.99 (12)	O4—N3—C11—C12	-175.12 (12)
C5—C6—C7—N1	-5.54 (19)	C10—C11—C12—C13	2.3 (2)
N1—N2—C8—C9	177.09 (11)	N3—C11—C12—C13	-176.51 (12)
N1—N2—C8—C13	-2.85 (18)	C11—C12—C13—C8	0.8 (2)
N2—C8—C9—C10	-177.12 (11)	N2—C8—C13—C12	176.68 (12)
C13—C8—C9—C10	2.82 (18)	C9—C8—C13—C12	-3.3 (2)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N2—H2A···O1	0.88	2.02	2.6317 (15)	126
N2—H2A···O1 ⁱ	0.88	2.51	3.3424 (15)	158
C2—H2B···F1 ⁱⁱ	0.95	2.45	3.3386 (17)	156
C3—H3A···O4 ⁱⁱⁱ	0.95	2.48	3.3177 (19)	148
C5—H5A···O3 ^{iv}	0.95	2.43	3.2694 (17)	148

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+3, -y, -z+1$; (iii) $x+2, y-1, z$; (iv) $-x+1, -y, -z$.