

(3a*R*,4*S*,7*R*,7a*S*)-2-Phenyl-4-propyl-3a,4,7,7a-tetrahydro-1*H*-4,7-epithioisoindole-1,3-dione 8-oxide 8-oxide

Aydin Demircan,^a Ertan Sahin,^b Gozde Beyazova,^a Muhsin Karaaslan^c and Tuncer Hökelek^{d*}

^aDepartment of Chemistry, Niğde University, 51100 Niğde, Turkey, ^bDepartment of Chemistry, Atatürk University, 25240 Erzurum, Turkey, ^cDepartment of Chemistry, Aksaray University, 68100 Aksaray, Turkey, and ^dDepartment of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 23 March 2011; accepted 6 April 2011

Key indicators: single-crystal X-ray study; $T = 294\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.047; wR factor = 0.108; data-to-parameter ratio = 15.6.

In the tetrahydroisoindole moiety of the title compound, $C_{17}H_{17}NO_3S$, the six-membered ring assumes a boat configuration and the $\text{S}=\text{O}$ group bridges the prow and stern of the boat. The phenyl ring is oriented at a dihedral angle of $83.2(1)^\circ$ with respect to the pyrrole ring. In the crystal, intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into a three-dimensional network. A weak $\text{C}-\text{H}\cdots\pi$ interaction involving the phenyl ring is also found. The crystal studied was an inversion twin.

Related literature

For background to the thiophenen system, see: Lert & Trindle (1971). For the conditions of cycloaddition reactions of thiophene, see: Al-Omran *et al.* (1996); Kuhn & Gollnick (1972); Kotsuki *et al.* (1978); Thiemann *et al.* (1995). For the biological activity of some thiophene 1,1-dioxide derivatives, see: Thiemann *et al.* (2009). For thiophene *s*-oxides with alkyl groups at positions 2,3,4 and 5, see: Rajappa (1984). For related structures, see: Arslan & Demircan (2007); Koşar *et al.* (2006).



Experimental

Crystal data

$C_{17}H_{17}NO_3S$

$M_r = 315.39$

Orthorhombic, $P2_12_12_1$

$a = 7.7712(3)\text{ \AA}$

$b = 10.8413(3)\text{ \AA}$

$c = 18.9762(4)\text{ \AA}$

$V = 1598.74(8)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.21\text{ mm}^{-1}$

$T = 294\text{ K}$

$0.30 \times 0.25 \times 0.20\text{ mm}$

Data collection

Rigaku R-AXIS RAPID-S

diffractometer

Absorption correction: multi-scan

(Blessing, 1995)

$R_{\min} = 0.807$, $T_{\max} = 0.865$

34450 measured reflections

3279 independent reflections

2686 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.088$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$

$wR(F^2) = 0.108$

$S = 1.08$

3279 reflections

210 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.20\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.22\text{ e \AA}^{-3}$

Absolute structure: Flack (1983),

1379 Friedel pairs

Flack parameter: 0.37 (9)

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$ is the centroid of the C12–C17 phenyl ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C5–H5 \cdots O1 ⁱ	0.95 (2)	2.51 (3)	3.326 (4)	144 (2)
C17–H17 \cdots O3 ⁱⁱ	0.93	2.57	3.315 (3)	137
C7–H7 \cdots Cg1 ⁱⁱⁱ	0.98	2.77	3.693 (3)	157

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x + \frac{1}{2}, -y + \frac{1}{2}, -z$; (iii) $x - 1, y, z$.

Data collection: *CrystalClear* (Rigaku/MSC, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).

The authors are indebted to the Department of Chemistry, Atatürk University, Erzurum, Turkey, for use of the X-ray diffractometer purchased under grant No. 2003/219 of the University Research Fund. We thank The Scientific & Technological Research Council of Turkey (TUBITAK) for financial support of this work (PN: 107T831).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5179).

References

- Al-Omran, F., Khalik, M. M. A., Al-Awadhi, H. & Elnagdi, M. H. (1996). *Tetrahedron*, **52**, 11915–11928.
- Arslan, H. & Demircan, A. (2007). *Acta Chim. Slov.* **54**, 341–353.
- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Koşar, B., Göktürk, E., Demircan, A. & Büyükgüngör, O. (2006). *Acta Cryst. E* **62**, o4192–o4193.
- Kotsuki, H., Kitagawa, S., Nishizawa, H. & Tokoroyama, T. (1978). *J. Org. Chem.*, **43**, 1471–1472.
- Kuhn, H. J. & Gollnick, K. (1972). *Tetrahedron Lett.*, pp. 1909–1912.

organic compounds

- Lert, P. W. & Trindle, C. (1971). *J. Am. Chem. Soc.* **93**, 6392–6395.
Rajappa, S. (1984). *Comprehensive Heterocyclic Chemistry*, Vol. 4, edited by A. R. Katritzky & C. W. Rees, p. 841. New York: Pergamon Press.
Rigaku/MSC (2005). *CrystalClear*. Rigaku/MSC, The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
Thiemann, T., Li, Y. Q., Mataka, S. & Tashiro, M. (1995). *J. Chem. Res. (M)*, pp. 6364–6379.
Thiemann, T., Tanaka, Y. & Iniesta, J. (2009). *Molecules*, **14**, 1013–1031.

supporting information

Acta Cryst. (2011). E67, o1085–o1086 [doi:10.1107/S1600536811012876]

(3a*R*,4*S*,7*R*,7a*S*)-2-Phenyl-4-propyl-3a,4,7,7a-tetrahydro-1*H*-4,7-epithioiso-indole-1,3-dione 8-oxide

Aydin Demircan, Ertan Şahin, Gözde Beyazova, Muhsin Karaaslan and Tuncer Hökelek

S1. Comment

Thiophene behaves as the least reactive diene among all aromatic five-membered heterocycles; the presence of 3d orbitals on sulfur contributes to the resonance stability of the thiophene system (Lert & Trindle, 1971). Cycloaddition reactions of thiophene also have limitations therefore its 4 + 2 cycloaddition can occur under following conditions; use of highly reactive dienophiles (Thiemann *et al.*, 1995), enhance electron density on the thiophene by having substituent at position 2 or 5 (Al-Omran *et al.*, 1996), high reaction temperature (Kuhn & Gollnick, 1972) or use of high pressure (Kotsuki *et al.*, 1978). However, the number of publications dealing with the chemistry of thiophene dioxide based new materials, namely, semiconductor organic transistors and light diodes first designed about 13 years ago. New data on the biological activities of some thiophene 1,1-dioxide derivatives were observed in recent years (Thiemann *et al.*, 2009).

Thiophene *s*-oxides are derivatives of thiophene, which belong to an important group of five-membered heterocyclic compounds having non-aromatic character. Thiophene *s*-oxides with alkyl groups at positions 2,3,4 and 5 have been the subject of extensive studies in recent past (Rajappa, 1984). Owing to the unique structure combined with high reactivity, they can be used to prepare various heterocyclic systems; therefore, these compounds are useful building blocks in synthetic organic chemistry. In continuation of our research program and following our previous interest in the syntheses of fused heterocyclic compounds using furan (Arslan & Demircan, 2007; Koşar *et al.*, 2006), it has been found that mono alkylated thiophene at position 2 will lead to an excellent building block for the synthesis of the title compound.

The title compound contains two non-planar five- and six-membered rings, which has a pyrrole (C1/N2/C3/C3a/C7a) ring (B) on one side and a propyl moiety at position 4. It also contains a phenyl (C12-C17) ring (A) bonded to the pyrrole ring (B) at position 2. The angles C4-C5-C6 [111.4 (3) $^{\circ}$] and C5-C6-C7 [111.1 (3) $^{\circ}$] about the double bond have an average value of 111.3 (3) $^{\circ}$. The dihedral angles between planes A, B, C (C3a/C4/C7a C7), D (C4-C7) and E (C4/S8/C7) are as follows: A/B = 83.2 (1) $^{\circ}$, B/C = 61.9 (1) $^{\circ}$, C/D = 60.6 (1) $^{\circ}$, C/E = 58.8 (1) $^{\circ}$ and D/E = 60.7 (1) $^{\circ}$.

In the crystal, intermolecular C—H \cdots O hydrogen bonds link the molecules to form a three-dimensional network (Table 1 and Fig. 2). There also exist a weak C—H \cdots π interaction involving the phenyl ring A (C12-C17) (Table 1).

S2. Experimental

For the preparation of the title compound, $\text{BF}_3\text{-Et}_2\text{O}$ (6.1 ml, 47.75 mmol) was added slowly to a solution of 2-propylthiophene (1.00 g, 7.90 mmol) and N-phenylmaleimide (2.06 g, 11.90 mmol) in dry dichloromethane (DCM) (40 ml) under an inert atmosphere at 253 K. The reaction mixture was stirred for 10 min at 253 K, and then a solution of meta chloroperbenzoic acid (*m*-CPBA) (2.05 g, 11.90 mmol) in dry DCM (40 ml) was added slowly. The reaction mixture was stirred for 2 h at 253 K, and then the suspension was poured into a mixture of aqueous NaHCO_3 solution (80 ml) and DCM (50 ml) and stirred at room temperature for 20 min. The organic phase was separated, and the aqueous phase was extracted with DCM (3 x 25 ml). The combined organic phase was washed with water and brine, and then dried over anhydrous

MgSO_4 . After removal of the solvent *in vacuo*, the residue was chromatographed on silica gel to give the title compound as colorless crystals.

S3. Refinement

The H atoms H5 and H6 were located in a difference Fourier map and were freely refined. Other C-bound H-atoms were positioned geometrically with C—H = 0.93, 0.96, 0.97 and 0.98 Å, for aromatic, methyl, methylene and methine H-atoms, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C})$, where $k = 1.5$ for methyl H-atoms and $k = 1.2$ for all other H-atoms.





Figure 1

The molecular structure of the title molecule with the atom-numbering scheme. Displacement ellipsoids are drawn at the 40% probability level.

**Figure 2**

A view of the crystal packing of the title compound. The C-H...O hydrogen bonds are shown as dashed lines [H-atoms not involved in hydrogen bonding have been omitted for clarity].

(3a*R*,4*S*,7*R*,7a*S*)-2-Phenyl-4-propyl-3*a*,4,7,7*a*-tetrahydro-1*H*-4,7-epithioisoindole-1,3-dione 8-oxide

Crystal data



$M_r = 315.39$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 7.7712 (3) \text{ \AA}$

$b = 10.8413 (3) \text{ \AA}$

$c = 18.9762 (4) \text{ \AA}$

$V = 1598.74 (8) \text{ \AA}^3$

$Z = 4$

$F(000) = 664$

$D_x = 1.310 \text{ Mg m}^{-3}$

Mo $\text{K}\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 6907 reflections

$\theta = 2.2\text{--}26.4^\circ$

$\mu = 0.21 \text{ mm}^{-1}$

$T = 294 \text{ K}$

Block, colorless

$0.30 \times 0.25 \times 0.20 \text{ mm}$

Data collection

Rigaku R-AXIS RAPID-S
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan
(Blessing, 1995)

$T_{\min} = 0.807$, $T_{\max} = 0.865$

34450 measured reflections

3279 independent reflections

2686 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.088$

$\theta_{\max} = 26.4^\circ$, $\theta_{\min} = 2.2^\circ$

$h = -9 \rightarrow 9$

$k = -13 \rightarrow 13$

$l = -23 \rightarrow 23$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.047$$

$$wR(F^2) = 0.108$$

$$S = 1.08$$

3279 reflections

210 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0453P)^2 + 0.1086P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.20 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$$

Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.018 (2)

Absolute structure: Flack (1983), 1379 Friedel
pairs

Absolute structure parameter: 0.37 (9)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.0424 (3)	0.0571 (2)	0.16742 (11)	0.0742 (6)
O2	0.3289 (2)	0.35385 (18)	0.09320 (10)	0.0708 (6)
O3	-0.3050 (2)	0.40176 (16)	0.00144 (9)	0.0633 (5)
C1	0.0032 (3)	0.1488 (3)	0.13751 (14)	0.0553 (6)
N2	0.1690 (3)	0.1987 (2)	0.14224 (10)	0.0528 (5)
C3	0.1923 (3)	0.3011 (2)	0.10005 (12)	0.0526 (6)
C3A	0.0230 (3)	0.3335 (2)	0.06695 (13)	0.0488 (6)
H3A	0.0324	0.3349	0.0155	0.059*
C4	-0.0495 (3)	0.4576 (2)	0.09505 (13)	0.0541 (6)
C5	-0.0650 (4)	0.4517 (3)	0.17396 (15)	0.0654 (8)
H5	-0.003 (3)	0.503 (2)	0.2053 (14)	0.060 (8)*
C6	-0.1750 (4)	0.3652 (3)	0.19369 (15)	0.0659 (8)
H6	-0.216 (4)	0.346 (3)	0.2393 (16)	0.083 (9)*
C7	-0.2475 (3)	0.2998 (2)	0.13186 (13)	0.0578 (7)
H7	-0.3499	0.2498	0.1414	0.069*
C7A	-0.1015 (3)	0.2327 (2)	0.09100 (12)	0.0509 (6)
H7A	-0.1479	0.1875	0.0505	0.061*
S8	-0.28413 (8)	0.43745 (6)	0.07643 (3)	0.0564 (2)
C9	0.0304 (4)	0.5709 (3)	0.06204 (17)	0.0716 (8)
H9A	0.1537	0.5675	0.0698	0.086*
H9B	0.0113	0.5671	0.0116	0.086*
C10	-0.0343 (5)	0.6940 (3)	0.0883 (2)	0.1081 (13)

H10A	-0.1584	0.6967	0.0831	0.130*
H10B	-0.0081	0.7014	0.1381	0.130*
C11	0.0429 (8)	0.8016 (4)	0.0500 (3)	0.176 (3)
H11A	-0.0063	0.8768	0.0675	0.264*
H11B	0.0192	0.7943	0.0005	0.264*
H11C	0.1650	0.8026	0.0574	0.264*
C12	0.3072 (3)	0.1422 (2)	0.18063 (14)	0.0580 (7)
C13	0.3496 (4)	0.1844 (3)	0.24562 (15)	0.0729 (9)
H13	0.2888	0.2488	0.2663	0.087*
C14	0.4881 (5)	0.1281 (4)	0.2808 (2)	0.0951 (12)
H14	0.5197	0.1554	0.3255	0.114*
C15	0.5753 (5)	0.0350 (4)	0.2504 (3)	0.1005 (14)
H15	0.6669	-0.0010	0.2743	0.121*
C16	0.5321 (5)	-0.0070 (4)	0.1856 (2)	0.0974 (12)
H16	0.5939	-0.0710	0.1651	0.117*
C17	0.3949 (4)	0.0460 (3)	0.15000 (16)	0.0732 (8)
H17	0.3626	0.0168	0.1059	0.088*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0578 (11)	0.0808 (14)	0.0841 (14)	-0.0097 (11)	-0.0041 (10)	0.0270 (12)
O2	0.0467 (10)	0.0775 (13)	0.0883 (14)	-0.0098 (9)	0.0005 (9)	0.0129 (11)
O3	0.0654 (12)	0.0712 (12)	0.0533 (10)	0.0027 (10)	-0.0088 (9)	-0.0030 (9)
C1	0.0522 (15)	0.0627 (17)	0.0509 (14)	-0.0031 (12)	0.0019 (12)	0.0021 (13)
N2	0.0457 (11)	0.0591 (12)	0.0537 (12)	-0.0010 (9)	-0.0012 (9)	0.0057 (10)
C3	0.0486 (15)	0.0587 (14)	0.0506 (13)	0.0003 (13)	0.0024 (11)	-0.0015 (11)
C3A	0.0463 (13)	0.0526 (14)	0.0476 (13)	0.0001 (10)	0.0013 (11)	0.0006 (11)
C4	0.0499 (14)	0.0552 (15)	0.0573 (15)	-0.0011 (12)	0.0010 (12)	-0.0058 (12)
C5	0.0577 (16)	0.081 (2)	0.0572 (16)	0.0093 (16)	-0.0064 (14)	-0.0204 (15)
C6	0.0559 (17)	0.090 (2)	0.0521 (16)	0.0140 (16)	0.0059 (13)	-0.0032 (15)
C7	0.0482 (15)	0.0675 (16)	0.0578 (15)	-0.0004 (12)	0.0037 (12)	0.0069 (13)
C7A	0.0458 (13)	0.0571 (14)	0.0499 (14)	-0.0020 (11)	-0.0026 (11)	0.0012 (11)
S8	0.0511 (4)	0.0604 (4)	0.0576 (4)	0.0037 (3)	-0.0014 (3)	-0.0029 (3)
C9	0.0703 (18)	0.0600 (17)	0.084 (2)	-0.0091 (15)	0.0010 (15)	-0.0013 (16)
C10	0.111 (3)	0.0577 (19)	0.156 (4)	-0.0059 (19)	0.021 (3)	-0.007 (2)
C11	0.196 (6)	0.061 (3)	0.272 (7)	-0.023 (3)	0.070 (5)	0.004 (3)
C12	0.0467 (14)	0.0694 (17)	0.0579 (15)	-0.0037 (13)	-0.0035 (12)	0.0137 (13)
C13	0.0675 (18)	0.090 (2)	0.0611 (17)	-0.0115 (17)	-0.0119 (14)	0.0085 (15)
C14	0.081 (3)	0.129 (3)	0.075 (2)	-0.035 (2)	-0.026 (2)	0.033 (2)
C15	0.062 (2)	0.127 (4)	0.112 (3)	0.000 (2)	-0.017 (2)	0.057 (3)
C16	0.076 (2)	0.113 (3)	0.103 (3)	0.025 (2)	-0.004 (2)	0.027 (2)
C17	0.0665 (18)	0.079 (2)	0.0737 (19)	0.0133 (17)	-0.0027 (15)	0.0094 (17)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.198 (3)	S8—C7	1.848 (3)
O2—C3	1.213 (3)	C9—C10	1.511 (4)

C1—C7A	1.506 (3)	C9—H9A	0.9700
N2—C1	1.401 (3)	C9—H9B	0.9700
N2—C3	1.380 (3)	C10—C11	1.500 (6)
N2—C12	1.435 (3)	C10—H10A	0.9700
C3A—C3	1.500 (3)	C10—H10B	0.9700
C3A—C4	1.553 (3)	C11—H11A	0.9600
C3A—C7A	1.529 (3)	C11—H11B	0.9600
C3A—H3A	0.9800	C11—H11C	0.9600
C4—C5	1.504 (4)	C12—C13	1.356 (4)
C4—C9	1.511 (4)	C12—C17	1.375 (4)
C5—H5	0.95 (3)	C13—C14	1.406 (5)
C6—C5	1.323 (5)	C13—H13	0.9300
C6—H6	0.95 (3)	C14—H14	0.9300
C7—C6	1.482 (4)	C15—C14	1.346 (5)
C7—C7A	1.555 (3)	C15—C16	1.353 (6)
C7—H7	0.9800	C15—H15	0.9300
C7A—H7A	0.9800	C16—H16	0.9300
S8—O3	1.4836 (18)	C17—C16	1.387 (4)
S8—C4	1.871 (3)	C17—H17	0.9300
O1—C1—N2	124.2 (2)	O3—S8—C4	108.56 (11)
O1—C1—C7A	128.2 (2)	O3—S8—C7	110.62 (11)
N2—C1—C7A	107.5 (2)	C7—S8—C4	80.60 (11)
C1—N2—C12	123.8 (2)	C4—C9—H9A	108.2
C3—N2—C1	113.2 (2)	C4—C9—H9B	108.2
C3—N2—C12	122.7 (2)	C10—C9—C4	116.4 (3)
O2—C3—N2	123.8 (2)	C10—C9—H9A	108.2
O2—C3—C3A	127.8 (2)	C10—C9—H9B	108.2
N2—C3—C3A	108.4 (2)	H9A—C9—H9B	107.3
C3—C3A—C4	112.2 (2)	C9—C10—H10A	108.9
C3—C3A—C7A	105.22 (19)	C9—C10—H10B	108.9
C3—C3A—H3A	110.8	C11—C10—C9	113.2 (3)
C4—C3A—H3A	110.8	C11—C10—H10A	108.9
C7A—C3A—C4	106.71 (19)	C11—C10—H10B	108.9
C7A—C3A—H3A	110.8	H10A—C10—H10B	107.7
C3A—C4—S8	100.79 (16)	C10—C11—H11A	109.5
C5—C4—S8	96.01 (18)	C10—C11—H11B	109.5
C5—C4—C3A	109.5 (2)	C10—C11—H11C	109.5
C5—C4—C9	118.7 (2)	H11A—C11—H11B	109.5
C9—C4—S8	114.65 (19)	H11A—C11—H11C	109.5
C9—C4—C3A	114.4 (2)	H11B—C11—H11C	109.5
C4—C5—H5	124.0 (16)	C13—C12—N2	120.0 (3)
C6—C5—C4	111.4 (3)	C13—C12—C17	121.3 (3)
C6—C5—H5	124.7 (15)	C17—C12—N2	118.7 (2)
C5—C6—C7	111.1 (3)	C12—C13—C14	118.2 (3)
C5—C6—H6	129.2 (19)	C12—C13—H13	120.9
C7—C6—H6	119.5 (19)	C14—C13—H13	120.9
S8—C7—H7	115.3	C13—C14—H14	119.8

C6—C7—S8	97.05 (19)	C15—C14—C13	120.4 (4)
C6—C7—C7A	110.0 (2)	C15—C14—H14	119.8
C6—C7—H7	115.3	C14—C15—C16	121.2 (4)
C7A—C7—S8	101.92 (16)	C14—C15—H15	119.4
C7A—C7—H7	115.3	C16—C15—H15	119.4
C1—C7A—C3A	105.4 (2)	C15—C16—C17	119.6 (4)
C1—C7A—C7	112.6 (2)	C15—C16—H16	120.2
C1—C7A—H7A	110.9	C17—C16—H16	120.2
C3A—C7A—C7	106.0 (2)	C12—C17—C16	119.3 (3)
C3A—C7A—H7A	110.9	C12—C17—H17	120.3
C7—C7A—H7A	110.9	C16—C17—H17	120.3
O1—C1—C7A—C3A	-179.5 (3)	C3A—C4—C5—C6	-62.6 (3)
O1—C1—C7A—C7	65.4 (4)	C9—C4—C5—C6	163.5 (3)
N2—C1—C7A—C3A	2.2 (3)	S8—C4—C9—C10	65.0 (3)
N2—C1—C7A—C7	-112.9 (2)	C3A—C4—C9—C10	-179.3 (3)
C3—N2—C1—O1	176.9 (3)	C5—C4—C9—C10	-47.5 (4)
C3—N2—C1—C7A	-4.8 (3)	C7—C6—C5—C4	0.7 (3)
C12—N2—C1—O1	3.5 (4)	S8—C7—C6—C5	-42.9 (3)
C12—N2—C1—C7A	-178.1 (2)	C7A—C7—C6—C5	62.5 (3)
C1—N2—C3—O2	-175.5 (2)	S8—C7—C7A—C1	156.02 (18)
C1—N2—C3—C3A	5.3 (3)	S8—C7—C7A—C3A	41.3 (2)
C12—N2—C3—O2	-2.1 (4)	C6—C7—C7A—C1	53.9 (3)
C12—N2—C3—C3A	178.8 (2)	C6—C7—C7A—C3A	-60.8 (3)
C1—N2—C12—C13	-100.8 (3)	O3—S8—C4—C3A	-51.85 (18)
C1—N2—C12—C17	79.6 (3)	O3—S8—C4—C5	-163.06 (17)
C3—N2—C12—C13	86.5 (3)	O3—S8—C4—C9	71.5 (2)
C3—N2—C12—C17	-93.1 (3)	C7—S8—C4—C3A	56.87 (16)
C4—C3A—C3—O2	-67.0 (3)	C7—S8—C4—C5	-54.33 (18)
C4—C3A—C3—N2	112.1 (2)	C7—S8—C4—C9	-179.8 (2)
C7A—C3A—C3—O2	177.3 (2)	O3—S8—C7—C6	161.81 (16)
C7A—C3A—C3—N2	-3.5 (3)	C4—S8—C7—C6	55.41 (17)
C3—C3A—C4—S8	-156.84 (17)	O3—S8—C7—C7A	49.62 (19)
C3—C3A—C4—C5	-56.4 (3)	C4—S8—C7—C7A	-56.78 (16)
C3—C3A—C4—C9	79.6 (3)	C4—C9—C10—C11	-176.4 (4)
C7A—C3A—C4—S8	-42.1 (2)	N2—C12—C13—C14	-178.7 (3)
C7A—C3A—C4—C5	58.3 (3)	C17—C12—C13—C14	0.9 (4)
C7A—C3A—C4—C9	-165.7 (2)	N2—C12—C17—C16	178.1 (3)
C3—C3A—C7A—C1	0.7 (3)	C13—C12—C17—C16	-1.5 (4)
C3—C3A—C7A—C7	120.3 (2)	C12—C13—C14—C15	0.0 (5)
C4—C3A—C7A—C1	-118.6 (2)	C16—C15—C14—C13	-0.2 (6)
C4—C3A—C7A—C7	1.0 (3)	C14—C15—C16—C17	-0.4 (6)
S8—C4—C5—C6	41.1 (3)	C12—C17—C16—C15	1.3 (5)

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C12–C17 phenyl ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C5—H5···O1 ⁱ	0.95 (2)	2.51 (3)	3.326 (4)	144 (2)
C17—H17···O3 ⁱⁱ	0.93	2.57	3.315 (3)	137
C7—H7···Cg1 ⁱⁱⁱ	0.98	2.77	3.693 (3)	157

Symmetry codes: (i) $-x, y+1/2, -z+1/2$; (ii) $x+1/2, -y+1/2, -z$; (iii) $x-1, y, z$.