

4,6-Bis(4-fluorophenyl)-2-phenyl-1*H*-indazol-3(2*H*)-one

R. J. Butcher,^a M. Akkurt,^{b*} S. Samshuddin,^c B. Narayana^c and H. S. Yathirajan^d

^aDepartment of Chemistry, Howard University, 525 College Street NW, Washington, DC 20059, USA, ^bDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^cDepartment of Studies in Chemistry, Mangalore University, Mangalagangotri 574 199, India, and ^dDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India
Correspondence e-mail: akkurt@erciyes.edu.tr


Received 26 March 2011; accepted 29 April 2011

Key indicators: single-crystal X-ray study; $T = 123\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.057; wR factor = 0.135; data-to-parameter ratio = 13.9.

In the title compound, $\text{C}_{25}\text{H}_{16}\text{F}_2\text{N}_2\text{O}$, the pyrazole ring is almost planar (r.m.s. deviation = 0.028 \AA) and makes a dihedral angle of $5.86(11)^\circ$ with the indazole benzene ring. The dihedral angle between the pyrazole ring and the unsubstituted phenyl ring is $28.19(11)^\circ$. The dihedral angles between the unsubstituted phenyl and the two fluorophenyl groups are $57.69(10)$ and $18.01(10)^\circ$. In the crystal, molecules are linked by intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{F}$ interactions, forming infinite chains along the b axis with graph-set motif $R_3^2(19)$. The crystal structure is further consolidated by $\pi-\pi$ stacking [centroid–centroid distances = $3.5916(13)$ and $3.6890(13)\text{ \AA}$] and $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For the pharmacological activity of indazole derivatives, see: Beylin *et al.* (1991); George *et al.* (1998); Jain *et al.* (1987); Palazzo *et al.* (1966); Popat *et al.* (2003); Roman (1990). For related structures, see: van der Helm *et al.* (1979); Fun *et al.* (2010). For hybridization and electron delocalization around N atoms, see: Susindran *et al.* (2010); Jin *et al.* (2004). For graph-set analysis, see: Etter (1990); Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{25}\text{H}_{16}\text{F}_2\text{N}_2\text{O}$	$V = 3723.23(15)\text{ \AA}^3$
$M_r = 398.40$	$Z = 8$
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation
$a = 15.2947(4)\text{ \AA}$	$\mu = 0.10\text{ mm}^{-1}$
$b = 11.6259(2)\text{ \AA}$	$T = 123\text{ K}$
$c = 20.9388(5)\text{ \AA}$	$0.49 \times 0.38 \times 0.23\text{ mm}$

Data collection

Oxford Diffraction Xcalibur Ruby Gemini diffractometer	19870 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2007)	3827 independent reflections
$R_{\text{int}} = 0.029$	3416 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.895$, $T_{\max} = 0.977$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.135$	$\Delta\rho_{\max} = 0.65\text{ e \AA}^{-3}$
$S = 1.09$	$\Delta\rho_{\min} = -0.35\text{ e \AA}^{-3}$
3827 reflections	
275 parameters	
1 restraint	

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg5$ is the centroid of the C20–C25 phenyl ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—HN2 \cdots O1 ⁱ	0.86 (2)	2.00 (2)	2.830 (2)	162 (2)
C6—H6 \cdots F1 ⁱⁱ	0.93	2.49	3.362 (2)	156
C15—H15 \cdots Cg5 ⁱⁱⁱ	0.93	2.85	3.656 (2)	145

Symmetry codes: (i) $-x + \frac{3}{2}, y + \frac{1}{2}, z$; (ii) $x, y + 1, z$; (iii) $x, -y, z + \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2007); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2007); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).

SS and BN thank Mangalore University and the UGC SAP for financial assistance for the purchase of chemicals. HSY thanks the UOM for sabbatical leave.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: QK2004).

References

- Altomare, A., Burla, M. C., Camalli, M., Carrozzini, B., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Rizzi, R. (1999). *J. Appl. Cryst.* **32**, 339–340.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Beylin, V. G., Colbry, N. L., Giordani, A. B., Goel, O. P., Johnson, D. R., Leeds, R. L., Leja, B., Lewis, E. P., Lustgarten, D. M., Showalter, H. D. H., Serel, A. D., Reily, M. D., Uhlenhorst, S. E. & Zisek, K. A. (1991). *J. Heterocycl. Chem.* **28**, 517–527.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.

- Fun, H.-K., Hemamalini, M., Samshuddin, S., Narayana, B. & Yathirajan, H. S. (2010). *Acta Cryst.* **E66**, o864–o865.
- George, V. D., Kim, U. T., Liang, J., Cordova, B., Klabe, R. M., Garber, S., Bacheler, L. T., Lam, G. N., Wright, M. R., Logue, K. A., Viitanen, S. E., Ko, S. S. & Trainor, G. L. (1998). *J. Med. Chem.* **41**, 2411–2423.
- Helm, D. van der, Wu, K. K., Ealick, S. E., Berlin, K. D. & Ramalingam, K. (1979). *Acta Cryst.* **B35**, 2804–2806.
- Jain, A. C., Mehta, A. & Arya, P. (1987). *Indian J. Chem. Sect. B*, **26**, 150–153.
- Jin, Z.-M., Li, L., Li, M.-C., Hu, M.-L. & Shen, L. (2004). *Acta Cryst.* **C60**, o642–o643.
- Oxford Diffraction (2007). *CrysAlis PRO* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
- Palazzo, G., Corsi, G., Baiocchi, L. & Silnerstrini, B. (1966). *J. Med. Chem.* **9**, 38–41.
- Popat, K. H., Nimavat, K. S., Vasoya, S. L. & Joshi, H. S. (2003). *Indian J. Chem. Sect. B*, **42**, 1497–1501.
- Roman, B. (1990). *Pharmazie*, **45**, 214–217.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Susindran, V., Athimoolam, S., Bahadur, S. A., Manikannan, R. & Muthusubramanian, S. (2010). *Acta Cryst.* **E66**, o2594–o2595.

supporting information

Acta Cryst. (2011). E67, o1346–o1347 [doi:10.1107/S1600536811016369]

4,6-Bis(4-fluorophenyl)-2-phenyl-1*H*-indazol-3(2*H*)-one

R. J. Butcher, M. Akkurt, S. Samshuddin, B. Narayana and H. S. Yathirajan

S1. Comment

In last few decades, much attention has been paid to the synthesis of heterocycles containing 1,2-diazole systems like indazole mainly due to their broad spectrum of pharmacological properties. Indazole derivatives possess variety of pharmacological activities such as analgesic, anti inflammatory, antidepressant, antihypertensive, antiviral and anticancer activities (Jain *et al.*, 1987; Palazzo *et al.*, 1966; Popat *et al.*, 2003; Beylin *et al.*, 1991; George *et al.*, 1998; Roman, 1990).

The crystal structure of indazole derivative, *viz.*, 1,2,4,5-tetrahydro-7-methoxy-3*H*-benz[g]indazol-3-one monohydrate (van der Helm *et al.*, 1979) has been reported. Also the crystal structure of methyl 4,6-bis(4-fluorophenyl)-2-oxocyclohex-3-ene-1-carboxylate, the precursor of the title compound, has been reported (Fun *et al.*, 2010). In view of the importance of indazole derivatives, the title compound (**I**) is synthesized and its crystal structure is reported here.

The pyrazole ring (N1/N2/C1/C2/C7) in (**I**), (Fig. 1), is almost planar with the largest deviations from the mean plane being 0.039 (2) Å for C7 and -0.035 (2) Å for N2. The sum of the surrounding angles around N1 in the pyrazole ring is 358.15 (15)°, in accordance with the *sp*² hybridization of the N1 atom (Susindran *et al.*, 2010). The C1—N1 and C7—N2 bond lengths in the pyrazole ring are 1.390 (2) and 1.367 (3) Å, respectively. The values of these distances are shorter than the pertinent single bond length of 1.443 Å and are longer than the double bond length of 1.269 Å (Jin *et al.*, 2004). This case indicates electron delocalization.

The dihedral angle between the pyrazole ring and the indazole benzene ring in (**I**), (Fig. 1), is 5.86 (11)°. The dihedral angle between the two fluorophenyl groups is 42.56 (11)°, and the dihedral angle between the five-membered pyrazole ring and the unsubstituted phenyl ring is 28.19 (11)°. The unsubstituted phenyl ring and the two fluorophenyl groups make dihedral angles of 57.69 (10) and 18.01 (10)°, respectively, with each other.


In the crystal structure, molecules are linked by intermolecular N—H···O and C—H···F interactions, forming *R*₂₃(19) graph-set motifs (Etter, 1990; Bernstein *et al.*, 1995) along the *b* axis of the unit cell (Table 1, Fig. 2). In addition, the crystal structure is consolidated by C—H···π and π···π stacking [*Cg*1···*Cg*3(3/2 - *x*, 1/2 + *y*, *z*) = 3.5916 (13) Å and *Cg*3···*Cg*3(1 - *x*, *y*, 1/2 - *z*) = 3.6890 (13) Å; *Cg*1 and *Cg*3 are the centroids of the N1/N2/C1/C2/C7 pyrazole ring and C8—C13 benzene ring, respectively] interactions.

S2. Experimental


A mixture of methyl 4,6-bis(4-fluorophenyl)-2-oxocyclohex-3-ene-1-carboxylate (3.42 g, 0.01 mol) and phenyl hydrazine (1.08 g, 0.01 mol) in 50 ml ethanol containing 1 ml glacial acetic acid was refluxed for 10 h. The reaction mixture was cooled and poured into 50 ml ice-cold water. The precipitate was collected by filtration and purified by recrystallization from ethanol. Yellow prisms of (**I**) were grown from DMF by slow evaporation (m.p.: > 523 K, yield: 58%).

S3. Refinement

All H atoms attached to C atoms were placed in their calculated positions (aromatic C—H = 0.93 Å) and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The N-bound H atom was located from a difference map and refined with a distance restraint N—H = 0.86±0.01 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$.

**Figure 1**

Molecular structure of the title compound showing the atom labeling scheme. Displacement ellipsoids for non-H atoms are drawn at the 50% probability level.

**Figure 2**

A partial packing diagram of the title structure viewed down the c axis. N—H···O and C—H···F hydrogen bondings (dashed lines) link the molecules, forming $R^2_3(19)$ graph-set motifs along the b axis of the unit cell. H atoms not involved in hydrogen bonds have been omitted for clarity.

4,6-Bis(4-fluorophenyl)-2-phenyl-1*H*-indazol-3(2*H*)-one

Crystal data

$C_{25}H_{16}F_2N_2O$
 $M_r = 398.40$
Orthorhombic, $Pbcn$
Hall symbol: -P 2n 2ab
 $a = 15.2947 (4)$ Å
 $b = 11.6259 (2)$ Å
 $c = 20.9388 (5)$ Å
 $V = 3723.23 (15)$ Å³
 $Z = 8$

$F(000) = 1648$
 $D_x = 1.421 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 11209 reflections
 $\theta = 5.2\text{--}37.5^\circ$
 $\mu = 0.10 \text{ mm}^{-1}$
 $T = 123$ K
Prism, colourless
 $0.49 \times 0.38 \times 0.23$ mm

Data collection

Oxford Diffraction Xcalibur Ruby Gemini diffractometer
Radiation source: Enhance (Mo) X-ray Source
Graphite monochromator
Detector resolution: 10.5081 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(CrysAlis PRO; Oxford Diffraction, 2007)
 $T_{\min} = 0.895$, $T_{\max} = 0.977$

19870 measured reflections
3827 independent reflections
3416 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$
 $\theta_{\max} = 26.5^\circ$, $\theta_{\min} = 5.2^\circ$
 $h = -19 \rightarrow 19$
 $k = -14 \rightarrow 14$
 $l = -26 \rightarrow 26$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.135$
 $S = 1.09$

3827 reflections
275 parameters
1 restraint
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

$$w = 1/[\sigma^2(F_o^2) + (0.0462P)^2 + 4.0317P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

Hydrogen site location: inferred from neighbouring sites

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.65 \text{ e \AA}^{-3}$$

H atoms treated by a mixture of independent and constrained refinement

$$\Delta\rho_{\min} = -0.35 \text{ e \AA}^{-3}$$

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors.

Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating - R -factor-obs etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.60370 (13)	-0.57691 (11)	0.22016 (9)	0.0656 (6)
F2	0.48986 (10)	0.29173 (14)	0.53611 (7)	0.0525 (5)
O1	0.73203 (11)	-0.14990 (12)	0.11749 (7)	0.0351 (5)
N1	0.72290 (11)	0.04878 (13)	0.11220 (7)	0.0239 (5)
N2	0.68782 (11)	0.13815 (13)	0.14893 (8)	0.0241 (5)
C1	0.70924 (15)	-0.05733 (16)	0.14107 (9)	0.0269 (6)
C2	0.66455 (14)	-0.03003 (16)	0.20027 (9)	0.0272 (6)
C3	0.63295 (16)	-0.10034 (17)	0.25063 (10)	0.0323 (6)
C4	0.60213 (13)	-0.04532 (16)	0.30476 (9)	0.0246 (5)
C5	0.60116 (12)	0.07752 (16)	0.31012 (9)	0.0230 (5)
C6	0.62766 (13)	0.14448 (16)	0.25935 (9)	0.0248 (6)
C7	0.65858 (13)	0.08914 (16)	0.20431 (9)	0.0241 (5)
C8	0.62753 (14)	-0.22733 (16)	0.24316 (9)	0.0257 (6)
C9	0.58886 (14)	-0.27250 (18)	0.18834 (10)	0.0291 (6)
C10	0.57996 (15)	-0.3901 (2)	0.18045 (11)	0.0355 (7)
C11	0.61074 (16)	-0.46066 (17)	0.22766 (12)	0.0382 (7)
C12	0.64919 (16)	-0.42035 (19)	0.28203 (12)	0.0391 (7)
C13	0.65705 (15)	-0.30264 (18)	0.28970 (10)	0.0315 (6)
C14	0.57027 (12)	0.13239 (17)	0.37010 (9)	0.0246 (6)
C15	0.58997 (14)	0.08541 (18)	0.42973 (10)	0.0302 (6)
C16	0.56348 (15)	0.1384 (2)	0.48586 (10)	0.0365 (7)
C17	0.51680 (14)	0.2389 (2)	0.48129 (10)	0.0355 (7)
C18	0.49559 (14)	0.28874 (19)	0.42389 (11)	0.0338 (6)
C19	0.52226 (13)	0.23481 (18)	0.36821 (10)	0.0280 (6)
C20	0.74707 (13)	0.07083 (15)	0.04813 (9)	0.0216 (5)
C21	0.71378 (14)	0.16740 (16)	0.01763 (9)	0.0257 (6)
C22	0.73814 (15)	0.19020 (17)	-0.04494 (9)	0.0303 (6)
C23	0.79396 (15)	0.11777 (17)	-0.07719 (10)	0.0303 (6)
C24	0.82668 (15)	0.02163 (18)	-0.04644 (10)	0.0320 (6)

C25	0.80400 (13)	-0.00225 (17)	0.01629 (10)	0.0279 (6)
HN2	0.7174 (14)	0.2009 (14)	0.1477 (12)	0.038 (7)*
H4	0.58140	-0.08940	0.33860	0.0300*
H6	0.62510	0.22430	0.26150	0.0300*
H9	0.56880	-0.22300	0.15670	0.0350*
H10	0.55380	-0.42030	0.14400	0.0430*
H12	0.66960	-0.47070	0.31310	0.0470*
H13	0.68250	-0.27360	0.32670	0.0380*
H15	0.62150	0.01710	0.43190	0.0360*
H16	0.57700	0.10670	0.52540	0.0440*
H18	0.46410	0.35710	0.42240	0.0400*
H19	0.50800	0.26720	0.32890	0.0340*
H21	0.67550	0.21630	0.03890	0.0310*
H22	0.71650	0.25530	-0.06540	0.0360*
H23	0.80960	0.13330	-0.11920	0.0360*
H24	0.86430	-0.02760	-0.06810	0.0380*
H25	0.82670	-0.06660	0.03680	0.0330*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0954 (13)	0.0186 (7)	0.0829 (12)	-0.0145 (7)	0.0149 (10)	-0.0064 (7)
F2	0.0579 (9)	0.0652 (10)	0.0345 (7)	0.0125 (8)	0.0096 (7)	-0.0210 (7)
O1	0.0646 (11)	0.0155 (7)	0.0251 (7)	0.0055 (7)	0.0088 (7)	0.0007 (6)
N1	0.0364 (9)	0.0141 (7)	0.0213 (8)	0.0030 (7)	0.0009 (7)	-0.0001 (6)
N2	0.0366 (9)	0.0142 (7)	0.0216 (8)	0.0004 (7)	0.0023 (7)	-0.0007 (6)
C1	0.0440 (12)	0.0172 (9)	0.0195 (9)	0.0009 (8)	0.0002 (8)	0.0024 (7)
C2	0.0421 (11)	0.0179 (9)	0.0216 (9)	0.0008 (8)	0.0008 (8)	-0.0011 (7)
C3	0.0489 (13)	0.0216 (10)	0.0265 (10)	-0.0039 (9)	0.0052 (9)	0.0002 (8)
C4	0.0311 (10)	0.0216 (9)	0.0210 (9)	-0.0026 (8)	0.0005 (8)	0.0014 (7)
C5	0.0253 (9)	0.0225 (9)	0.0213 (9)	0.0023 (7)	-0.0030 (8)	-0.0027 (7)
C6	0.0344 (11)	0.0149 (8)	0.0250 (10)	0.0026 (7)	-0.0019 (8)	-0.0011 (7)
C7	0.0326 (10)	0.0177 (9)	0.0220 (9)	-0.0008 (8)	-0.0024 (8)	0.0011 (7)
C8	0.0365 (11)	0.0184 (9)	0.0222 (9)	-0.0025 (8)	0.0058 (8)	-0.0002 (7)
C9	0.0345 (11)	0.0292 (10)	0.0237 (10)	0.0015 (9)	0.0011 (9)	0.0034 (8)
C10	0.0384 (12)	0.0360 (12)	0.0321 (11)	-0.0130 (10)	0.0035 (9)	-0.0104 (9)
C11	0.0496 (14)	0.0153 (9)	0.0496 (14)	-0.0070 (9)	0.0130 (11)	-0.0019 (9)
C12	0.0503 (14)	0.0262 (11)	0.0407 (13)	0.0018 (10)	0.0019 (11)	0.0134 (10)
C13	0.0417 (12)	0.0291 (11)	0.0238 (10)	-0.0061 (9)	-0.0023 (9)	0.0046 (8)
C14	0.0247 (9)	0.0246 (10)	0.0244 (10)	-0.0009 (7)	-0.0005 (8)	-0.0038 (8)
C15	0.0351 (11)	0.0303 (10)	0.0253 (10)	0.0051 (9)	0.0007 (9)	-0.0019 (8)
C16	0.0419 (13)	0.0449 (13)	0.0228 (10)	0.0039 (10)	0.0002 (9)	-0.0021 (9)
C17	0.0329 (11)	0.0451 (13)	0.0284 (11)	0.0006 (10)	0.0059 (9)	-0.0150 (10)
C18	0.0283 (10)	0.0330 (11)	0.0400 (12)	0.0068 (9)	0.0025 (9)	-0.0083 (10)
C19	0.0261 (10)	0.0292 (10)	0.0287 (10)	0.0027 (8)	-0.0029 (8)	-0.0039 (8)
C20	0.0276 (9)	0.0180 (9)	0.0191 (9)	-0.0041 (7)	-0.0018 (7)	0.0010 (7)
C21	0.0368 (11)	0.0191 (9)	0.0213 (9)	0.0043 (8)	-0.0013 (8)	-0.0022 (7)
C22	0.0479 (13)	0.0206 (9)	0.0223 (10)	0.0003 (9)	-0.0040 (9)	0.0026 (8)

C23	0.0452 (12)	0.0243 (10)	0.0215 (9)	-0.0072 (9)	0.0054 (9)	0.0021 (8)
C24	0.0370 (11)	0.0273 (11)	0.0318 (11)	0.0024 (9)	0.0095 (9)	-0.0007 (9)
C25	0.0317 (10)	0.0221 (9)	0.0298 (10)	0.0028 (8)	0.0011 (8)	0.0037 (8)

Geometric parameters (\AA , $^{\circ}$)

F1—C11	1.365 (2)	C15—C16	1.387 (3)
F2—C17	1.366 (3)	C16—C17	1.373 (3)
O1—C1	1.234 (2)	C17—C18	1.373 (3)
N1—N2	1.400 (2)	C18—C19	1.385 (3)
N1—C1	1.390 (2)	C20—C25	1.387 (3)
N1—C20	1.415 (2)	C20—C21	1.388 (3)
N2—C7	1.367 (3)	C21—C22	1.388 (3)
N2—HN2	0.859 (18)	C22—C23	1.376 (3)
C1—C2	1.451 (3)	C23—C24	1.384 (3)
C2—C3	1.419 (3)	C24—C25	1.387 (3)
C2—C7	1.391 (3)	C4—H4	0.9300
C3—C4	1.384 (3)	C6—H6	0.9300
C3—C8	1.487 (3)	C9—H9	0.9300
C4—C5	1.433 (3)	C10—H10	0.9300
C5—C14	1.486 (3)	C12—H12	0.9300
C5—C6	1.379 (3)	C13—H13	0.9300
C6—C7	1.402 (3)	C15—H15	0.9300
C8—C13	1.386 (3)	C16—H16	0.9300
C8—C9	1.394 (3)	C18—H18	0.9300
C9—C10	1.384 (3)	C19—H19	0.9300
C10—C11	1.368 (3)	C21—H21	0.9300
C11—C12	1.364 (3)	C22—H22	0.9300
C12—C13	1.383 (3)	C23—H23	0.9300
C14—C15	1.396 (3)	C24—H24	0.9300
C14—C19	1.400 (3)	C25—H25	0.9300
N2—N1—C1	111.25 (15)	C17—C18—C19	118.4 (2)
N2—N1—C20	119.12 (14)	C14—C19—C18	121.05 (19)
C1—N1—C20	127.78 (15)	N1—C20—C21	119.13 (17)
N1—N2—C7	106.38 (14)	C21—C20—C25	120.29 (18)
C7—N2—HN2	123.5 (17)	N1—C20—C25	120.58 (17)
N1—N2—HN2	114.3 (15)	C20—C21—C22	119.36 (18)
N1—C1—C2	104.38 (15)	C21—C22—C23	120.86 (19)
O1—C1—C2	131.73 (18)	C22—C23—C24	119.36 (19)
O1—C1—N1	123.89 (18)	C23—C24—C25	120.8 (2)
C1—C2—C3	132.02 (18)	C20—C25—C24	119.32 (18)
C1—C2—C7	107.51 (16)	C3—C4—H4	119.00
C3—C2—C7	120.41 (18)	C5—C4—H4	119.00
C2—C3—C4	117.28 (18)	C5—C6—H6	121.00
C4—C3—C8	121.73 (18)	C7—C6—H6	121.00
C2—C3—C8	120.85 (18)	C8—C9—H9	120.00
C3—C4—C5	121.89 (18)	C10—C9—H9	120.00

C4—C5—C6	119.97 (17)	C9—C10—H10	121.00
C4—C5—C14	119.84 (17)	C11—C10—H10	121.00
C6—C5—C14	120.19 (17)	C11—C12—H12	121.00
C5—C6—C7	118.29 (17)	C13—C12—H12	121.00
C2—C7—C6	121.95 (17)	C8—C13—H13	120.00
N2—C7—C6	128.04 (17)	C12—C13—H13	119.00
N2—C7—C2	110.00 (16)	C14—C15—H15	119.00
C3—C8—C13	122.36 (18)	C16—C15—H15	119.00
C9—C8—C13	118.64 (18)	C15—C16—H16	121.00
C3—C8—C9	118.97 (18)	C17—C16—H16	121.00
C8—C9—C10	120.81 (19)	C17—C18—H18	121.00
C9—C10—C11	118.2 (2)	C19—C18—H18	121.00
F1—C11—C10	118.9 (2)	C14—C19—H19	119.00
C10—C11—C12	123.0 (2)	C18—C19—H19	119.00
F1—C11—C12	118.0 (2)	C20—C21—H21	120.00
C11—C12—C13	118.3 (2)	C22—C21—H21	120.00
C8—C13—C12	121.0 (2)	C21—C22—H22	120.00
C15—C14—C19	118.13 (18)	C23—C22—H22	119.00
C5—C14—C15	121.30 (18)	C22—C23—H23	120.00
C5—C14—C19	120.55 (17)	C24—C23—H23	120.00
C14—C15—C16	121.40 (19)	C23—C24—H24	120.00
C15—C16—C17	118.1 (2)	C25—C24—H24	120.00
F2—C17—C18	118.3 (2)	C20—C25—H25	120.00
C16—C17—C18	122.9 (2)	C24—C25—H25	120.00
F2—C17—C16	118.77 (19)		
C1—N1—N2—C7	5.3 (2)	C6—C5—C14—C19	35.8 (3)
C20—N1—N2—C7	171.01 (17)	C4—C5—C14—C19	-143.24 (19)
N2—N1—C1—O1	178.1 (2)	C4—C5—C6—C7	-2.7 (3)
C20—N1—C1—O1	14.0 (3)	C4—C5—C14—C15	38.4 (3)
N2—N1—C1—C2	-1.4 (2)	C5—C6—C7—N2	179.9 (2)
C20—N1—C1—C2	-165.57 (19)	C5—C6—C7—C2	-1.2 (3)
C1—N1—C20—C25	-35.8 (3)	C3—C8—C9—C10	-177.9 (2)
N2—N1—C20—C21	-18.2 (3)	C3—C8—C13—C12	178.4 (2)
C1—N1—C20—C21	144.9 (2)	C9—C8—C13—C12	0.5 (3)
N2—N1—C20—C25	161.15 (18)	C13—C8—C9—C10	0.1 (3)
N1—N2—C7—C6	171.78 (19)	C8—C9—C10—C11	-0.5 (3)
N1—N2—C7—C2	-7.2 (2)	C9—C10—C11—F1	-179.0 (2)
N1—C1—C2—C3	-180.0 (2)	C9—C10—C11—C12	0.4 (4)
N1—C1—C2—C7	-3.0 (2)	C10—C11—C12—C13	0.2 (4)
O1—C1—C2—C3	0.5 (4)	F1—C11—C12—C13	179.6 (2)
O1—C1—C2—C7	177.5 (2)	C11—C12—C13—C8	-0.7 (4)
C1—C2—C7—C6	-172.62 (19)	C19—C14—C15—C16	-0.4 (3)
C1—C2—C7—N2	6.4 (2)	C5—C14—C19—C18	-177.86 (19)
C1—C2—C3—C4	172.4 (2)	C5—C14—C15—C16	178.00 (19)
C3—C2—C7—N2	-176.14 (19)	C15—C14—C19—C18	0.5 (3)
C3—C2—C7—C6	4.8 (3)	C14—C15—C16—C17	0.2 (3)
C7—C2—C3—C8	171.4 (2)	C15—C16—C17—F2	179.7 (2)

C1—C2—C3—C8	−11.9 (4)	C15—C16—C17—C18	−0.1 (3)
C7—C2—C3—C4	−4.3 (3)	C16—C17—C18—C19	0.3 (3)
C4—C3—C8—C9	127.1 (2)	F2—C17—C18—C19	−179.50 (19)
C2—C3—C4—C5	0.5 (3)	C17—C18—C19—C14	−0.5 (3)
C4—C3—C8—C13	−50.8 (3)	N1—C20—C21—C22	179.20 (18)
C2—C3—C8—C13	133.7 (2)	C25—C20—C21—C22	−0.1 (3)
C2—C3—C8—C9	−48.4 (3)	N1—C20—C25—C24	−179.86 (18)
C8—C3—C4—C5	−175.18 (19)	C21—C20—C25—C24	−0.6 (3)
C3—C4—C5—C6	3.0 (3)	C20—C21—C22—C23	0.7 (3)
C3—C4—C5—C14	−177.90 (19)	C21—C22—C23—C24	−0.6 (3)
C6—C5—C14—C15	−142.5 (2)	C22—C23—C24—C25	−0.1 (3)
C14—C5—C6—C7	178.28 (17)	C23—C24—C25—C20	0.7 (3)

Hydrogen-bond geometry (Å, °)

Cg5 is the centroid of the C20—C25 phenyl ring.

D—H···A	D—H	H···A	D···A	D—H···A
N2—HN2···O1 ⁱ	0.86 (2)	2.00 (2)	2.830 (2)	162 (2)
C6—H6···F1 ⁱⁱ	0.93	2.49	3.362 (2)	156
C21—H21···N2	0.93	2.48	2.799 (3)	100
C25—H25···O1	0.93	2.43	2.941 (3)	115
C15—H15···Cg5 ⁱⁱⁱ	0.93	2.85	3.656 (2)	145

Symmetry codes: (i) $-x+3/2, y+1/2, z$; (ii) $x, y+1, z$; (iii) $x, -y, z+1/2$.