

# Di- $\mu$ -benzoato- $\kappa^3$ O,O':O; $\kappa^3$ O:O,O'-bis[(benzoato- $\kappa^2$ O,O')(1,10-phenanthroline- $\kappa^2$ N,N')lead(II)]

Hong-Jin Li,<sup>a</sup> Zhu-Qing Gao<sup>a\*</sup> and Jin-Zhong Gu<sup>b</sup>

<sup>a</sup>School of Chemistry and Biology Engineering, Taiyuan University of Science and Technology, Taiyuan 030021, People's Republic of China, and <sup>b</sup>College of Chemistry and Chemical Engineering, Lanzhou University, Lanzhou 730000, People's Republic of China

Correspondence e-mail: zqgao2008@163.com


Received 4 May 2011; accepted 15 May 2011

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(C-C) = 0.011$  Å;  $R$  factor = 0.034;  $wR$  factor = 0.069; data-to-parameter ratio = 13.6.

In the centrosymmetric dinuclear title compound,  $[Pb_2(C_7H_5O_2)_4(C_{12}H_8N_2)_2]$ , two  $Pb^{2+}$  ions are connected by two tridentate bridging benzoate anions. The  $Pb^{2+}$  ion is seven-coordinated by five O atoms from three benzoate anions and two N atoms from the 1,10-phenanthroline ligands. The benzoate anions adopt two different coordination modes, one bidentate-chelating and one tridentate bridging-chelating. The three-dimensional supramolecular framework is achieved by intermolecular  $\pi-\pi$  stacking interactions, with a shortest centroid–centroid distance of 3.617 (4) Å.

## Related literature

For bond lengths and angles in other lead(II) compounds, see: Fan *et al.* (2006); Hu *et al.* (2011).



## Experimental

### Crystal data

$[Pb_2(C_7H_5O_2)_4(C_{12}H_8N_2)_2]$	$\gamma = 71.601 (3)^\circ$
$M_r = 1259.23$	$V = 1106.6 (6) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 1$
$a = 9.011 (3)$ Å	Mo $K\alpha$ radiation
$b = 10.923 (3)$ Å	$\mu = 7.66 \text{ mm}^{-1}$
$c = 11.920 (4)$ Å	$T = 293$ K
$\alpha = 83.760 (3)^\circ$	$0.28 \times 0.26 \times 0.24$ mm
$\beta = 87.626 (3)^\circ$	

### Data collection

Bruker APEXII CCD diffractometer	7969 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2004)	4059 independent reflections
$T_{\min} = 0.223$ , $T_{\max} = 0.261$	3296 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.041$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$	298 parameters
$wR(F^2) = 0.069$	H-atom parameters constrained
$S = 0.96$	$\Delta\rho_{\max} = 1.95 \text{ e \AA}^{-3}$
4059 reflections	$\Delta\rho_{\min} = -1.35 \text{ e \AA}^{-3}$

**Table 1**  
Selected bond lengths (Å).

Pb1—O4	2.394 (4)	Pb1—O2	2.723 (5)
Pb1—N1	2.578 (5)	Pb1—O3	2.788 (5)
Pb1—O1	2.584 (4)	Pb1—O3 <sup>i</sup>	2.924 (5)
Pb1—N2	2.703 (5)		

Symmetry code: (i)  $-x + 1, -y + 2, -z + 1$ .

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2005); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2486).

## References

- Brandenburg, K. & Putz, H. (2005). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2004). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fan, S. R. & Zhu, L. G. (2006). *Inorg. Chem. Commun.* **45**, 7935–7942.
- Hu, R. R., Cai, H. & Luo, J. H. (2011). *Inorg. Chem. Commun.* **14**, 433–436.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

# supporting information

*Acta Cryst.* (2011). E67, m778 [doi:10.1107/S160053681101840X]

## Di- $\mu$ -benzoato- $\kappa^3O,O':O;\kappa^3O:O,O'$ -bis[(benzoato- $\kappa^2O,O'$ )(1,10-phenanthroline- $\kappa^2N,N'$ )lead(II)]

Hong-Jin Li, Zhu-Qing Gao and Jin-Zhong Gu

### S1. Comment

Lead(II) compounds have been increasingly studied owing to their interesting physical and chemical properties (Fan *et al.*, 2006; Hu *et al.*, 2011). In order to extend our investigations in this field, we crystallised the lead(II) title compound [ $Pb_2(C_7H_5O_2)_4(C_{12}H_8N_2)_2$ ], and report its structure here.

The asymmetric unit of the title complex (Fig. 1) contains one  $Pb^{2+}$  ion, two benzoate anions, and one 1,10-phenanthroline ligand. The  $Pb^{2+}$  ion is seven-coordinated by five O atoms from three benzoate ligands and by two N atoms from 1,10-phenanthroline. The coordination environment around the  $Pb^{2+}$  ion may be described as a distorted mono-capped trigonal prism. Two adjacent  $Pb^{II}$  complexes are connected by two bridging benzoate anions to generate a centrosymmetric dinuclear unit. The benzoate anions adopt two kinds of coordination modes, *viz.* a bidentate chelating and a tridentate bridging-chelating mode.

The Pb—N and Pb—O bond lengths range between 2.578 (5)–2.703 (5) Å and 2.394 (4)–2.924 (5) Å, respectively. These values are in good agreement with those reported for other Pb(II)—O and Pb(II)—N donor complexes (Fan *et al.*, 2006; Hu *et al.*, 2011).


In the crystal structure,  $\pi$ — $\pi$  stacking interactions between adjacent 1,10-phenanthroline ligands [centroid—centroid distance = 3.617 (4) Å] are observed. Furthermore, adjacent benzene rings from benzoate anions are also involved in  $\pi$ — $\pi$  stacking interactions [centroid—centroid distance = 4.083 (3) Å].  $\pi$ — $\pi$  stacking interactions between adjacent 1,10-phenanthroline ligands and benzene rings from benzoate anions [centroid—centroid distance = 3.945 (4) Å] are also observed. These interactions of the discrete neutral molecules lead to a three-dimensional supramolecular framework (Fig. 2).

### S2. Experimental


A mixture of  $Pb(CH_3COO)_2 \cdot 3H_2O$  (0.20 g, 0.54 mmol), benzoic acid (0.12 g, 1.0 mmol), 1,10-phenanthroline (0.11 g, 0.54 mmol), NaOH (0.04 g, 1.0 mmol), and water (10 ml) was stirred at room temperature for 15 min, and then sealed in a 25 ml Teflon-lined, stainless-steel Parr autoclave. The autoclave was heated at 433 K for 3 d. Upon cooling, the solution contained single crystals of the title complex in *ca* 80% yield. Anal./calc. for  $C_{26}H_{18}N_2O_4Pb$ : C, 49.60; H, 2.88; N, 4.45; found: C, 49.43; H, 3.07; N, 4.13.

### S3. Refinement

The carbon-bound H atoms were placed in geometrically idealized positions, with C—H = 0.93 Å, and constrained to ride on their respective parent atoms, with  $U_{iso}(H) = 1.2 U_{eq}(C)$ . The highest peak and the deepest hole in the final difference map are 0.98 Å and 0.89 Å, respectively, from Pb1.

**Figure 1**


The asymmetric unit in the structure of the title complex, showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 30% probability level.

**Figure 2**

View approximately along the  $a$  axis, showing the three-dimensional framework structure in the title complex.

### Di- $\mu$ -benzoato- $\kappa^3O,O':O;\kappa^3O;O,O'$ - bis[(benzoato- $\kappa^2O,O'$ )(1,10-phenanthroline- $\kappa^2N,N'$ )lead(II)]

#### Crystal data



$M_r = 1259.23$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a = 9.011 (3)$  Å

$b = 10.923 (3)$  Å

$c = 11.920 (4)$  Å

$\alpha = 83.760 (3)^\circ$

$\beta = 87.626 (3)^\circ$

$\gamma = 71.601 (3)^\circ$

$V = 1106.6 (6)$  Å<sup>3</sup>

$Z = 1$

$F(000) = 604$

$D_x = 1.889 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 3450 reflections

$\theta = 2.4\text{--}24.1^\circ$

$\mu = 7.66 \text{ mm}^{-1}$

$T = 293$  K

Block, colorless

$0.28 \times 0.26 \times 0.24$  mm

#### Data collection

Bruker APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2004)

$T_{\min} = 0.223$ ,  $T_{\max} = 0.261$

7969 measured reflections

4059 independent reflections

3296 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.041$

$\theta_{\max} = 25.5^\circ$ ,  $\theta_{\min} = 2.4^\circ$

$h = -10 \rightarrow 10$

$k = -12 \rightarrow 13$

$l = -14 \rightarrow 14$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.034$  $wR(F^2) = 0.069$  $S = 0.96$ 

4059 reflections

298 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0271P)^2]$   
where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.001$  $\Delta\rho_{\max} = 1.95 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\min} = -1.35 \text{ e } \text{\AA}^{-3}$ *Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Pb1	0.24985 (3)	0.97230 (2)	0.484550 (19)	0.03770 (9)
C3	0.1241 (8)	0.5411 (7)	0.4132 (6)	0.060 (2)
H3	0.0979	0.4679	0.4009	0.072*
C2	0.0639 (8)	0.6531 (8)	0.3469 (7)	0.064 (2)
H2	-0.0042	0.6577	0.2890	0.077*
C1	0.1049 (8)	0.7616 (7)	0.3660 (6)	0.0556 (18)
H1	0.0621	0.8384	0.3202	0.067*
N1	0.2020 (5)	0.7600 (5)	0.4465 (4)	0.0415 (12)
C5	0.2638 (7)	0.6483 (6)	0.5128 (5)	0.0383 (14)
C9	0.3737 (7)	0.6449 (6)	0.5979 (5)	0.0392 (14)
N2	0.4171 (6)	0.7506 (5)	0.6046 (4)	0.0414 (12)
C12	0.5255 (8)	0.7451 (6)	0.6789 (6)	0.0515 (17)
H12	0.5580	0.8175	0.6821	0.062*
C11	0.5919 (8)	0.6347 (7)	0.7520 (6)	0.062 (2)
H11	0.6669	0.6340	0.8036	0.074*
C10	0.5474 (8)	0.5294 (7)	0.7478 (6)	0.0603 (19)
H10	0.5890	0.4562	0.7981	0.072*
C8	0.4388 (8)	0.5296 (6)	0.6680 (6)	0.0484 (17)
C7	0.3938 (9)	0.4183 (6)	0.6538 (7)	0.0591 (19)
H7	0.4332	0.3431	0.7023	0.071*
C6	0.2968 (9)	0.4201 (7)	0.5725 (7)	0.060 (2)
H6	0.2746	0.3444	0.5624	0.072*
C4	0.2253 (7)	0.5353 (6)	0.5000 (5)	0.0460 (16)
O1	0.0004 (5)	1.0690 (4)	0.3639 (4)	0.0515 (11)
O2	0.1272 (6)	1.2117 (5)	0.3714 (4)	0.0702 (15)

C13	0.0280 (8)	1.1731 (6)	0.3290 (6)	0.0466 (16)
C14	-0.0619 (7)	1.2544 (6)	0.2281 (5)	0.0417 (15)
C19	-0.1491 (8)	1.2081 (8)	0.1642 (6)	0.068 (2)
H19	-0.1557	1.1250	0.1825	0.081*
C15	-0.0538 (8)	1.3785 (7)	0.1983 (6)	0.0580 (19)
H15	0.0068	1.4112	0.2406	0.070*
C18	-0.2279 (11)	1.2843 (12)	0.0720 (8)	0.104 (3)
H18	-0.2879	1.2521	0.0287	0.125*
C17	-0.2193 (12)	1.4049 (12)	0.0435 (8)	0.108 (4)
H17	-0.2718	1.4546	-0.0197	0.130*
C16	-0.1348 (11)	1.4533 (9)	0.1065 (7)	0.086 (3)
H16	-0.1312	1.5372	0.0880	0.103*
O3	0.5456 (6)	0.9341 (5)	0.3871 (4)	0.0672 (14)
O4	0.3436 (6)	0.9290 (5)	0.2975 (4)	0.0689 (14)
C20	0.4843 (8)	0.9206 (6)	0.3015 (6)	0.0436 (15)
C21	0.5768 (7)	0.8906 (5)	0.1958 (5)	0.0410 (15)
C26	0.5161 (9)	0.8558 (7)	0.1053 (6)	0.0612 (19)
H26	0.4149	0.8506	0.1100	0.073*
C25	0.5999 (13)	0.8290 (8)	0.0091 (7)	0.089 (3)
H25	0.5563	0.8068	-0.0519	0.106*
C22	0.7265 (8)	0.8974 (7)	0.1879 (6)	0.067 (2)
H22	0.7698	0.9208	0.2485	0.080*
C23	0.8124 (11)	0.8697 (9)	0.0907 (10)	0.097 (3)
H23	0.9136	0.8749	0.0856	0.116*
C24	0.7511 (14)	0.8349 (9)	0.0027 (8)	0.100 (4)
H24	0.8105	0.8150	-0.0623	0.120*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Pb1	0.04097 (14)	0.03437 (14)	0.04007 (15)	-0.01488 (10)	0.00285 (9)	-0.00557 (10)
C3	0.060 (5)	0.050 (5)	0.081 (6)	-0.027 (4)	0.015 (4)	-0.028 (4)
C2	0.059 (5)	0.068 (5)	0.074 (5)	-0.028 (4)	-0.009 (4)	-0.023 (4)
C1	0.058 (4)	0.053 (5)	0.059 (5)	-0.017 (4)	-0.007 (4)	-0.012 (4)
N1	0.041 (3)	0.038 (3)	0.049 (3)	-0.015 (2)	0.002 (2)	-0.012 (3)
C5	0.042 (3)	0.034 (3)	0.044 (4)	-0.017 (3)	0.015 (3)	-0.014 (3)
C9	0.041 (3)	0.036 (4)	0.042 (4)	-0.015 (3)	0.014 (3)	-0.009 (3)
N2	0.053 (3)	0.033 (3)	0.039 (3)	-0.015 (3)	0.008 (2)	-0.007 (2)
C12	0.058 (4)	0.044 (4)	0.055 (4)	-0.018 (3)	-0.002 (3)	-0.011 (3)
C11	0.064 (5)	0.055 (5)	0.060 (5)	-0.010 (4)	-0.011 (4)	-0.001 (4)
C10	0.078 (5)	0.045 (4)	0.048 (4)	-0.008 (4)	-0.007 (4)	0.007 (3)
C8	0.057 (4)	0.036 (4)	0.050 (4)	-0.013 (3)	0.015 (3)	-0.005 (3)
C7	0.074 (5)	0.034 (4)	0.067 (5)	-0.015 (4)	0.014 (4)	-0.003 (4)
C6	0.071 (5)	0.037 (4)	0.078 (6)	-0.023 (4)	0.026 (4)	-0.021 (4)
C4	0.047 (4)	0.041 (4)	0.059 (4)	-0.023 (3)	0.014 (3)	-0.019 (3)
O1	0.057 (3)	0.040 (3)	0.058 (3)	-0.017 (2)	0.003 (2)	-0.001 (2)
O2	0.072 (3)	0.063 (3)	0.082 (4)	-0.034 (3)	-0.032 (3)	0.012 (3)
C13	0.047 (4)	0.037 (4)	0.054 (4)	-0.011 (3)	0.007 (3)	-0.008 (3)

C14	0.040 (4)	0.050 (4)	0.038 (4)	-0.018 (3)	0.008 (3)	-0.006 (3)
C19	0.073 (5)	0.091 (6)	0.053 (5)	-0.046 (5)	0.000 (4)	-0.005 (4)
C15	0.063 (5)	0.057 (5)	0.054 (4)	-0.021 (4)	0.004 (4)	-0.001 (4)
C18	0.122 (8)	0.155 (11)	0.062 (6)	-0.081 (8)	-0.024 (5)	0.000 (7)
C17	0.123 (9)	0.143 (10)	0.057 (6)	-0.050 (8)	-0.034 (6)	0.034 (7)
C16	0.094 (7)	0.084 (6)	0.068 (6)	-0.023 (5)	0.004 (5)	0.024 (5)
O3	0.093 (4)	0.074 (4)	0.046 (3)	-0.038 (3)	0.002 (3)	-0.019 (3)
O4	0.069 (3)	0.093 (4)	0.050 (3)	-0.032 (3)	0.018 (2)	-0.017 (3)
C20	0.058 (4)	0.027 (3)	0.048 (4)	-0.017 (3)	0.004 (3)	-0.005 (3)
C21	0.051 (4)	0.030 (3)	0.042 (4)	-0.013 (3)	0.012 (3)	-0.006 (3)
C26	0.081 (5)	0.061 (5)	0.047 (4)	-0.029 (4)	0.014 (4)	-0.013 (4)
C25	0.129 (9)	0.086 (7)	0.056 (5)	-0.038 (6)	0.025 (5)	-0.027 (5)
C22	0.060 (5)	0.075 (5)	0.064 (5)	-0.021 (4)	0.017 (4)	-0.012 (4)
C23	0.073 (6)	0.094 (7)	0.114 (8)	-0.018 (5)	0.044 (6)	-0.010 (7)
C24	0.137 (10)	0.069 (6)	0.074 (7)	-0.007 (6)	0.060 (7)	-0.017 (5)

Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )

Pb1—O4	2.394 (4)	O1—C13	1.263 (7)
Pb1—N1	2.578 (5)	O1—Pb1 <sup>ii</sup>	2.946 (4)
Pb1—O1	2.584 (4)	O2—C13	1.247 (8)
Pb1—N2	2.703 (5)	C13—C14	1.512 (9)
Pb1—O2	2.723 (5)	C13—Pb1 <sup>ii</sup>	3.874 (6)
Pb1—O3	2.788 (5)	C14—C19	1.356 (9)
Pb1—O3 <sup>i</sup>	2.924 (5)	C14—C15	1.387 (9)
C3—C2	1.350 (10)	C19—C18	1.377 (11)
C3—C4	1.390 (9)	C19—H19	0.9300
C3—H3	0.9300	C15—C16	1.376 (10)
C2—C1	1.390 (9)	C15—H15	0.9300
C2—H2	0.9300	C18—C17	1.350 (13)
C1—N1	1.320 (8)	C18—H18	0.9300
C1—H1	0.9300	C17—C16	1.345 (12)
N1—C5	1.350 (7)	C17—H17	0.9300
C5—C4	1.408 (8)	C16—H16	0.9300
C5—C9	1.436 (8)	O3—C20	1.224 (7)
C9—N2	1.342 (7)	O3—Pb1 <sup>i</sup>	2.923 (5)
C9—C8	1.404 (8)	O4—C20	1.244 (7)
N2—C12	1.329 (8)	C20—C21	1.490 (8)
C12—C11	1.389 (9)	C21—C26	1.370 (9)
C12—H12	0.9300	C21—C22	1.374 (9)
C11—C10	1.339 (10)	C26—C25	1.353 (10)
C11—H11	0.9300	C26—H26	0.9300
C10—C8	1.392 (9)	C25—C24	1.383 (13)
C10—H10	0.9300	C25—H25	0.9300
C8—C7	1.425 (9)	C22—C23	1.374 (11)
C7—C6	1.326 (10)	C22—H22	0.9300
C7—H7	0.9300	C23—C24	1.348 (13)
C6—C4	1.429 (9)	C23—H23	0.9300

C6—H6	0.9300	C24—H24	0.9300
O4—Pb1—N1	73.03 (16)	C6—C7—H7	119.4
O4—Pb1—O1	76.92 (16)	C8—C7—H7	119.4
N1—Pb1—O1	80.92 (15)	C7—C6—C4	121.7 (6)
O4—Pb1—N2	100.85 (16)	C7—C6—H6	119.1
N1—Pb1—N2	62.22 (16)	C4—C6—H6	119.1
O1—Pb1—N2	141.48 (15)	C3—C4—C5	117.5 (6)
O4—Pb1—O2	79.49 (17)	C3—C4—C6	123.6 (6)
N1—Pb1—O2	127.45 (15)	C5—C4—C6	118.9 (6)
O1—Pb1—O2	49.27 (13)	C13—O1—Pb1	95.5 (4)
N2—Pb1—O2	169.20 (13)	C13—O1—Pb1 <sup>ii</sup>	129.6 (4)
O4—Pb1—O3	48.70 (15)	Pb1—O1—Pb1 <sup>ii</sup>	103.74 (14)
N1—Pb1—O3	100.12 (14)	C13—O2—Pb1	89.4 (4)
O1—Pb1—O3	120.86 (14)	O2—C13—O1	124.0 (6)
N2—Pb1—O3	78.88 (14)	O2—C13—C14	117.9 (6)
O2—Pb1—O3	93.68 (15)	O1—C13—C14	118.1 (6)
O4—Pb1—O3 <sup>i</sup>	113.45 (15)	C19—C14—C15	118.7 (7)
N1—Pb1—O3 <sup>i</sup>	140.87 (15)	C19—C14—C13	121.5 (6)
O1—Pb1—O3 <sup>i</sup>	137.97 (14)	C15—C14—C13	119.8 (6)
N2—Pb1—O3 <sup>i</sup>	78.83 (15)	C14—C19—C18	119.9 (8)
O2—Pb1—O3 <sup>i</sup>	91.08 (14)	C14—C19—H19	120.1
O3—Pb1—O3 <sup>i</sup>	66.84 (16)	C18—C19—H19	120.1
C2—C3—C4	120.0 (6)	C16—C15—C14	120.3 (7)
C2—C3—H3	120.0	C16—C15—H15	119.8
C4—C3—H3	120.0	C14—C15—H15	119.8
C3—C2—C1	119.3 (7)	C17—C18—C19	121.0 (9)
C3—C2—H2	120.3	C17—C18—H18	119.5
C1—C2—H2	120.3	C19—C18—H18	119.5
N1—C1—C2	122.8 (7)	C16—C17—C18	120.1 (9)
N1—C1—H1	118.6	C16—C17—H17	120.0
C2—C1—H1	118.6	C18—C17—H17	120.0
Pb1—C1—H1	78.0	C17—C16—C15	120.0 (9)
Pb1 <sup>ii</sup> —C1—H1	66.1	C17—C16—H16	120.0
C1—N1—C5	118.4 (5)	C15—C16—H16	120.0
C1—N1—Pb1	119.7 (4)	C20—O3—Pb1	85.0 (4)
C5—N1—Pb1	121.7 (4)	C20—O3—Pb1 <sup>i</sup>	155.1 (4)
N1—C5—C4	122.0 (6)	Pb1—O3—Pb1 <sup>i</sup>	113.16 (16)
N1—C5—C9	118.6 (5)	C20—O4—Pb1	103.6 (4)
C4—C5—C9	119.3 (6)	O3—C20—O4	122.8 (6)
N2—C9—C8	121.7 (6)	O3—C20—C21	121.1 (6)
N2—C9—C5	118.8 (5)	O4—C20—C21	116.1 (6)
C8—C9—C5	119.4 (6)	C26—C21—C22	118.7 (6)
C12—N2—C9	118.9 (5)	C26—C21—C20	121.5 (6)
C12—N2—Pb1	123.4 (4)	C22—C21—C20	119.8 (6)
C9—N2—Pb1	117.2 (4)	C25—C26—C21	121.5 (8)
N2—C12—C11	122.0 (6)	C25—C26—H26	119.2
N2—C12—H12	119.0	C21—C26—H26	119.2

C11—C12—H12	119.0	C26—C25—C24	119.2 (9)
Pb1—C12—H12	80.8	C26—C25—H25	120.4
C10—C11—C12	119.7 (7)	C24—C25—H25	120.4
C10—C11—H11	120.2	C23—C22—C21	120.1 (8)
C12—C11—H11	120.2	C23—C22—H22	120.0
C11—C10—C8	119.9 (6)	C21—C22—H22	120.0
C11—C10—H10	120.0	C24—C23—C22	120.4 (9)
C8—C10—H10	120.0	C24—C23—H23	119.8
C10—C8—C9	117.7 (6)	C22—C23—H23	119.8
C10—C8—C7	122.9 (6)	C23—C24—C25	120.1 (8)
C9—C8—C7	119.4 (6)	C23—C24—H24	120.0
C6—C7—C8	121.1 (7)	C25—C24—H24	120.0

Symmetry codes: (i)  $-x+1, -y+2, -z+1$ ; (ii)  $-x, -y+2, -z+1$ .