

Bis[μ -N'-(2-oxidobenzylidene)thiophene-2-carbohydrazidato]bis[dimethanol-nickel(II)]

Jian-Hua Ma, Wen-Shi Wu,* Xiao-Qing Zhang, Sheng-Jiao Tang and Xin-Yong Lin

College of Materials Science and Engineering, Huaqiao University, Xiamen, Fujian 361021, People's Republic of China

Correspondence e-mail: wwws@hqu.edu.cn


Received 2 March 2011; accepted 3 June 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; disorder in main residue; R factor = 0.037; wR factor = 0.089; data-to-parameter ratio = 16.0.

In the crystal structure of the centrosymmetric binuclear title complex, $[\text{Ni}_2(\text{C}_{12}\text{H}_8\text{N}_2\text{O}_2\text{S})_2(\text{CH}_3\text{OH})_4]$, there are intermolecular $\text{O}-\text{H}\cdots\text{O}$, $\text{O}-\text{H}\cdots\text{N}$ and $\text{O}-\text{H}\cdots\text{S}$ hydrogen bonds. These help to stabilize the structure and link the molecules, forming a three-dimensional network structure. The Ni^{2+} cation exists in a slightly distorted octahedral NiNO_5 coordination environment. The thiophene rings are disordered over two equivalent conformations with occupancies of 0.881 (3) and 0.119 (3).

Related literature

For the structure of the related Cu complex, see: Lu *et al.* (2006). For the synthesis of the ligand, see: Wu *et al.* (2004).



Experimental

Crystal data

$[\text{Ni}_2(\text{C}_{12}\text{H}_8\text{N}_2\text{O}_2\text{S})_2(\text{CH}_3\text{OH})_4]$

$M_r = 734.12$

Monoclinic, $P2_1/n$

$a = 13.7958$ (14) \AA

$b = 7.8880$ (8) \AA

$c = 14.4219$ (16) \AA

$\beta = 104.672$ (1) $^\circ$

$V = 1518.2$ (3) \AA^3

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 1.43$ mm $^{-1}$

$T = 293$ K

$0.60 \times 0.41 \times 0.39$ mm

Data collection

Bruker SMART CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.499$, $T_{\max} = 0.572$

11412 measured reflections

3484 independent reflections

3137 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.089$

$S = 1.11$

3469 reflections

217 parameters

10 restraints

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.36$ e \AA^{-3}

$\Delta\rho_{\text{min}} = -0.34$ e \AA^{-3}

Table 1
Selected bond lengths (\AA).

Ni1–N2	1.9906 (18)	Ni1–O2 ⁱ	2.0521 (15)
Ni1–O2	2.0165 (16)	Ni1–O3	2.1425 (17)
Ni1–O1	2.0244 (16)	Ni1–O4	2.1897 (16)

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O3–H16 \cdots O4 ⁱ	0.82	2.13	2.920 (2)	163
O4–H15 \cdots N1 ⁱⁱ	0.82	2.07	2.850 (2)	159
O4–H15 \cdots S1A ⁱⁱ	0.82	2.92	3.452 (3)	125

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 1999); cell refinement: *SAINT* (Bruker, 1999); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors are grateful for financial support from The National Science Foundation of Fujian Province of China (No. 2010J01288) and Huaqiao University Basic Research Special Fund operating expenses (No.JB-JC1003).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BV2181).

References

- Bruker (1999). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Lu, Y., Chen, J., Wu, W.-S., Dai, J.-C. & Lin, J.-M. (2006). *Acta Cryst. E62*, m1291–m1292.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Wu, W. S., Feng, Y. L., Lan, X. R. & Huang, T. T. (2004). *Chin. J. Appl. Chem. A21*, 135–139.

supporting information

Acta Cryst. (2011). E67, m910 [doi:10.1107/S160053681102143X]

Bis[μ -N'-(2-oxidobenzylidene)thiophene-2-carbohydrazidato]bis[dimethano-Nnickel(II)]

Jian-Hua Ma, Wen-Shi Wu, Xiao-Qing Zhang, Sheng-Jiao Tang and Xin-Yong Lin

S1. Comment

The structure of $[\text{Cu}(\text{C}_{12}\text{H}_9\text{N}_2\text{O}_2\text{S})\text{Cl}]\cdot\text{H}_2\text{O}$ (II), which is closely related to the title complex has already been reported (Lu *et al.*, 2006). In this article, we report the crystal structure of the realted nickel complex, I.

In the title complex, $[\text{Ni}_2(\text{C}_{24}\text{H}_{16}\text{N}_4\text{O}_4\text{S}_2)\text{C}_4\text{H}_{16}\text{O}_4]$ (Fig.1 and Table 1), the Ni(II) cation is six-coordinated by an amide N, a phenoxide O and a carbonyl O atom derived from the first tridentate ligand, another phenoxide O atom derived from the second tridentate ligand and two other O atoms from two molecules of methanol. The five-and six-membered chelate rings are coplanar, the mean deviation of 0.0497 \AA from the least-squares plane through both of them, slightly larger than that of the structure of II ($0.037(18)\text{ \AA}$, Lu *et al.*). The thiophene rings are slightly twisted with respect to the above mentioned plane, making a larger dihedral angle of $15.4(2)^\circ$ than that of II ($8.77(9)^\circ$). The Ni-O and Ni-N distances [$\text{Ni}-\text{O}1(2.024(2)\text{ \AA})$, $\text{Ni}-\text{O}2(2.016(2)\text{ \AA})$ and $\text{Ni}-\text{N}2(1.991(2)\text{ \AA})$] are longer than the corresponding distances for II (Lu *et al.*, 2006). The $\text{O}1-\text{Ni}1-\text{N}2(79.53(7)^\circ)$ and $\text{O}2-\text{Ni}1-\text{N}2(91.24(7)^\circ)$ angles differ slightly from the corresponding angles found in II. The thiophene rings are disordered over two eqivalent conformations with occupancies of 0.881 (3) and 0.119 (3) as is commonly found for this moiety.


N—H \cdots O, N—H \cdots S and O—H \cdots O intermolecular hydrogen bonds in the compound stabilize the structure and link the molecules in a three-dimensional network structure (shown in Fig.2) and detailed in Table 2.

S2. Experimental


The ligand was synthesized according to the method of Wu *et al.*,(2004). NiCl_2 (1 mmol) and the ligand were separately dissolved in 20 ml methanol and the resulting solutions mixed slowly, stirred at room temperature for one hour, and then filtered. The filtrate was allowed to stand at room temperature for ten days, yielding deep-blue crystals of the title compound by slow evaporation.

S3. Refinement

The C-bound H atoms were included in the riding model approximation with C—H = $0.93 - 0.96\text{ \AA}$, all these H atoms included in the final refinement. The U_{iso} of each H atom = $1.2U_{\text{eq}}(\text{C})$ [$1.5U_{\text{eq}}(\text{C})$ for CH_3]. The methanol H atoms were refined isotropically.

**Figure 1**


The molecular structure (at 30% probability) of the title compound.

**Figure 2**

Packing diagram of the title complex, showing hydrogen bonds as dashed lines.

Bis[μ -N'-(2-oxidobenzylidene)thiophene-2- carbohydrazidato]bis[dimethanolnickel(II)]

Crystal data



$M_r = 734.12$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 13.7958 (14)$ Å

$b = 7.8880 (8)$ Å

$c = 14.4219 (16)$ Å

$\beta = 104.672 (1)^\circ$

$V = 1518.2 (3)$ Å³

$Z = 2$

$F(000) = 760$

$D_x = 1.606 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1552 reflections

$\theta = 3.1\text{--}27.5^\circ$

$\mu = 1.43 \text{ mm}^{-1}$

$T = 293$ K

Columnar, blue
 $0.60 \times 0.41 \times 0.39$ mm

Data collection

Bruker SMART CCD diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: 0 pixels mm⁻¹
 ω scans
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.499$, $T_{\max} = 0.572$

11412 measured reflections
 3484 independent reflections
 3137 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.1^\circ$
 $h = -17 \rightarrow 17$
 $k = 0 \rightarrow 10$
 $l = 0 \rightarrow 18$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.089$
 $S = 1.11$
 3469 reflections
 217 parameters
 10 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0375P)^2 + 0.8784P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.36 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. 2011-03-01 # Formatted by IUCr publCIF system

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ni1	0.39217 (2)	0.56046 (3)	0.449805 (19)	0.02409 (10)	
S1A	0.03964 (10)	0.9026 (2)	0.32607 (15)	0.0563 (5)	0.881 (3)
C1A	0.1016 (3)	0.7392 (5)	0.3958 (3)	0.0273 (7)	0.881 (3)
C2A	0.0491 (6)	0.6804 (10)	0.4605 (6)	0.0408 (11)	0.881 (3)
H2AA	0.0709	0.5952	0.5056	0.049*	0.881 (3)
C3A	-0.0430 (3)	0.7708 (5)	0.4469 (4)	0.0436 (9)	0.881 (3)
H3AA	-0.0897	0.7471	0.4816	0.052*	0.881 (3)
C4A	-0.0570 (3)	0.8928 (6)	0.3797 (3)	0.0495 (10)	0.881 (3)
H4AA	-0.1127	0.9635	0.3643	0.059*	0.881 (3)
S1B	0.0546 (13)	0.687 (2)	0.4652 (14)	0.0563 (5)	0.119 (3)
C1B	0.104 (3)	0.767 (5)	0.379 (3)	0.0273 (7)	0.119 (3)
C2B	0.053 (3)	0.914 (6)	0.338 (4)	0.0408 (11)	0.119 (3)
H2BA	0.0773	1.0035	0.3088	0.049*	0.119 (3)
C3B	-0.046 (2)	0.896 (5)	0.352 (2)	0.0436 (9)	0.119 (3)
H3BA	-0.1026	0.9367	0.3086	0.052*	0.119 (3)

C4B	-0.049 (3)	0.814 (6)	0.433 (3)	0.0495 (10)	0.119 (3)
H4BA	-0.0992	0.8237	0.4652	0.059*	0.119 (3)
O1	0.24595 (12)	0.5752 (2)	0.45082 (12)	0.0313 (4)	
O2	0.53297 (11)	0.57985 (18)	0.43430 (11)	0.0263 (3)	
O3	0.42607 (13)	0.7737 (2)	0.54454 (13)	0.0401 (4)	
H16	0.4807	0.7471	0.5794	0.060*	
O4	0.36612 (12)	0.3220 (2)	0.36958 (12)	0.0322 (4)	
H15	0.3433	0.3266	0.3113	0.048*	
N1	0.24055 (14)	0.7622 (2)	0.32434 (14)	0.0288 (4)	
N2	0.34061 (13)	0.7130 (2)	0.33827 (13)	0.0259 (4)	
C5	0.20199 (16)	0.6862 (3)	0.38915 (16)	0.0262 (4)	
C6	0.39124 (17)	0.7867 (3)	0.28636 (16)	0.0285 (5)	
H6A	0.3575	0.8637	0.2406	0.034*	
C7	0.49684 (16)	0.7591 (3)	0.29352 (16)	0.0260 (4)	
C8	0.56252 (16)	0.6577 (3)	0.36440 (15)	0.0244 (4)	
C9	0.66258 (17)	0.6423 (3)	0.35929 (17)	0.0317 (5)	
H9A	0.7066	0.5765	0.4047	0.038*	
C10	0.69723 (19)	0.7222 (3)	0.28877 (19)	0.0374 (6)	
H10A	0.7638	0.7088	0.2873	0.045*	
C11	0.63412 (19)	0.8220 (3)	0.22015 (18)	0.0375 (6)	
H11A	0.6578	0.8766	0.1731	0.045*	
C12	0.53578 (18)	0.8385 (3)	0.22323 (18)	0.0330 (5)	
H12A	0.4932	0.9049	0.1770	0.040*	
C13	0.3631 (3)	0.8355 (4)	0.6013 (3)	0.0574 (8)	
H13A	0.3956	0.9281	0.6402	0.086*	
H13B	0.3503	0.7462	0.6419	0.086*	
H13C	0.3009	0.8738	0.5602	0.086*	
C14	0.3199 (2)	0.1845 (3)	0.4077 (2)	0.0448 (6)	
H14A	0.2951	0.1025	0.3582	0.067*	
H14B	0.2654	0.2271	0.4311	0.067*	
H14C	0.3685	0.1319	0.4593	0.067*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.02149 (15)	0.02730 (16)	0.02132 (16)	0.00196 (10)	0.00142 (11)	0.00409 (11)
S1A	0.0438 (7)	0.0593 (7)	0.0688 (10)	0.0245 (6)	0.0200 (7)	0.0337 (8)
C1A	0.0267 (11)	0.023 (2)	0.028 (2)	0.0005 (12)	0.0005 (12)	0.0056 (11)
C2A	0.036 (2)	0.047 (2)	0.043 (2)	-0.0097 (15)	0.0153 (15)	-0.0102 (16)
C3A	0.0327 (16)	0.051 (3)	0.050 (2)	0.0043 (15)	0.0156 (16)	0.0032 (17)
C4A	0.0349 (18)	0.054 (2)	0.062 (3)	0.0181 (15)	0.0165 (17)	0.009 (2)
S1B	0.0438 (7)	0.0593 (7)	0.0688 (10)	0.0245 (6)	0.0200 (7)	0.0337 (8)
C1B	0.0267 (11)	0.023 (2)	0.028 (2)	0.0005 (12)	0.0005 (12)	0.0056 (11)
C2B	0.036 (2)	0.047 (2)	0.043 (2)	-0.0097 (15)	0.0153 (15)	-0.0102 (16)
C3B	0.0327 (16)	0.051 (3)	0.050 (2)	0.0043 (15)	0.0156 (16)	0.0032 (17)
C4B	0.0349 (18)	0.054 (2)	0.062 (3)	0.0181 (15)	0.0165 (17)	0.009 (2)
O1	0.0240 (8)	0.0361 (9)	0.0320 (9)	0.0034 (6)	0.0040 (7)	0.0105 (7)
O2	0.0248 (8)	0.0310 (8)	0.0210 (8)	0.0017 (6)	0.0018 (6)	0.0074 (6)

O3	0.0373 (9)	0.0399 (10)	0.0386 (10)	0.0025 (7)	0.0015 (8)	-0.0098 (8)
O4	0.0360 (9)	0.0334 (8)	0.0236 (8)	-0.0058 (7)	0.0008 (7)	-0.0022 (7)
N1	0.0247 (9)	0.0312 (10)	0.0276 (10)	0.0051 (7)	0.0012 (8)	0.0040 (8)
N2	0.0222 (9)	0.0284 (9)	0.0239 (10)	0.0017 (7)	0.0000 (7)	0.0028 (7)
C5	0.0247 (10)	0.0264 (10)	0.0244 (11)	0.0013 (8)	0.0003 (9)	-0.0003 (8)
C6	0.0308 (11)	0.0272 (11)	0.0238 (11)	0.0025 (9)	-0.0001 (9)	0.0051 (9)
C7	0.0301 (11)	0.0241 (10)	0.0220 (11)	-0.0029 (8)	0.0034 (9)	0.0003 (8)
C8	0.0275 (11)	0.0249 (10)	0.0195 (10)	-0.0028 (8)	0.0035 (8)	-0.0023 (8)
C9	0.0274 (11)	0.0382 (12)	0.0280 (12)	0.0012 (9)	0.0045 (9)	0.0032 (10)
C10	0.0277 (12)	0.0489 (14)	0.0360 (14)	-0.0027 (10)	0.0088 (10)	0.0009 (11)
C11	0.0397 (14)	0.0458 (14)	0.0292 (13)	-0.0085 (11)	0.0127 (11)	0.0070 (11)
C12	0.0367 (13)	0.0327 (12)	0.0273 (12)	-0.0011 (10)	0.0042 (10)	0.0070 (9)
C13	0.065 (2)	0.0542 (18)	0.059 (2)	-0.0006 (15)	0.0263 (17)	-0.0225 (15)
C14	0.0552 (17)	0.0366 (13)	0.0403 (15)	-0.0131 (12)	0.0080 (13)	0.0013 (11)

Geometric parameters (\AA , $^\circ$)

Ni1—N2	1.9906 (18)	O2—Ni1 ⁱ	2.0521 (15)
Ni1—O2	2.0165 (16)	O3—C13	1.422 (3)
Ni1—O1	2.0244 (16)	O3—H16	0.8200
Ni1—O2 ⁱ	2.0521 (15)	O4—C14	1.436 (3)
Ni1—O3	2.1425 (17)	O4—H15	0.8206
Ni1—O4	2.1897 (16)	N1—C5	1.330 (3)
S1A—C4A	1.704 (4)	N1—N2	1.399 (3)
S1A—C1A	1.722 (3)	N2—C6	1.285 (3)
C1A—C2A	1.397 (8)	C6—C7	1.451 (3)
C1A—C5	1.474 (4)	C6—H6A	0.9300
C2A—C3A	1.427 (8)	C7—C12	1.409 (3)
C2A—H2AA	0.9300	C7—C8	1.427 (3)
C3A—C4A	1.345 (5)	C8—C9	1.406 (3)
C3A—H3AA	0.9300	C9—C10	1.381 (3)
C4A—H4AA	0.9300	C9—H9A	0.9300
S1B—C1B	1.680 (18)	C10—C11	1.386 (4)
S1B—C4B	1.710 (18)	C10—H10A	0.9300
C1B—C2B	1.41 (2)	C11—C12	1.375 (4)
C1B—C5	1.47 (3)	C11—H11A	0.9300
C2B—C3B	1.43 (2)	C12—H12A	0.9300
C2B—H2BA	0.9300	C13—H13A	0.9600
C3B—C4B	1.344 (19)	C13—H13B	0.9600
C3B—H3BA	0.9300	C13—H13C	0.9600
C4B—H4BA	0.9300	C14—H14A	0.9600
O1—C5	1.285 (3)	C14—H14B	0.9600
O2—C8	1.330 (3)	C14—H14C	0.9600
N2—Ni1—O2	91.24 (7)	Ni1—O3—H16	102.0
N2—Ni1—O1	79.53 (7)	C14—O4—Ni1	118.42 (15)
O2—Ni1—O1	170.42 (6)	C14—O4—H15	109.5
N2—Ni1—O2 ⁱ	170.91 (7)	Ni1—O4—H15	118.2

O2—Ni1—O2 ⁱ	80.27 (7)	C5—N1—N2	109.27 (17)
O1—Ni1—O2 ⁱ	108.78 (6)	C6—N2—N1	116.97 (18)
N2—Ni1—O3	90.93 (8)	C6—N2—Ni1	127.48 (15)
O2—Ni1—O3	87.16 (7)	N1—N2—Ni1	114.92 (14)
O1—Ni1—O3	90.41 (7)	O1—C5—N1	126.2 (2)
O2 ⁱ —Ni1—O3	85.43 (7)	O1—C5—C1B	128.6 (10)
N2—Ni1—O4	96.61 (7)	N1—C5—C1B	105.0 (9)
O2—Ni1—O4	92.32 (6)	O1—C5—C1A	115.8 (2)
O1—Ni1—O4	91.30 (6)	N1—C5—C1A	117.9 (2)
O2 ⁱ —Ni1—O4	87.05 (6)	N2—C6—C7	125.1 (2)
O3—Ni1—O4	172.44 (7)	N2—C6—H6A	117.4
C4A—S1A—C1A	91.74 (17)	C7—C6—H6A	117.4
C2A—C1A—C5	127.2 (4)	C12—C7—C8	118.4 (2)
C2A—C1A—S1A	112.0 (4)	C12—C7—C6	116.2 (2)
C5—C1A—S1A	120.6 (2)	C8—C7—C6	125.4 (2)
C1A—C2A—C3A	109.6 (6)	O2—C8—C9	119.57 (19)
C1A—C2A—H2AA	125.2	O2—C8—C7	122.76 (19)
C3A—C2A—H2AA	125.2	C9—C8—C7	117.7 (2)
C4A—C3A—C2A	114.6 (5)	C10—C9—C8	121.8 (2)
C4A—C3A—H3AA	122.7	C10—C9—H9A	119.1
C2A—C3A—H3AA	122.7	C8—C9—H9A	119.1
C3A—C4A—S1A	112.0 (3)	C9—C10—C11	120.9 (2)
C3A—C4A—H4AA	124.0	C9—C10—H10A	119.6
S1A—C4A—H4AA	124.0	C11—C10—H10A	119.6
C1B—S1B—C4B	92.8 (12)	C12—C11—C10	118.5 (2)
C2B—C1B—C5	139 (2)	C12—C11—H11A	120.7
C2B—C1B—S1B	111.3 (17)	C10—C11—H11A	120.7
C5—C1B—S1B	107.6 (17)	C11—C12—C7	122.7 (2)
C1B—C2B—C3B	105 (2)	C11—C12—H12A	118.6
C1B—C2B—H2BA	127.7	C7—C12—H12A	118.6
C3B—C2B—H2BA	127.7	O3—C13—H13A	109.5
C4B—C3B—C2B	114 (2)	O3—C13—H13B	109.5
C4B—C3B—H3BA	122.8	H13A—C13—H13B	109.5
C2B—C3B—H3BA	122.8	O3—C13—H13C	109.5
C3B—C4B—S1B	107.8 (18)	H13A—C13—H13C	109.5
C3B—C4B—H4BA	126.1	H13B—C13—H13C	109.5
S1B—C4B—H4BA	126.1	O4—C14—H14A	109.5
C5—O1—Ni1	109.30 (14)	O4—C14—H14B	109.5
C8—O2—Ni1	127.45 (13)	H14A—C14—H14B	109.5
C8—O2—Ni1 ⁱ	132.76 (14)	O4—C14—H14C	109.5
Ni1—O2—Ni1 ⁱ	99.73 (7)	H14A—C14—H14C	109.5
C13—O3—Ni1	124.72 (17)	H14B—C14—H14C	109.5
C13—O3—H16	109.6		
C4A—S1A—C1A—C2A	-1.0 (5)	O1—Ni1—N2—N1	5.62 (14)
C4A—S1A—C1A—C5	-176.0 (4)	O3—Ni1—N2—N1	-84.62 (15)
C5—C1A—C2A—C3A	176.7 (5)	O4—Ni1—N2—N1	95.73 (15)
S1A—C1A—C2A—C3A	2.0 (7)	Ni1—O1—C5—N1	9.3 (3)

C1A—C2A—C3A—C4A	−2.3 (7)	Ni1—O1—C5—C1B	−165 (3)
C2A—C3A—C4A—S1A	1.6 (6)	Ni1—O1—C5—C1A	−167.0 (3)
C1A—S1A—C4A—C3A	−0.4 (4)	N2—N1—C5—O1	−4.8 (3)
C4B—S1B—C1B—C2B	12 (5)	N2—N1—C5—C1B	171 (2)
C4B—S1B—C1B—C5	178 (3)	N2—N1—C5—C1A	171.5 (3)
C5—C1B—C2B—C3B	175 (5)	C2B—C1B—C5—O1	159 (5)
S1B—C1B—C2B—C3B	−26 (5)	S1B—C1B—C5—O1	0 (4)
C1B—C2B—C3B—C4B	33 (6)	C2B—C1B—C5—N1	−17 (7)
C2B—C3B—C4B—S1B	−25 (5)	S1B—C1B—C5—N1	−176 (2)
C1B—S1B—C4B—C3B	7 (4)	C2B—C1B—C5—C1A	166 (16)
N2—Ni1—O1—C5	−7.40 (15)	S1B—C1B—C5—C1A	7 (8)
O2 ⁱ —Ni1—O1—C5	168.79 (14)	C2A—C1A—C5—O1	−1.0 (7)
O3—Ni1—O1—C5	83.47 (15)	S1A—C1A—C5—O1	173.2 (3)
O4—Ni1—O1—C5	−103.89 (15)	C2A—C1A—C5—N1	−177.7 (5)
N2—Ni1—O2—C8	−5.77 (17)	S1A—C1A—C5—N1	−3.4 (5)
O2 ⁱ —Ni1—O2—C8	177.5 (2)	C2A—C1A—C5—C1B	−174 (11)
O3—Ni1—O2—C8	−96.64 (17)	S1A—C1A—C5—C1B	0 (10)
O4—Ni1—O2—C8	90.90 (17)	N1—N2—C6—C7	176.82 (19)
N2—Ni1—O2—Ni1 ⁱ	176.73 (7)	Ni1—N2—C6—C7	6.4 (3)
O2 ⁱ —Ni1—O2—Ni1 ⁱ	0.0	N2—C6—C7—C12	173.4 (2)
O3—Ni1—O2—Ni1 ⁱ	85.85 (7)	N2—C6—C7—C8	−5.7 (4)
O4—Ni1—O2—Ni1 ⁱ	−86.60 (7)	Ni1—O2—C8—C9	−172.88 (15)
N2—Ni1—O3—C13	97.2 (2)	Ni1 ⁱ —O2—C8—C9	3.8 (3)
O2—Ni1—O3—C13	−171.6 (2)	Ni1—O2—C8—C7	7.7 (3)
O1—Ni1—O3—C13	17.7 (2)	Ni1 ⁱ —O2—C8—C7	−175.67 (15)
O2 ⁱ —Ni1—O3—C13	−91.1 (2)	C12—C7—C8—O2	179.0 (2)
N2—Ni1—O4—C14	−134.19 (17)	C6—C7—C8—O2	−1.8 (3)
O2—Ni1—O4—C14	134.29 (17)	C12—C7—C8—C9	−0.4 (3)
O1—Ni1—O4—C14	−54.58 (17)	C6—C7—C8—C9	178.7 (2)
O2 ⁱ —Ni1—O4—C14	54.16 (17)	O2—C8—C9—C10	−179.3 (2)
C5—N1—N2—C6	−174.3 (2)	C7—C8—C9—C10	0.2 (3)
C5—N1—N2—Ni1	−2.7 (2)	C8—C9—C10—C11	0.3 (4)
O2—Ni1—N2—C6	−1.2 (2)	C9—C10—C11—C12	−0.6 (4)
O1—Ni1—N2—C6	176.2 (2)	C10—C11—C12—C7	0.4 (4)
O3—Ni1—N2—C6	85.9 (2)	C8—C7—C12—C11	0.1 (4)
O4—Ni1—N2—C6	−93.7 (2)	C6—C7—C12—C11	−179.1 (2)
O2—Ni1—N2—N1	−171.80 (14)		

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^{\circ}$)

$D\cdots H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O3—H16 \cdots O4 ⁱ	0.82	2.13	2.920 (2)	163
O4—H15 \cdots N1 ⁱⁱ	0.82	2.07	2.850 (2)	159
O4—H15 \cdots S1A ⁱⁱ	0.82	2.92	3.452 (3)	125

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+1/2, y-1/2, -z+1/2$.