

N-(4-Chlorobutanoyl)-N'-(2-fluoro-phenyl)thiourea

M. Sukeri M. Yusof,^{a*} Nur Farhana Embong,^a Eliyanti A. Othman^b and Bohari M. Yamin^b

^aDepartment of Chemical Sciences, Faculty of Science and Technology, Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Terengganu, Malaysia, and ^bSchool of Chemical Sciences and Food Technology, Universiti Kebangsaan Malaysia, UKM 43500 Bangi Selangor, Malaysia

Correspondence e-mail: mohdsukeri@umt.edu.my


Received 6 June 2011; accepted 23 June 2011

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$; R factor = 0.063; wR factor = 0.138; data-to-parameter ratio = 14.7.

In the title compound, $\text{C}_{11}\text{H}_{12}\text{ClFN}_2\text{OS}$, the asymmetric unit consists of two independent molecules. Both molecules maintain a *trans-cis* configuration of the positions of the butanoyl and fluorophenyl groups with respect to the thiono group across their C–N bonds and are stabilized by classical intramolecular N–H···O hydrogen bonds. In the crystal, intermolecular N–H···O, C–H···S and N–H···S hydrogen bonds link the molecules into infinite chains along the c axis.

Related literature

For a related structure, see: Yamin *et al.* (2011). For standard bond lengths, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{11}\text{H}_{12}\text{ClFN}_2\text{OS}$
 $M_r = 274.75$
Monoclinic, $P2_1/c$
 $a = 14.818 (7)\text{ \AA}$

$b = 10.291 (5)\text{ \AA}$
 $c = 18.201 (9)\text{ \AA}$
 $\beta = 112.599 (12)^\circ$
 $V = 2562 (2)\text{ \AA}^3$

$Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.46\text{ mm}^{-1}$

$T = 298\text{ K}$
 $0.50 \times 0.22 \times 0.07\text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.803$, $T_{\max} = 0.969$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.063$
 $wR(F^2) = 0.138$
 $S = 0.89$
4501 reflections

307 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.34\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.30\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

| $D-\text{H} \cdots A$ | $D-\text{H}$ | $\text{H} \cdots A$ | $D \cdots A$ | $D-\text{H} \cdots A$ |
|------------------------------|--------------|---------------------|--------------|-----------------------|
| N2–H2···O1 | 0.86 | 2.02 | 2.691 (5) | 134 |
| N2–H2···O2 ⁱ | 0.86 | 2.40 | 3.128 (5) | 143 |
| N4–H4···O2 | 0.86 | 2.02 | 2.676 (5) | 133 |
| N4–H4···O1 ⁱⁱ | 0.86 | 2.32 | 3.033 (5) | 141 |
| N3–H3···S1 ⁱⁱⁱ | 0.86 | 2.59 | 3.447 (4) | 176 |
| N1–H1···S2 ⁱⁱⁱ | 0.86 | 2.52 | 3.364 (4) | 169 |
| C14–H14A···S1 ⁱⁱⁱ | 0.97 | 2.96 | 3.784 (5) | 143 |
| C14–H14B···S2 ^{iv} | 0.97 | 2.74 | 3.691 (5) | 168 |

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, -y + 1, -z$; (iv) $-x, -y + 2, -z$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*, *PARST* (Nardelli, 1995) and *PLATON* (Spek, 2009).

The authors thank the Malaysian Government, Universiti Kebangsaan Malaysia, Universiti Malaysia Terengganu and the Ministry of Higher Education, Malaysia, for research grants UKM-GUP-NBT-08-27-110 and FRGS 59178.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2280).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2000). *SADABS*, *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Yamin, B. M., Othman, N. E. A., Yusof, M. S. M. & Embong, F. (2011). *Acta Cryst. E* **67**, o419.

supporting information

Acta Cryst. (2011). E67, o1849 [doi:10.1107/S1600536811024743]

N-(4-Chlorobutanoyl)-N'-(2-fluorophenyl)thiourea

M. Sukeri M. Yusof, Nur Farhana Embong, Eliyanti A. Othman and Bohari M. Yamin

S1. Comment


The title compound, is analogous to the previously reported *N*-(4-chlorobutanoyl)-*N'*-(phenyl)thiourea (Yamin *et al.*, 2011) except the fluoro atom is attached at the *ortho*-position of the phenyl ring. The asymmetric unit consists of two independent molecules (Fig. 1). The whole molecule is not planar. However, the thiourea N1/C5/S1/N2/C6, N3/C16/S2/N4/C17 fragments and the benzene rings, (C6–C11) and (C17–C22) are each planar with maximum deviation of 0.020 (3) Å for N4 atom from the least square plane. In each molecule, the benzene ring and thiourea moiety forms dihedral angle of 74.78 (19)° and 82.3 (2)°, respectively. The same dihedral angle in *N*-(4-chlorobutanoyl)-*N'*-(phenyl)-thiourea are 72.98 (12)° and 81.47 (14)°, respectively. The bond lengths and angles are in normal ranges (Allen *et al.*, 1987) and comparable to those in the analog. Both molecules maintain their *trans*–*cis*-configuration with respect to the position of the butanoyl and fluorophenyl groups against the thiono C=S-group bond across the C—N bonds. Like in most of the carbonylthiourea derivatives, the classical intramolecular hydrogen bonds between the carbonyl oxygen atom and thioamide hydrogen atom, N2—H2···O1 and N4—H4···O2, in both molecules are present. In the crystal packing, the molecules are linked by N3—H3···S1ⁱⁱⁱ and N1—H1···S2ⁱⁱⁱⁱ, N2—H2···O2ⁱ and N4—H4···O1ⁱⁱ; C14—H14A···S1ⁱⁱⁱ and C14—H14B···S2^{iv} intermolecular hydrogen bonds (symmetry codes as in Table 1) and form infinite chains along the *c*-axis (Fig. 2).

S2. Experimental


A solution of 4-chlorobutanoylisothiocyanate (1.25 g, 6.33 mmol) in 30 ml acetone was added into a flask containing 30 ml acetone solution of 2-floroaniline (0.71 g, 6.33 mmol). The mixture was refluxed for 1 h. Then, the solution was filtered-off and left to evaporate at room temperature. The colourless solid was obtained after one day of evaporation (yield 83%, m.p. 411.7 K–415.5 K)

S3. Refinement

All H atoms attached to C and N atoms were fixed geometrically and treated as riding with C—H = 0.93 Å or 0.97 Å (aromatic and methylene) and N—H = 0.86 Å (amino) with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$.

**Figure 1**

The molecular structure of title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as a small spheres of arbitrary radius. intramolecular H bonds are presented by dashed lines.

**Figure 2**

A packing diagram of title compound viewed down the a -axis. H bonds are shown by dashed lines.

N-(4-Chlorobutanoyl)-*N'*-(2-fluorophenyl)thiourea

Crystal data

$C_{11}H_{12}ClFN_2OS$

$M_r = 274.75$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 14.818 (7)$ Å

$b = 10.291 (5)$ Å

$c = 18.201 (9)$ Å

$\beta = 112.599 (12)^\circ$

$V = 2562 (2)$ Å³

$Z = 8$

$F(000) = 1136$

$D_x = 1.425$ Mg m⁻³

Melting point = 411.7–415.5 K

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1404 reflections

$\theta = 1.5\text{--}25.0^\circ$

$\mu = 0.46$ mm⁻¹

$T = 298$ K

Plate, colourless

$0.50 \times 0.22 \times 0.07$ mm

Data collection

Bruker SMART APEX CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube
Graphite monochromator

Detector resolution: 83.66 pixels mm⁻¹

ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2000)

$T_{\min} = 0.803$, $T_{\max} = 0.969$
 13771 measured reflections
 4501 independent reflections
 2231 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.119$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.5^\circ$
 $h = -17 \rightarrow 16$
 $k = -12 \rightarrow 12$
 $l = -13 \rightarrow 21$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.063$
 $wR(F^2) = 0.138$
 $S = 0.89$
 4501 reflections
 307 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0451P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.34 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|---------------|--------------|----------------------------------|
| Cl1 | 0.49078 (11) | 0.21242 (19) | 0.00799 (10) | 0.1269 (7) |
| F1 | 1.0669 (2) | -0.1688 (3) | 0.35144 (16) | 0.0795 (9) |
| F2 | 0.3894 (2) | 0.7354 (3) | 0.12270 (17) | 0.0868 (9) |
| S1 | 1.02276 (8) | -0.03047 (12) | 0.17159 (6) | 0.0509 (3) |
| S2 | 0.24427 (8) | 1.01331 (12) | 0.02342 (6) | 0.0502 (3) |
| Cl2 | -0.23288 (9) | 0.68710 (13) | -0.21630 (7) | 0.0804 (5) |
| O1 | 0.80228 (19) | 0.1449 (3) | 0.26210 (18) | 0.0562 (9) |
| O2 | 0.0637 (2) | 0.6917 (3) | 0.07106 (17) | 0.0552 (9) |
| N1 | 0.8543 (2) | 0.0465 (3) | 0.17333 (18) | 0.0424 (9) |
| H1 | 0.8335 | 0.0212 | 0.1246 | 0.051* |
| N2 | 0.9869 (2) | 0.0693 (3) | 0.29214 (19) | 0.0428 (9) |
| H2 | 0.9467 | 0.1033 | 0.3103 | 0.051* |
| N3 | 0.0918 (2) | 0.8641 (3) | 0.00426 (18) | 0.0421 (9) |
| H3 | 0.0658 | 0.9086 | -0.0388 | 0.051* |
| N4 | 0.2217 (2) | 0.8449 (3) | 0.12431 (18) | 0.0447 (9) |
| H4 | 0.1883 | 0.7860 | 0.1360 | 0.054* |
| C1 | 0.5036 (3) | 0.1269 (6) | 0.0958 (3) | 0.0858 (19) |
| H1A | 0.4571 | 0.1608 | 0.1167 | 0.103* |
| H1B | 0.4885 | 0.0359 | 0.0831 | 0.103* |
| C2 | 0.6047 (3) | 0.1386 (5) | 0.1580 (3) | 0.0718 (16) |

| | | | | |
|------|-------------|-------------|-------------|-------------|
| H2A | 0.6195 | 0.2299 | 0.1700 | 0.086* |
| H2B | 0.6065 | 0.0966 | 0.2063 | 0.086* |
| C3 | 0.6818 (3) | 0.0806 (5) | 0.1350 (2) | 0.0547 (13) |
| H3A | 0.6790 | 0.1208 | 0.0860 | 0.066* |
| H3B | 0.6682 | -0.0113 | 0.1246 | 0.066* |
| C4 | 0.7839 (3) | 0.0961 (4) | 0.1974 (3) | 0.0453 (11) |
| C5 | 0.9539 (3) | 0.0312 (4) | 0.2162 (2) | 0.0410 (11) |
| C6 | 1.0875 (3) | 0.0553 (4) | 0.3442 (2) | 0.0407 (11) |
| C7 | 1.1249 (3) | -0.0633 (5) | 0.3737 (3) | 0.0522 (12) |
| C8 | 1.2206 (4) | -0.0794 (6) | 0.4252 (3) | 0.0739 (16) |
| H8 | 1.2448 | -0.1613 | 0.4445 | 0.089* |
| C9 | 1.2793 (4) | 0.0285 (7) | 0.4472 (3) | 0.0822 (19) |
| H9 | 1.3443 | 0.0198 | 0.4816 | 0.099* |
| C10 | 1.2431 (3) | 0.1484 (6) | 0.4192 (3) | 0.0786 (17) |
| H10 | 1.2835 | 0.2210 | 0.4346 | 0.094* |
| C11 | 1.1459 (3) | 0.1625 (5) | 0.3675 (3) | 0.0610 (14) |
| H11 | 1.1209 | 0.2444 | 0.3489 | 0.073* |
| C12 | -0.2143 (3) | 0.6210 (5) | -0.1206 (3) | 0.0691 (15) |
| H12A | -0.2532 | 0.6696 | -0.0977 | 0.083* |
| H12B | -0.2369 | 0.5316 | -0.1269 | 0.083* |
| C13 | -0.1093 (3) | 0.6249 (4) | -0.0646 (2) | 0.0530 (13) |
| H13A | -0.0713 | 0.5723 | -0.0865 | 0.064* |
| H13B | -0.1036 | 0.5862 | -0.0145 | 0.064* |
| C14 | -0.0659 (3) | 0.7590 (4) | -0.0487 (2) | 0.0490 (12) |
| H14A | -0.0639 | 0.7934 | -0.0977 | 0.059* |
| H14B | -0.1086 | 0.8147 | -0.0334 | 0.059* |
| C15 | 0.0351 (3) | 0.7652 (4) | 0.0149 (2) | 0.0418 (11) |
| C16 | 0.1846 (3) | 0.9006 (4) | 0.0537 (2) | 0.0386 (10) |
| C17 | 0.3156 (3) | 0.8794 (4) | 0.1821 (2) | 0.0453 (12) |
| C18 | 0.3983 (3) | 0.8244 (5) | 0.1795 (3) | 0.0566 (13) |
| C19 | 0.4902 (3) | 0.8541 (6) | 0.2342 (3) | 0.0781 (17) |
| H19 | 0.5457 | 0.8151 | 0.2319 | 0.094* |
| C20 | 0.4970 (4) | 0.9431 (7) | 0.2921 (4) | 0.091 (2) |
| H20 | 0.5584 | 0.9657 | 0.3291 | 0.109* |
| C21 | 0.4160 (4) | 0.9992 (5) | 0.2966 (3) | 0.0777 (17) |
| H21 | 0.4221 | 1.0588 | 0.3366 | 0.093* |
| C22 | 0.3243 (3) | 0.9664 (4) | 0.2409 (3) | 0.0571 (13) |
| H22 | 0.2686 | 1.0039 | 0.2438 | 0.069* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|-------------|-------------|-------------|-------------|--------------|
| Cl1 | 0.0706 (11) | 0.1612 (18) | 0.1063 (14) | 0.0426 (11) | -0.0131 (9) | 0.0101 (12) |
| F1 | 0.077 (2) | 0.0548 (19) | 0.098 (2) | 0.0034 (15) | 0.0232 (16) | 0.0093 (16) |
| F2 | 0.068 (2) | 0.111 (3) | 0.081 (2) | 0.0213 (17) | 0.0283 (16) | -0.0051 (19) |
| S1 | 0.0373 (6) | 0.0695 (9) | 0.0458 (7) | 0.0083 (6) | 0.0157 (5) | -0.0007 (6) |
| S2 | 0.0383 (6) | 0.0665 (9) | 0.0432 (7) | -0.0099 (6) | 0.0128 (5) | 0.0030 (6) |
| Cl2 | 0.0695 (9) | 0.0838 (10) | 0.0632 (9) | -0.0010 (7) | -0.0020 (7) | -0.0005 (7) |

| | | | | | | |
|-----|-------------|-----------|-----------|--------------|-------------|--------------|
| O1 | 0.0393 (18) | 0.072 (2) | 0.051 (2) | 0.0063 (15) | 0.0105 (15) | -0.0195 (17) |
| O2 | 0.0491 (19) | 0.050 (2) | 0.053 (2) | -0.0104 (15) | 0.0043 (15) | 0.0103 (16) |
| N1 | 0.0292 (19) | 0.059 (2) | 0.035 (2) | 0.0022 (17) | 0.0072 (16) | -0.0047 (16) |
| N2 | 0.0299 (19) | 0.052 (2) | 0.041 (2) | 0.0042 (16) | 0.0075 (16) | -0.0039 (17) |
| N3 | 0.032 (2) | 0.052 (2) | 0.036 (2) | -0.0023 (17) | 0.0056 (16) | 0.0003 (16) |
| N4 | 0.031 (2) | 0.054 (2) | 0.043 (2) | -0.0076 (16) | 0.0067 (16) | 0.0077 (17) |
| C1 | 0.030 (3) | 0.135 (5) | 0.086 (4) | -0.007 (3) | 0.014 (3) | -0.035 (4) |
| C2 | 0.037 (3) | 0.110 (5) | 0.065 (3) | -0.002 (3) | 0.017 (2) | -0.014 (3) |
| C3 | 0.031 (2) | 0.078 (4) | 0.051 (3) | 0.006 (2) | 0.011 (2) | -0.003 (2) |
| C4 | 0.037 (3) | 0.051 (3) | 0.045 (3) | 0.003 (2) | 0.013 (2) | -0.001 (2) |
| C5 | 0.033 (2) | 0.040 (3) | 0.045 (3) | -0.0013 (19) | 0.009 (2) | 0.003 (2) |
| C6 | 0.032 (2) | 0.053 (3) | 0.033 (2) | 0.004 (2) | 0.0073 (18) | -0.004 (2) |
| C7 | 0.046 (3) | 0.055 (3) | 0.051 (3) | 0.007 (3) | 0.013 (2) | -0.002 (2) |
| C8 | 0.058 (4) | 0.089 (4) | 0.061 (3) | 0.032 (3) | 0.007 (3) | 0.008 (3) |
| C9 | 0.034 (3) | 0.133 (6) | 0.063 (4) | 0.022 (3) | 0.001 (3) | -0.007 (4) |
| C10 | 0.036 (3) | 0.100 (5) | 0.089 (4) | -0.013 (3) | 0.011 (3) | -0.026 (3) |
| C11 | 0.047 (3) | 0.055 (3) | 0.070 (3) | 0.003 (2) | 0.010 (2) | -0.007 (3) |
| C12 | 0.050 (3) | 0.082 (4) | 0.068 (3) | -0.022 (3) | 0.015 (3) | -0.010 (3) |
| C13 | 0.046 (3) | 0.059 (3) | 0.048 (3) | -0.012 (2) | 0.011 (2) | -0.004 (2) |
| C14 | 0.032 (2) | 0.052 (3) | 0.060 (3) | 0.003 (2) | 0.014 (2) | -0.004 (2) |
| C15 | 0.032 (2) | 0.042 (3) | 0.045 (3) | 0.000 (2) | 0.008 (2) | -0.007 (2) |
| C16 | 0.028 (2) | 0.048 (3) | 0.038 (3) | -0.0008 (19) | 0.0109 (19) | -0.003 (2) |
| C17 | 0.031 (3) | 0.053 (3) | 0.043 (3) | -0.005 (2) | 0.005 (2) | 0.010 (2) |
| C18 | 0.045 (3) | 0.069 (4) | 0.053 (3) | 0.000 (3) | 0.016 (2) | 0.005 (3) |
| C19 | 0.037 (3) | 0.104 (5) | 0.082 (4) | 0.006 (3) | 0.011 (3) | 0.025 (4) |
| C20 | 0.047 (4) | 0.105 (5) | 0.089 (5) | -0.022 (4) | -0.007 (3) | 0.022 (4) |
| C21 | 0.068 (4) | 0.077 (4) | 0.068 (4) | -0.016 (3) | 0.005 (3) | -0.013 (3) |
| C22 | 0.050 (3) | 0.053 (3) | 0.058 (3) | 0.001 (2) | 0.009 (2) | 0.000 (3) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|---------|-----------|----------|-----------|
| Cl1—C1 | 1.769 (6) | C6—C7 | 1.361 (6) |
| F1—C7 | 1.347 (5) | C6—C11 | 1.365 (5) |
| F2—C18 | 1.350 (5) | C7—C8 | 1.377 (6) |
| S1—C5 | 1.654 (4) | C8—C9 | 1.372 (7) |
| S2—C16 | 1.675 (4) | C8—H8 | 0.9300 |
| Cl2—C12 | 1.791 (5) | C9—C10 | 1.364 (7) |
| O1—C4 | 1.212 (4) | C9—H9 | 0.9300 |
| O2—C15 | 1.210 (4) | C10—C11 | 1.393 (6) |
| N1—C4 | 1.374 (5) | C10—H10 | 0.9300 |
| N1—C5 | 1.388 (4) | C11—H11 | 0.9300 |
| N1—H1 | 0.8600 | C12—C13 | 1.498 (5) |
| N2—C5 | 1.336 (5) | C12—H12A | 0.9700 |
| N2—C6 | 1.433 (5) | C12—H12B | 0.9700 |
| N2—H2 | 0.8600 | C13—C14 | 1.502 (5) |
| N3—C16 | 1.375 (4) | C13—H13A | 0.9700 |
| N3—C15 | 1.380 (5) | C13—H13B | 0.9700 |
| N3—H3 | 0.8600 | C14—C15 | 1.501 (5) |

| | | | |
|------------|-----------|---------------|-----------|
| N4—C16 | 1.318 (4) | C14—H14A | 0.9700 |
| N4—C17 | 1.430 (5) | C14—H14B | 0.9700 |
| N4—H4 | 0.8600 | C17—C22 | 1.363 (6) |
| C1—C2 | 1.496 (5) | C17—C18 | 1.366 (6) |
| C1—H1A | 0.9700 | C18—C19 | 1.377 (6) |
| C1—H1B | 0.9700 | C19—C20 | 1.369 (7) |
| C2—C3 | 1.485 (6) | C19—H19 | 0.9300 |
| C2—H2A | 0.9700 | C20—C21 | 1.363 (8) |
| C2—H2B | 0.9700 | C20—H20 | 0.9300 |
| C3—C4 | 1.511 (5) | C21—C22 | 1.389 (6) |
| C3—H3A | 0.9700 | C21—H21 | 0.9300 |
| C3—H3B | 0.9700 | C22—H22 | 0.9300 |
| | | | |
| C4—N1—C5 | 129.5 (4) | C9—C10—C11 | 120.2 (5) |
| C4—N1—H1 | 115.2 | C9—C10—H10 | 119.9 |
| C5—N1—H1 | 115.2 | C11—C10—H10 | 119.9 |
| C5—N2—C6 | 121.9 (4) | C6—C11—C10 | 119.5 (5) |
| C5—N2—H2 | 119.0 | C6—C11—H11 | 120.2 |
| C6—N2—H2 | 119.0 | C10—C11—H11 | 120.2 |
| C16—N3—C15 | 128.2 (3) | C13—C12—Cl2 | 112.4 (4) |
| C16—N3—H3 | 115.9 | C13—C12—H12A | 109.1 |
| C15—N3—H3 | 115.9 | Cl2—C12—H12A | 109.1 |
| C16—N4—C17 | 122.1 (3) | C13—C12—H12B | 109.1 |
| C16—N4—H4 | 118.9 | Cl2—C12—H12B | 109.1 |
| C17—N4—H4 | 118.9 | H12A—C12—H12B | 107.8 |
| C2—C1—Cl1 | 111.9 (4) | C12—C13—C14 | 114.3 (4) |
| C2—C1—H1A | 109.2 | C12—C13—H13A | 108.7 |
| Cl1—C1—H1A | 109.2 | C14—C13—H13A | 108.7 |
| C2—C1—H1B | 109.2 | C12—C13—H13B | 108.7 |
| Cl1—C1—H1B | 109.2 | C14—C13—H13B | 108.7 |
| H1A—C1—H1B | 107.9 | H13A—C13—H13B | 107.6 |
| C3—C2—C1 | 114.2 (4) | C15—C14—C13 | 114.4 (3) |
| C3—C2—H2A | 108.7 | C15—C14—H14A | 108.7 |
| C1—C2—H2A | 108.7 | C13—C14—H14A | 108.7 |
| C3—C2—H2B | 108.7 | C15—C14—H14B | 108.7 |
| C1—C2—H2B | 108.7 | C13—C14—H14B | 108.7 |
| H2A—C2—H2B | 107.6 | H14A—C14—H14B | 107.6 |
| C2—C3—C4 | 113.7 (4) | O2—C15—N3 | 123.0 (4) |
| C2—C3—H3A | 108.8 | O2—C15—C14 | 123.2 (4) |
| C4—C3—H3A | 108.8 | N3—C15—C14 | 113.8 (4) |
| C2—C3—H3B | 108.8 | N4—C16—N3 | 117.2 (4) |
| C4—C3—H3B | 108.8 | N4—C16—S2 | 123.2 (3) |
| H3A—C3—H3B | 107.7 | N3—C16—S2 | 119.7 (3) |
| O1—C4—N1 | 123.3 (4) | C22—C17—C18 | 119.0 (4) |
| O1—C4—C3 | 124.0 (4) | C22—C17—N4 | 120.8 (4) |
| N1—C4—C3 | 112.6 (4) | C18—C17—N4 | 120.3 (4) |
| N2—C5—N1 | 115.8 (4) | F2—C18—C17 | 118.9 (4) |
| N2—C5—S1 | 124.8 (3) | F2—C18—C19 | 118.9 (5) |

| | | | |
|-----------|-----------|-------------|-----------|
| N1—C5—S1 | 119.4 (3) | C17—C18—C19 | 122.2 (5) |
| C7—C6—C11 | 119.3 (4) | C20—C19—C18 | 117.7 (5) |
| C7—C6—N2 | 120.9 (4) | C20—C19—H19 | 121.2 |
| C11—C6—N2 | 119.8 (4) | C18—C19—H19 | 121.2 |
| F1—C7—C6 | 119.2 (4) | C21—C20—C19 | 121.6 (5) |
| F1—C7—C8 | 118.6 (5) | C21—C20—H20 | 119.2 |
| C6—C7—C8 | 122.2 (5) | C19—C20—H20 | 119.2 |
| C9—C8—C7 | 118.2 (5) | C20—C21—C22 | 119.4 (5) |
| C9—C8—H8 | 120.9 | C20—C21—H21 | 120.3 |
| C7—C8—H8 | 120.9 | C22—C21—H21 | 120.3 |
| C10—C9—C8 | 120.6 (5) | C17—C22—C21 | 120.2 (5) |
| C10—C9—H9 | 119.7 | C17—C22—H22 | 119.9 |
| C8—C9—H9 | 119.7 | C21—C22—H22 | 119.9 |

Hydrogen-bond geometry (Å, °)

| D—H···A | D—H | H···A | D···A | D—H···A |
|------------------------------|------|-------|-----------|---------|
| N2—H2···O1 | 0.86 | 2.02 | 2.691 (5) | 134 |
| N2—H2···O2 ⁱ | 0.86 | 2.40 | 3.128 (5) | 143 |
| N4—H4···O2 | 0.86 | 2.02 | 2.676 (5) | 133 |
| N4—H4···O1 ⁱⁱ | 0.86 | 2.32 | 3.033 (5) | 141 |
| N3—H3···S1 ⁱⁱⁱ | 0.86 | 2.59 | 3.447 (4) | 176 |
| N1—H1···S2 ⁱⁱⁱ | 0.86 | 2.52 | 3.364 (4) | 169 |
| C3—H3A···Cl1 | 0.97 | 2.76 | 3.189 (5) | 107 |
| C14—H14A···Cl2 | 0.97 | 2.82 | 3.190 (4) | 103 |
| C14—H14A···S1 ⁱⁱⁱ | 0.97 | 2.96 | 3.784 (5) | 143 |
| C14—H14B···S2 ^{iv} | 0.97 | 2.74 | 3.691 (5) | 168 |

Symmetry codes: (i) $-x+1, y-1/2, -z+1/2$; (ii) $-x+1, y+1/2, -z+1/2$; (iii) $-x+1, -y+1, -z$; (iv) $-x, -y+2, -z$.