

Bis(4-amino-1-hexylpyridinium) bis(1,2-dicyanoethene-1,2-dithiolato)cuprate(II)

Qi Liu* and Jianlan Liu

Department of Applied Chemistry, College of Science, Nanjing University of Technology, No.5 Xinmofan Road, Nanjing, Nanjing 210009, People's Republic of China

Correspondence e-mail: liuqi198721@163.com


Received 28 May 2011; accepted 4 June 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.031; wR factor = 0.074; data-to-parameter ratio = 15.4.

The complete complex anion in the title salt, $(\text{C}_{11}\text{H}_{19}\text{N}_2)_2[\text{Cu}(\text{C}_4\text{N}_2\text{S}_2)_2]$, has $2/m$ symmetry while the complete cation is generated by mirror symmetry with the non-H atoms of the alkyl chain lying on the plane. A square-planar geometry based on an S_4 donor set is found in the anion; the $\text{Cu}-\text{S}$ distance is $2.2663(5)\text{ \AA}$. In the crystal, intermolecular $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds link the ions into layers in the bc plane comprising alternating rows of cations and anions.


Related literature

For square-planar $M[\text{dithiolene}]_2$ complexes acting as magnetic materials or showing nonlinear optical properties, see: Cassoux *et al.* (1991); Robertson & Cronin (2002).



Experimental

Crystal data



$M_r = 702.51$

Monoclinic, $C2/m$

$a = 13.3648(9)\text{ \AA}$

$b = 10.0768(4)\text{ \AA}$

$c = 13.8550(8)\text{ \AA}$

$\beta = 111.902(8)^\circ$

$V = 1731.24(17)\text{ \AA}^3$

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 0.91\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.3 \times 0.2 \times 0.1\text{ mm}$

Data collection

Siemens SMART CCD area-detector diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.939$, $T_{\max} = 1.000$

4194 measured reflections

1805 independent reflections

1299 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$

$wR(F^2) = 0.074$

$S = 0.92$

1805 reflections

117 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.22\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.17\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}2-\text{H}2\text{A}\cdots\text{N}1^i$	0.81 (2)	2.39 (2)	3.157 (2)	160 (2)

Symmetry code: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + 1$.

Data collection: *SMART* (Siemens, 1996); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Science and Technology Department of Jiangsu Province, People's Republic of China, for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2752).

References

- Cassoux, P., Valade, L., Kobayashi, H., Kobayashi, A., Clark, R. A. & Underhill, A. E. (1991). *Coord. Chem. Rev.* **110**, 115–160.
- Robertson, N. & Cronin, L. (2002). *Coord. Chem. Rev.* **227**, 93–127.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siemens (1996). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

supporting information

Acta Cryst. (2011). E67, m916 [doi:10.1107/S1600536811021611]

Bis(4-amino-1-hexylpyridinium) bis(1,2-dicyanoethene-1,2-dithiolato)cuprate(II)

Qi Liu and Jianlan Liu

S1. Comment

Square-planar $M[\text{dithiolene}]_2$ complexes have attracted extensive interest in the areas of conducting and magnetic materials, dyes, non-linear optics and catalysis (Robertson *et al.*, 2002; Cassoux *et al.*, 1991). Herein, we report the crystal structure of the title compound, Fig.1.

The $[\text{Cu}(\text{mnt})_2]^{2-}$ dianion is located about a site of symmetry $2/m$. The 1-hexyl-4-aminopyridinium cation lies on a mirror plane whereby the non-H atoms of the alkyl chain lie on the plane which bisects the 1,4 atoms of the benzene ring.


In the crystal structure, intermolecular N—H···N hydrogen bonds (Table 1) link the cations and anions to form a layer in the bc plane comprising alternating cations and anions.

S2. Experimental


Disodium maleonitriledithiobate (468 mg, 2.5 mmol) and cupric nitrate trihydrate (302 mg, 1.25 mmol) were mixed under stirring in water (20 mL) at room temperature. Subsequently, a solution of 1-hexyl-4-aminopyridinium iodide (765 mg, 2.5 mmol) in water (10 mL) was added to the mixture. The brown precipitate that formed immediately was filtered off and washed with water. The crude product was recrystallized from acetone (20 mL) to give brown crystals. The crystals suitable for X-ray diffraction measurements were obtained by diffusing diethyl ether into the solution of the salt in acetone for 6 days.

S3. Refinement

The C-bound H atoms were geometrically placed ($\text{C—H} = 0.93$ or 0.96 \AA) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2-1.5 U_{\text{eq}}(\text{C})$. The N-bound H atom was refined freely.

**Figure 1**

Molecular structure of the ions comprising (I) showing atom labelling. Displacement ellipsoids are drawn at the 50% probability level. The cation has mirror symmetry with $i: x, 1-y, z$. The Cu atom in the anion is located on a site of symmetry $2/m$. Symmetry operations $ii: -x, y, -z$ and $iii: -x, 1-y, -z$.

**Figure 2**

Partial packing view showing the layer in the bc plane. Dashed lines indicate intermolecular N—H···N hydrogen bonds.

Bis(4-amino-1-hexylpyridinium) bis(1,2-dicyanoethene-1,2-dithiolato)cuprate(II)*Crystal data*

$M_r = 702.51$

Monoclinic, $C2/m$

Hall symbol: -C 2y

$a = 13.3648 (9) \text{ \AA}$

$b = 10.0768 (4) \text{ \AA}$

$c = 13.8550 (8) \text{ \AA}$

$\beta = 111.902 (8)^\circ$

$V = 1731.24 (17) \text{ \AA}^3$

$Z = 2$

$F(000) = 734$

$D_x = 1.348 \text{ Mg m}^{-3}$

Melting point = 430–432 K

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2059 reflections

$\theta = 3.1\text{--}29.2^\circ$

$\mu = 0.91 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, brown

$0.3 \times 0.2 \times 0.1 \text{ mm}$

Data collection

Siemens SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.939$, $T_{\max} = 1.000$

4194 measured reflections

1805 independent reflections

1299 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$

$\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 3.1^\circ$

$h = -12 \rightarrow 16$

$k = -11 \rightarrow 12$

$l = -17 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.031$

$wR(F^2) = 0.074$

$S = 0.92$

1805 reflections

117 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0411P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.22 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.17 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.0000	0.5000	0.0000	0.04510 (17)
S1	0.04362 (5)	0.66007 (5)	0.12376 (4)	0.0623 (2)
N1	0.12555 (15)	0.7075 (2)	0.40284 (14)	0.0721 (6)

N2	0.3513 (2)	0.5000	0.5099 (2)	0.0615 (8)
N3	0.39247 (19)	0.5000	0.23171 (18)	0.0536 (6)
C1	0.07750 (14)	0.56692 (18)	0.23653 (14)	0.0445 (4)
C2	0.10516 (16)	0.6424 (2)	0.33068 (15)	0.0507 (5)
C3	0.3637 (2)	0.5000	0.4193 (2)	0.0472 (7)
C4	0.37096 (16)	0.38155 (19)	0.36957 (15)	0.0544 (5)
H4A	0.3666	0.3005	0.3998	0.065*
C5	0.38435 (18)	0.3850 (2)	0.27783 (17)	0.0586 (6)
H5A	0.3881	0.3054	0.2454	0.070*
C6	0.4079 (3)	0.5000	0.1312 (2)	0.0678 (9)
H6A	0.4487	0.4229	0.1280	0.081*
C7	0.3047 (3)	0.5000	0.0398 (3)	0.0810 (10)
H7A	0.2635	0.4226	0.0414	0.097*
C8	0.3282 (3)	0.5000	-0.0612 (3)	0.0819 (11)
H8A	0.3740	0.4117	-0.0657	0.098*
C9	0.2314 (3)	0.5000	-0.1571 (3)	0.0917 (12)
H9A	0.1892	0.4230	-0.1569	0.110*
C10	0.2529 (4)	0.5000	-0.2572 (3)	0.0930 (12)
H10A	0.2950	0.4229	-0.2577	0.112*
C11	0.1534 (3)	0.5000	-0.3517 (3)	0.0931 (12)
H11A	0.1716	0.5000	-0.4125	0.140*
H11B	0.1119	0.4222	-0.3519	0.140*
H2A	0.3461 (18)	0.432 (2)	0.5381 (16)	0.075 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0613 (3)	0.0335 (3)	0.0420 (3)	0.000	0.0211 (2)	0.000
S1	0.1042 (5)	0.0323 (3)	0.0463 (3)	0.0005 (3)	0.0235 (3)	0.0001 (2)
N1	0.0822 (14)	0.0679 (13)	0.0574 (11)	0.0058 (11)	0.0159 (10)	-0.0184 (10)
N2	0.076 (2)	0.0505 (19)	0.0559 (16)	0.000	0.0218 (15)	0.000
N3	0.0594 (15)	0.0435 (14)	0.0669 (15)	0.000	0.0340 (13)	0.000
C1	0.0492 (11)	0.0407 (10)	0.0423 (10)	0.0004 (9)	0.0155 (9)	-0.0020 (8)
C2	0.0538 (13)	0.0454 (11)	0.0486 (11)	0.0054 (10)	0.0143 (10)	0.0004 (10)
C3	0.0386 (16)	0.0423 (17)	0.0544 (17)	0.000	0.0100 (14)	0.000
C4	0.0649 (14)	0.0343 (11)	0.0652 (14)	0.0001 (10)	0.0257 (12)	0.0042 (10)
C5	0.0692 (15)	0.0352 (11)	0.0769 (15)	0.0012 (10)	0.0338 (13)	-0.0054 (11)
C6	0.080 (2)	0.058 (2)	0.083 (2)	0.000	0.051 (2)	0.000
C7	0.090 (3)	0.096 (3)	0.070 (2)	0.000	0.045 (2)	0.000
C8	0.092 (3)	0.087 (3)	0.079 (2)	0.000	0.046 (2)	0.000
C9	0.108 (3)	0.096 (3)	0.084 (3)	0.000	0.051 (3)	0.000
C10	0.114 (3)	0.099 (3)	0.080 (3)	0.000	0.053 (3)	0.000
C11	0.122 (3)	0.073 (3)	0.097 (3)	0.000	0.055 (3)	0.000

Geometric parameters (\AA , $^\circ$)

Cu1—S1 ⁱ	2.2663 (5)	C4—C5	1.349 (3)
Cu1—S1	2.2663 (5)	C4—H4A	0.9300

Cu1—S1 ⁱⁱ	2.2663 (5)	C5—H5A	0.9300
Cu1—S1 ⁱⁱⁱ	2.2663 (5)	C6—C7	1.483 (4)
S1—C1	1.7319 (19)	C6—H6A	0.9600
N1—C2	1.141 (2)	C7—C8	1.545 (4)
N2—C3	1.327 (4)	C7—H7A	0.9600
N2—H2A	0.80 (2)	C8—C9	1.468 (5)
N3—C5	1.347 (2)	C8—H8A	1.0943
N3—C5 ⁱⁱⁱ	1.347 (2)	C9—C10	1.519 (4)
N3—C6	1.482 (3)	C9—H9A	0.9601
C1—C1 ⁱⁱⁱ	1.349 (4)	C10—C11	1.478 (5)
C1—C2	1.434 (3)	C10—H10A	0.9600
C3—C4	1.399 (2)	C11—H11A	0.9600
C3—C4 ⁱⁱⁱ	1.399 (2)	C11—H11B	0.9600
S1 ⁱ —Cu1—S1	180.00 (2)	N3—C5—C4	122.1 (2)
S1 ⁱ —Cu1—S1 ⁱⁱ	90.75 (3)	N3—C5—H5A	118.9
S1—Cu1—S1 ⁱⁱ	89.25 (3)	C4—C5—H5A	118.9
S1 ⁱ —Cu1—S1 ⁱⁱⁱ	89.25 (3)	N3—C6—C7	113.0 (2)
S1—Cu1—S1 ⁱⁱⁱ	90.75 (3)	N3—C6—H6A	108.9
S1 ⁱⁱ —Cu1—S1 ⁱⁱⁱ	180.00 (3)	C7—C6—H6A	108.9
C1—S1—Cu1	101.76 (6)	C6—C7—C8	109.6 (3)
C3—N2—H2A	121.7 (17)	C6—C7—H7A	109.8
C5—N3—C5 ⁱⁱⁱ	118.7 (2)	C8—C7—H7A	109.5
C5—N3—C6	120.65 (12)	C9—C8—C7	114.3 (3)
C5 ⁱⁱⁱ —N3—C6	120.65 (12)	C9—C8—H8A	105.7
C1 ⁱⁱⁱ —C1—C2	122.02 (11)	C7—C8—H8A	111.0
C1 ⁱⁱⁱ —C1—S1	122.82 (6)	C8—C9—C10	115.0 (3)
C2—C1—S1	115.16 (14)	C8—C9—H9A	108.5
N1—C2—C1	176.9 (2)	C10—C9—H9A	108.3
N2—C3—C4	121.46 (13)	C11—C10—C9	113.2 (3)
N2—C3—C4 ⁱⁱⁱ	121.46 (13)	C11—C10—H10A	109.3
C4—C3—C4 ⁱⁱⁱ	117.1 (3)	C9—C10—H10A	108.5
C5—C4—C3	120.0 (2)	C10—C11—H11A	109.8
C5—C4—H4A	120.0	C10—C11—H11B	109.3
C3—C4—H4A	120.0	H11A—C11—H11B	109.5

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x, y, -z$; (iii) $x, -y+1, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
N2—H2A \cdots N1 ^{iv}	0.81 (2)	2.39 (2)	3.157 (2)	160 (2)

Symmetry code: (iv) $-x+1/2, y-1/2, -z+1$.