

## 2-Amino-6-nitro-1,3-benzothiazol-3-i um hydrogen sulfate

Hui-Fen Qian<sup>a\*</sup> and Wei Huang<sup>b</sup>

<sup>a</sup>College of Sciences, Nanjing University of Technology, Nanjing 210009, People's Republic of China, and <sup>b</sup>State Key Laboratory of Coordination Chemistry, Nanjing National Laboratory of Microstructures, School of Chemistry and Chemical Engineering, Nanjing University, Nanjing 210093, People's Republic of China  
Correspondence e-mail: qhf@njut.edu.cn


Received 8 July 2011; accepted 11 July 2011

Key indicators: single-crystal X-ray study;  $T = 291\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$ ;  $R$  factor = 0.061;  $wR$  factor = 0.159; data-to-parameter ratio = 12.0.

In the title molecular salt,  $\text{C}_7\text{H}_6\text{N}_3\text{O}_2\text{S}^+\cdot\text{HSO}_4^-$ , the 2-amino-6-nitro-1,3-benzothiazole ring system is essentially planar [mean deviation = 0.0605 (4)  $\text{\AA}$ ]. In the crystal,  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{O}-\text{H}\cdots\text{O}$  hydrogen-bonding interactions result in a layer motif.

### Related literature

For related compounds, see Glidewell *et al.* (2001); Lynch (2002); Lynch & Duckhouse (2001); You *et al.* (2009).



### Experimental

#### Crystal data

$\text{C}_7\text{H}_6\text{N}_3\text{O}_2\text{S}^+\cdot\text{HSO}_4^-$

$M_r = 293.28$

Monoclinic,  $P_{\bar{2}}1/n$

$a = 7.849 (6)\text{ \AA}$

$b = 16.219 (12)\text{ \AA}$

$c = 9.191 (7)\text{ \AA}$

$\beta = 108.584 (10)^\circ$

$V = 1109.0 (14)\text{ \AA}^3$

$Z = 4$

Mo  $K\alpha$  radiation

$\mu = 0.51\text{ mm}^{-1}$

$T = 291\text{ K}$

$0.16 \times 0.14 \times 0.12\text{ mm}$

### Data collection

Bruker 1K CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Bruker, 2000)  
 $T_{\min} = 0.923$ ,  $T_{\max} = 0.942$

5436 measured reflections  
1958 independent reflections  
1586 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.097$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$   
 $wR(F^2) = 0.159$   
 $S = 0.98$   
1958 reflections

163 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.52\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.60\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots\text{A}$	$D\cdots\text{A}$	$D-\text{H}\cdots\text{A}$
N1—H1A $\cdots$ O4 <sup>i</sup>	0.86	1.97	2.825 (4)	171
N2—H2A $\cdots$ O6 <sup>i</sup>	0.86	2.02	2.867 (4)	170
N2—H2B $\cdots$ O6 <sup>ii</sup>	0.86	2.10	2.888 (4)	151
O3—H3A $\cdots$ O4 <sup>iii</sup>	0.82	1.86	2.664 (4)	166
Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$ ; (ii) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (iii) $-x + 1, -y + 2, -z + 2$ .				

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

We would like to acknowledge the National Natural Science Foundation of China (No. 20871065) and the Jiangsu Province Department of Science and Technology (No. BK2009226) for financial aid.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FF2020).

### References

- Bruker (2000). *SMART, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Glidewell, C., Low, J. N., McWilliam, S. A., Skakle, J. M. S. & Wardell, J. L. (2001). *Acta Cryst. C57*, 1209–1211.
- Lynch, D. E. (2002). *Acta Cryst. E58*, o1139–o1141.
- Lynch, D. E. & Duckhouse, H. L. (2001). *Acta Cryst. C57*, 1036–1038.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- You, W., Fan, Y., Qian, H.-F., Yao, C. & Huang, W. (2009). *Acta Cryst. E65*, o115.

# supporting information

*Acta Cryst.* (2011). E67, o2044 [doi:10.1107/S1600536811027620]

## 2-Amino-6-nitro-1,3-benzothiazol-3-ium hydrogen sulfate

Hui-Fen Qian and Wei Huang

### S1. Comment

There have been three single-crystal structural investigations on 2-amino-6-nitro-1,3-benzothiazole, namely 2-amino-6-nitro-1,3-benzothiazole (Glidewell *et al.*, 2001), its monohydrate (Lynch *et al.*, 2002) and its PtCl<sub>2</sub> complex (Lynch & Duckhouse, 2001). We have previously reported the single-crystal structure of 2-aminobenzimidazolium hydrogen sulfate (You *et al.*, 2009). In this work, we describe the single-crystal structure of a hydrogen sulfate salt of 2-amino-6-nitro-1,3-benzothiazole.


The atom-numbering scheme of the title compound is shown in Fig. 1, while selected bond distances and bond angles are given in Table 1. The 2-amino-6-nitro-1,3-benzothiazole skeleton of the title compound is essentially planar with the mean deviation of 0.0605 (4) Å. The proton is delocalized within the thiazole ring although it is added to the nitrogen atom. With regard to the hydrogen sulfate anion, the hydrogen atom is added to the O3 atom of SO<sub>4</sub> group due to the obviously longer O3–S2 bond length. In the crystal packing, N—H···O and O—H···O hydrogen-bond interactions are found between adjacent molecules giving rise to a layer motif.

### S2. Experimental

The treatment of 2-amino-6-nitro-1,3-benzothiazole dissolved in methanol with an excess of sulfuric acid yields the title compound. Single crystals suitable for X-ray diffraction measurement were obtained after 7 days' slow evaporation of the mother liquid at room temperature in air. Anal. Calcd. For C<sub>7</sub>H<sub>6</sub>N<sub>3</sub>O<sub>2</sub>S<sup>+</sup>.HSO<sub>4</sub><sup>-</sup>: C, 28.67; H, 2.41; N, 14.33%. Found: C, 28.53; H, 2.66; N, 14.44%.

### S3. Refinement

The non-hydrogen atoms were refined anisotropically, whereas the H atoms bonded with carbon, nitrogen and oxygen atoms were placed in geometrically idealized positions (C—H = 0.93 Å, N—H = 0.86 Å and O—H = 0.82 Å) and refined as riding atoms, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  and  $1.2U_{\text{eq}}(\text{N})$  and  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ .

**Figure 1**

An *ORTEP* drawing of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

### 2-Amino-6-nitro-1,3-benzothiazol-3-ium hydrogen sulfate

#### Crystal data



$M_r = 293.28$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 7.849(6)$  Å

$b = 16.219(12)$  Å

$c = 9.191(7)$  Å

$\beta = 108.584(10)^\circ$

$V = 1109.0(14)$  Å<sup>3</sup>

$Z = 4$

$F(000) = 600$

$D_x = 1.757$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 2618 reflections

$\theta = 2.3\text{--}28.0^\circ$

$\mu = 0.51$  mm<sup>-1</sup>

$T = 291$  K

Block, colourless

$0.16 \times 0.14 \times 0.12$  mm

#### Data collection

Bruker 1K CCD area-detector

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2000)

$T_{\min} = 0.923$ ,  $T_{\max} = 0.942$

5436 measured reflections

1958 independent reflections

1586 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.097$

$\theta_{\max} = 25.0^\circ$ ,  $\theta_{\min} = 2.5^\circ$

$h = -9 \rightarrow 6$

$k = -19 \rightarrow 18$

$l = -8 \rightarrow 10$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.061$

$wR(F^2) = 0.159$

$S = 0.98$

1958 reflections

163 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.1126P)^2]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.52 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.60 \text{ e } \text{\AA}^{-3}$

### Special details

**Experimental.** The structure was solved by direct methods (Bruker, 2000) and successive difference Fourier syntheses.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7190 (4)	0.12918 (17)	0.4541 (3)	0.0400 (7)
C2	0.7416 (4)	-0.00938 (16)	0.5027 (3)	0.0379 (7)
C3	0.7086 (5)	-0.09352 (18)	0.4792 (4)	0.0483 (8)
H3	0.6294	-0.1131	0.3874	0.058*
C4	0.7959 (5)	-0.14661 (19)	0.5950 (4)	0.0501 (8)
H4	0.7790	-0.2032	0.5815	0.060*
C5	0.9089 (4)	-0.11571 (17)	0.7315 (4)	0.0427 (7)
C6	0.9441 (4)	-0.03251 (17)	0.7588 (3)	0.0427 (7)
H6	1.0207	-0.0132	0.8520	0.051*
C7	0.8588 (4)	0.02045 (17)	0.6393 (3)	0.0387 (7)
N1	0.6668 (3)	0.05352 (15)	0.4010 (3)	0.0412 (6)
H1A	0.5927	0.0447	0.3107	0.049*
N2	0.6630 (4)	0.19655 (15)	0.3774 (3)	0.0524 (8)
H2A	0.5874	0.1941	0.2861	0.063*
H2B	0.7014	0.2436	0.4178	0.063*
N3	0.9911 (4)	-0.17264 (18)	0.8562 (3)	0.0538 (7)
O1	1.0531 (3)	-0.14566 (16)	0.9864 (3)	0.0651 (7)
O2	0.9914 (5)	-0.24573 (18)	0.8252 (3)	0.0885 (10)
O3	0.4172 (3)	0.88694 (13)	1.0011 (3)	0.0602 (7)
H3A	0.3934	0.9305	1.0357	0.090*
O4	0.6040 (3)	0.96348 (13)	0.8861 (2)	0.0479 (6)
O5	0.7360 (3)	0.90117 (14)	1.1335 (3)	0.0607 (7)
O6	0.6226 (3)	0.81597 (13)	0.9086 (3)	0.0549 (7)
S1	0.87176 (11)	0.12757 (5)	0.63686 (9)	0.0506 (3)
S2	0.60859 (9)	0.89227 (4)	0.98544 (8)	0.0401 (3)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0477 (16)	0.0359 (15)	0.0280 (16)	-0.0021 (12)	0.0003 (13)	0.0006 (12)
C2	0.0509 (16)	0.0330 (14)	0.0274 (15)	-0.0023 (12)	0.0092 (13)	0.0000 (11)

C3	0.067 (2)	0.0398 (16)	0.0321 (16)	-0.0075 (15)	0.0077 (14)	-0.0101 (13)
C4	0.072 (2)	0.0347 (15)	0.0430 (19)	0.0010 (15)	0.0172 (16)	0.0014 (14)
C5	0.0461 (16)	0.0434 (17)	0.0365 (17)	0.0047 (13)	0.0103 (14)	0.0073 (13)
C6	0.0451 (16)	0.0452 (16)	0.0315 (16)	-0.0033 (13)	0.0034 (13)	0.0035 (13)
C7	0.0449 (15)	0.0351 (15)	0.0312 (15)	-0.0024 (12)	0.0053 (12)	-0.0006 (11)
N1	0.0530 (14)	0.0383 (13)	0.0239 (12)	-0.0053 (11)	0.0005 (10)	-0.0023 (10)
N2	0.0697 (18)	0.0341 (13)	0.0363 (15)	-0.0036 (12)	-0.0073 (13)	-0.0003 (11)
N3	0.0579 (16)	0.0541 (17)	0.0479 (19)	0.0046 (13)	0.0149 (14)	0.0168 (14)
O1	0.0650 (15)	0.0766 (18)	0.0433 (16)	-0.0019 (13)	0.0025 (12)	0.0191 (13)
O2	0.131 (3)	0.0496 (15)	0.073 (2)	0.0216 (16)	0.0161 (17)	0.0176 (14)
O3	0.0525 (14)	0.0479 (13)	0.081 (2)	-0.0029 (10)	0.0230 (14)	-0.0050 (12)
O4	0.0656 (14)	0.0382 (11)	0.0354 (12)	0.0087 (10)	0.0099 (10)	0.0037 (9)
O5	0.0633 (15)	0.0678 (16)	0.0346 (13)	-0.0024 (11)	-0.0077 (11)	-0.0017 (11)
O6	0.0696 (15)	0.0379 (12)	0.0450 (14)	0.0118 (10)	0.0013 (11)	-0.0044 (10)
S1	0.0641 (6)	0.0364 (5)	0.0329 (5)	-0.0078 (3)	-0.0104 (4)	-0.0014 (3)
S2	0.0452 (5)	0.0362 (5)	0.0297 (5)	0.0029 (3)	-0.0010 (4)	-0.0023 (3)

Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )

C1—N2	1.299 (4)	C6—H6	0.9300
C1—N1	1.336 (3)	C7—S1	1.741 (3)
C1—S1	1.725 (3)	N1—H1A	0.8600
C2—N1	1.382 (3)	N2—H2A	0.8600
C2—C7	1.386 (4)	N2—H2B	0.8600
C2—C3	1.393 (4)	N3—O1	1.220 (4)
C3—C4	1.370 (4)	N3—O2	1.220 (4)
C3—H3	0.9300	O3—S2	1.557 (3)
C4—C5	1.380 (5)	O3—H3A	0.8200
C4—H4	0.9300	O4—S2	1.466 (2)
C5—C6	1.384 (4)	O5—S2	1.416 (2)
C5—N3	1.453 (4)	O6—S2	1.446 (2)
C6—C7	1.387 (4)		
N2—C1—N1	124.2 (3)	C2—C7—S1	111.2 (2)
N2—C1—S1	123.5 (2)	C6—C7—S1	127.8 (2)
N1—C1—S1	112.3 (2)	C1—N1—C2	114.6 (2)
N1—C2—C7	111.8 (2)	C1—N1—H1A	122.7
N1—C2—C3	126.9 (3)	C2—N1—H1A	122.7
C7—C2—C3	121.3 (3)	C1—N2—H2A	120.0
C4—C3—C2	118.2 (3)	C1—N2—H2B	120.0
C4—C3—H3	120.9	H2A—N2—H2B	120.0
C2—C3—H3	120.9	O1—N3—O2	123.3 (3)
C3—C4—C5	119.7 (3)	O1—N3—C5	118.9 (3)
C3—C4—H4	120.2	O2—N3—C5	117.8 (3)
C5—C4—H4	120.2	S2—O3—H3A	109.5
C4—C5—C6	123.5 (3)	C1—S1—C7	90.14 (13)
C4—C5—N3	118.8 (3)	O5—S2—O6	114.58 (14)
C6—C5—N3	117.6 (3)	O5—S2—O4	112.84 (14)

C5—C6—C7	116.2 (3)	O6—S2—O4	111.16 (15)
C5—C6—H6	121.9	O5—S2—O3	108.89 (16)
C7—C6—H6	121.9	O6—S2—O3	102.90 (14)
C2—C7—C6	121.0 (3)	O4—S2—O3	105.57 (13)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1A···O4 <sup>i</sup>	0.86	1.97	2.825 (4)	171
N2—H2A···O6 <sup>i</sup>	0.86	2.02	2.867 (4)	170
N2—H2B···O6 <sup>ii</sup>	0.86	2.10	2.888 (4)	151
O3—H3A···O4 <sup>iii</sup>	0.82	1.86	2.664 (4)	166

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $-x+3/2, y-1/2, -z+3/2$ ; (iii)  $-x+1, -y+2, -z+2$ .