

## Bis(acetonitrile- $\kappa N$ )diaquabis-(perchlorato- $\kappa O$ )copper(II)

Viktor A. Tafeenko, Stanislav I. Gurskiy and Leonid A. Aslanov\*

Chemistry Department, Moscow State University, 119991 Moscow, Russian Federation

Correspondence e-mail: Aslanov@struct.chem.msu.ru


Received 27 May 2011; accepted 7 July 2011

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(C-C) = 0.011$  Å; disorder in main residue;  $R$  factor = 0.066;  $wR$  factor = 0.192; data-to-parameter ratio = 9.1.

In the title compound,  $[Cu(ClO_4)_2(CH_3CN)_2(H_2O)_2]$ , the  $Cu^{2+}$  ion, located on a special position (site symmetry  $\bar{1}$ ), is coordinated by six monodentate ligands, *viz.* an  $N$ -coordinated acetonitrile, a perchlorate anion and a water molecule, and their symmetry-related counterparts. The perchlorate anion is disordered over two sets of sites with occupancies of 0.53 (2) and 0.47 (2). The crystal structure is stabilized by  $O-H \cdots O$  hydrogen bonds involving the perchlorate ion and aqua H atoms.

### Related literature

For details of the changing Cu(II/I) redox potential with increasing acetonitrile contents in water–acetonitrile solution, see: Cox *et al.* (1988); Verma & Sood (1979); Sumalekshmy & Gopidas (2005); Ajayakumar *et al.* (2009); Drew *et al.* (1985). For the dependence of the luminescent properties (emission energy) of the 3-cyano-4-dicyanomethylene-5-oxo-4,5-dihydro-1H-pyrrol-2-olate (*A*)-based salts depend on the molecular environment around (*A*), see: Tafeenko *et al.* (2009, 2010). For transition metals as fluorescence quenchers, see: Xu *et al.* (2005, 2010). For a previous study on the formation of related compounds, see: Inamo *et al.* (2001).



### Experimental

#### Crystal data

|                                    |  |
|------------------------------------|--|
| $[Cu(ClO_4)_2(C_2H_3N)_2(H_2O)_2]$ | $\gamma = 77.12 (1)^\circ$               |
| $M_r = 380.59$                     | $V = 334.1 (1) \text{ \AA}^3$            |
| Triclinic, $P\bar{1}$              | $Z = 1$                                  |
| $a = 5.581 (1) \text{ \AA}$        | Ag $K\alpha$ radiation                   |
| $b = 7.244 (2) \text{ \AA}$        | $\lambda = 0.56085 \text{ \AA}$          |
| $c = 8.733 (2) \text{ \AA}$        | $\mu = 1.09 \text{ mm}^{-1}$             |
| $\alpha = 82.82 (2)^\circ$         | $T = 296 \text{ K}$                      |
| $\beta = 76.86 (1)^\circ$          | $0.15 \times 0.1 \times 0.08 \text{ mm}$ |

#### Data collection

|                                   |                                      |
|-----------------------------------|--------------------------------------|
| Enraf–Nonius CAD-4 diffractometer | 1020 reflections with $I > 2s(I)$    |
| 2518 measured reflections         | $R_{int} = 0.049$                    |
| 1259 independent reflections      | 2 standard reflections every 120 min |
|                                   | intensity decay: none                |

#### Refinement

|                                 |  |
|---------------------------------|--|
| $R[F^2 > 2\sigma(F^2)] = 0.066$ | H atoms treated by a mixture of independent and constrained refinement |
| $wR(F^2) = 0.192$               |  |
| $S = 1.08$                      |  |
| 1259 reflections                | $\Delta\rho_{max} = 0.70 \text{ e \AA}^{-3}$                           |
| 138 parameters                  | $\Delta\rho_{min} = -1.09 \text{ e \AA}^{-3}$                          |
| 11 restraints                   |  |

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H \cdots A$          | $D-H$    | $H \cdots A$ | $D \cdots A$ | $D-H \cdots A$ |
|-------------------------|----------|--------------|--------------|----------------|
| $O1-H11 \cdots O3^i$    | 0.83 (7) | 2.03 (7)     | 2.758 (17)   | 146 (6)        |
| $O1-H12 \cdots O3^{ii}$ | 0.79 (9) | 2.22 (10)    | 3.00 (3)     | 165 (8)        |

Symmetry codes: (i)  $x, y-1, z$ ; (ii)  $-x+1, -y+1, -z+1$ .

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2000); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FI2110).

### References

- Ajayakumar, G., Sreenath, K. & Gopidas, K. R. (2009). *Dalton Trans.* pp. 1180–1186.
- Brandenburg, K. (2000). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Cox, B. G., Jedral, W. & Palou, J. (1988). *J. Chem. Soc. Dalton Trans.* pp. 733–740.
- Drew, M. G. B., Yates, P. C., Trocha-Grimshaw, J., McKillop, K. P. & Nelson, S. M. (1985). *J. Chem. Soc. Chem. Commun.* pp. 262–263.
- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Inamo, M., Kamiya, N., Inada, Y., Nomura, M. & Funahashi, S. (2001). *Inorg. Chem.* **40**, 5636–5641.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sumalekshmy, S. & Gopidas, K. R. (2005). *Chem. Phys. Lett.* **413**, 294–299.
- Tafeenko, V. A., Gurskiy, S. I., Baranov, A. N., Kaisarova, T. V. & Aslanov, L. A. (2009). *Acta Cryst.* **C65**, m52–m55.

## metal-organic compounds

---

Tafeenko, V. A., Gurskiy, S. I., Fazyzbekov, M. F., Baranov, A. N. & Aslanov, L. A. (2010). *Acta Cryst.* **C66**, m32–m34.  
Verma, B. C. & Sood, R. K. (1979). *Talanta*, **26**, 906–907.

Xu, Z., Han, S. J., Lee, C., Yoon, J. & Spring, D. R. (2010). *Chem. Commun.* **46**, 1679–1681.  
Xu, Z., Xiao, V., Qian, X., Cui, J. & Cui, D. (2005). *Org. Lett.* **7**, 889–892.

## supporting information

*Acta Cryst.* (2011). E67, m1125–m1126 [doi:10.1107/S1600536811027309]

**Bis(acetonitrile- $\kappa$ N)diaquabis(perchlorato- $\kappa$ O)copper(II)**

**Viktor A. Tafeenko, Stanislav I. Gurskiy and Leonid A. Aslanov**

**S1. Comment**

It was found (Tafeenko *et al.*, 2009; Tafeenko *et al.*, 2010) that the luminescent properties (emission energy) of the 3-cyano-4-dicyanomethylene-5-oxo-4,5-dihydro-1*H*-pyrrol-2-olate (A)-based salts depend on the molecular environment around (A). To investigate the effect of transition metals, which act usually as fluorescence quenchers (Xu *et al.*, 2005, 2010), we made an attempt to synthesize (A)-based salts with Cu<sup>2+</sup>. Acetonitrile treatment of crude crystalline mass, obtained after evaporation of CuSO<sub>4</sub> + BaA<sub>2</sub> (equivalent amounts) mixture in water-ethanol (*v/v*=1/1) solution, and posterior evaporation of an acetonitrile solution in an air atmosphere at room temperature resulted in formation of (A)-based salts with Cu<sup>+</sup>. We saw several reasons for the reduction of Cu<sup>2+</sup> to Cu<sup>+</sup>:

- iodide anion I<sup>-</sup> reduces Cu<sup>2+</sup> in water-ethanol solution, as it was detected that BaA<sub>2</sub> salt used for synthesis contained I<sup>-</sup>;
- anion (A) exhibits of reducing properties;
- acetonitrile-water mixture causes reduction of copper(II) to copper(I), as it was documented that i) copper(II) salt solution in acetonitrile-water mixture is a powerful oxidizer of organic molecules (Verma *et al.*, 1979; Cox *et al.*, 1988; Sumalekshmy *et al.*, 2005; Ajayakumar *et al.*, 2009); ii) acetonitrile can reduce Cu<sup>2+</sup> to Cu<sup>+</sup> (Drew *et al.*, 1985).

To find out whether acetonitrile-water mixture can cause a reduction of Cu<sup>2+</sup> to Cu<sup>+</sup>, we prepared a Cu(ClO<sub>4</sub>)<sub>2</sub> solution in a mixture of 99.5% CH<sub>3</sub>CN and 0.5% H<sub>2</sub>O (volume percentage) and evaporated this solution, which yielded crystals of the title compound, [Cu(CH<sub>3</sub>CN)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>(ClO<sub>4</sub>)<sub>2</sub>]. No other phases could be detected using powder-XRD. The crystal and molecular structure of the title compound (Fig.1) is presented in this paper.

The structure is composed of monomeric units built up around a Cu<sup>2+</sup> on a special position (site symmetry  $\bar{1}$ ). The Cu<sup>2+</sup> cation is surrounded by six monodentate ligands, *viz.* an N-coordinated acetonitrile, a perchlorate anion and a water molecule, and their symmetry related counterparts. The perchlorate anion is disordered over two positions, with occupancies (0.53 (2) and 0.47 (2)), but it's O atoms displacement ellipsoids are still quite large, indicating possible rotational disorder, with the rotation axis passing through oxygen O1 of the perchlorate coordinated with Cu1 and Cl1. The complex adopts an elongated octahedral coordination geometry.

The axial Cu–O1 (perchlorate) bond length is 2.401 (15) and in plane Cu–N1(acetonitrile) 1.960 (5), Cu1–O2(aqua) 1.950 (5) Å respectively.

Besides ionic forces, the crystal structure is stabilized by hydrogen bonding interaction *via* the perchlorate and aqua H atoms (Fig. 2).

In conclusion, we have to note that the structure of the title compound differs from reported by Inamo with co-workers (Inamo *et al.*, 2001) compounds. They reported the formation of [Cu(H<sub>2</sub>O)<sub>n</sub>(CH<sub>3</sub>CN)<sub>(6-n)</sub>]<sup>2+</sup> (*n* = 0–3) in (water/acetonitrile) solutions with [Cu(H<sub>2</sub>O)<sub>m</sub>(CH<sub>3</sub>CN)<sub>(6-m)</sub>]<sup>2+</sup> (*m* < 3) as the dominant species for H<sub>2</sub>O concentration lower than 0.5 M (99% acetonitrile).


## S2. Experimental

Blue crystals of the title salt were obtained by slow evaporation of 0.045 M  $\text{Cu}(\text{H}_2\text{O})_6(\text{ClO}_4)_2$  solution in acetonitrile at room temperature in an air atmosphere. The crystals are not stable in the open air, so suitable for X-ray investigation crystal was placed in a sealed capillary. Copper(II) perchlorate hexahydrate of 98% (Aldrich) grade was used for synthesis. Acetonitrile was boiled with phosphorus pentoxide and then distilled at 353 K.

## S3. Refinement


During the refinement a difference maps showed peaks consistent with the perchlorate atoms Cl1, O2—O5 being unequally disorder over two interpenetrating sites. This was allowed for by use of the appropriate *SHELXL* SAME, EADP restraints. At convergence the perchlorate disorder was modelled with occupancies (0.53 (2) and 0.47 (2)).

The positions of the H atoms of the water molecule were determined from Fourier difference maps and refined freely; the positions of the H atoms of the methyl group were placed in calculated positions and allowed to ride on their parent atoms [ $\text{C—H} = 0.96 \text{ \AA}$ ].  $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{parent atom})$ , where  $x = 1.5$  for attached C atoms.



**Figure 1**

The molecular structure of the title compound, with displacement ellipsoids drawn at the 50% probability level. Only one of the two disordered positions of the perchlorate ion is shown. Symmetry code: (i)  $-x, 1 - y, -z$ .

**Figure 2**

The hydrogen bonding pattern in the title compound. H bonds are drawn as dashed lines. Symmetry codes (ii)  $x, 1-y, z$ ; (iii)  $1-x, 1-y, 1-z$ ; (iv)  $1-x, -y, 1-z$ .

### Bis(acetonitrile- $\kappa$ N)diaquabis(perchlorato- $\kappa$ O)copper(II)

#### Crystal data

$[\text{Cu}(\text{ClO}_4)_2(\text{C}_2\text{H}_3\text{N})_2(\text{H}_2\text{O})_2]$

$M_r = 380.59$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 5.581\ (1)\ \text{\AA}$

$b = 7.244\ (2)\ \text{\AA}$

$c = 8.733\ (2)\ \text{\AA}$

$\alpha = 82.82\ (2)^\circ$

$\beta = 76.86\ (1)^\circ$

$\gamma = 77.12\ (1)^\circ$

$V = 334.1\ (1)\ \text{\AA}^3$

$Z = 1$

$F(000) = 191$

$D_x = 1.891\ \text{Mg m}^{-3}$

Melting point: 422 K

Ag  $K\alpha$  radiation,  $\lambda = 0.56085\ \text{\AA}$

Cell parameters from 25 reflections

$\theta = 11\text{--}13^\circ$

$\mu = 1.09\ \text{mm}^{-1}$

$T = 296\ \text{K}$

Prism, light-blue

$0.15 \times 0.1 \times 0.08\ \text{mm}$

*Data collection*

|  |  |
|--|--|
| Enraf–Nonius CAD-4<br>diffractometer     | $R_{\text{int}} = 0.049$   |
| Radiation source: fine-focus sealed tube | $\theta_{\text{max}} = 20.0^\circ$ , $\theta_{\text{min}} = 1.9^\circ$ |
| Graphite monochromator                   | $h = -6 \rightarrow 6$   |
| non-profiled $\omega$ scans              | $k = -8 \rightarrow 8$   |
| 2518 measured reflections                | $l = -10 \rightarrow 10$   |
| 1259 independent reflections             | 2 standard reflections every 120 min                                   |
| 1020 reflections with $I > 2\sigma(I)$   | intensity decay: none  |

*Refinement*

|  |  |
|--|--|
| Refinement on $F^2$  | Secondary atom site location: difference Fourier map                   |
| Least-squares matrix: full                                     | Hydrogen site location: inferred from neighbouring sites               |
| $R[F^2 > 2\sigma(F^2)] = 0.066$                                | H atoms treated by a mixture of independent and constrained refinement |
| $wR(F^2) = 0.192$  | $w = 1/[\sigma^2(F_o^2) + (0.1161P)^2 + 0.4339P]$                      |
| $S = 1.08$   | where $P = (F_o^2 + 2F_c^2)/3$   |
| 1259 reflections   | $(\Delta/\sigma)_{\text{max}} = 0.042$                                 |
| 138 parameters   | $\Delta\rho_{\text{max}} = 0.70 \text{ e } \text{\AA}^{-3}$            |
| 11 restraints  | $\Delta\rho_{\text{min}} = -1.09 \text{ e } \text{\AA}^{-3}$           |
| Primary atom site location: structure-invariant direct methods |  |

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

|      | $x$         | $y$         | $z$         | $U_{\text{iso}}^*/U_{\text{eq}}$ | Occ. (<1) |
|------|-------------|-------------|-------------|----------------------------------|-----------|
| Cu1  | 0.0000      | 0.5000      | 0.5000      | 0.0477 (4)                       |           |
| O1   | 0.2556 (10) | 0.2741 (6)  | 0.5295 (6)  | 0.0574 (12)                      |           |
| N1   | 0.2362 (12) | 0.5987 (8)  | 0.3237 (7)  | 0.0631 (16)                      |           |
| C1   | 0.3717 (13) | 0.6605 (9)  | 0.2302 (8)  | 0.0555 (15)                      |           |
| C2   | 0.5546 (16) | 0.7387 (12) | 0.1089 (9)  | 0.072 (2)                        |           |
| H2A  | 0.7197      | 0.6899      | 0.1297      | 0.109*                           |           |
| H2B  | 0.5174      | 0.8747      | 0.1088      | 0.109*                           |           |
| H2C  | 0.5478      | 0.7034      | 0.0079      | 0.109*                           |           |
| Cl1  | 0.120 (3)   | 0.803 (2)   | 0.7550 (16) | 0.0538 (10)                      | 0.53 (2)  |
| O2   | 0.195 (3)   | 0.656 (3)   | 0.651 (2)   | 0.082 (6)                        | 0.53 (2)  |
| O3   | 0.295 (5)   | 0.921 (3)   | 0.694 (3)   | 0.142 (9)                        | 0.53 (2)  |
| O4   | 0.143 (3)   | 0.729 (3)   | 0.9072 (14) | 0.105 (6)                        | 0.53 (2)  |
| O5   | -0.124 (3)  | 0.902 (2)   | 0.755 (2)   | 0.096 (5)                        | 0.53 (2)  |
| Cl11 | 0.132 (3)   | 0.820 (2)   | 0.7453 (18) | 0.0538 (10)                      | 0.47 (2)  |
| O21  | 0.112 (5)   | 0.637 (2)   | 0.717 (2)   | 0.079 (6)                        | 0.47 (2)  |

|     |            |            |            |             |          |
|-----|------------|------------|------------|-------------|----------|
| O31 | 0.383 (2)  | 0.837 (3)  | 0.729 (2)  | 0.080 (5)   | 0.47 (2) |
| O41 | 0.008 (6)  | 0.853 (6)  | 0.899 (3)  | 0.22 (2)    | 0.47 (2) |
| O51 | 0.017 (6)  | 0.955 (2)  | 0.644 (4)  | 0.156 (12)  | 0.47 (2) |
| H11 | 0.212 (12) | 0.172 (10) | 0.565 (7)  | 0.041 (16)* |          |
| H12 | 0.357 (17) | 0.228 (12) | 0.458 (10) | 0.07 (3)*   |          |

*Atomic displacement parameters (Å<sup>2</sup>)*

|      | $U^{11}$    | $U^{22}$   | $U^{33}$    | $U^{12}$     | $U^{13}$    | $U^{23}$     |
|------|-------------|------------|-------------|--------------|-------------|--------------|
| Cu1  | 0.0492 (7)  | 0.0344 (6) | 0.0530 (7)  | -0.0110 (4)  | 0.0021 (4)  | 0.0024 (4)   |
| O1   | 0.063 (3)   | 0.034 (2)  | 0.064 (3)   | -0.004 (2)   | 0.001 (2)   | 0.002 (2)    |
| N1   | 0.060 (3)   | 0.045 (3)  | 0.068 (3)   | -0.013 (3)   | 0.014 (3)   | 0.008 (2)    |
| C1   | 0.057 (4)   | 0.042 (3)  | 0.060 (4)   | -0.006 (3)   | -0.005 (3)  | 0.002 (3)    |
| C2   | 0.073 (5)   | 0.075 (5)  | 0.060 (4)   | -0.028 (4)   | 0.005 (4)   | 0.016 (3)    |
| C11  | 0.0544 (13) | 0.040 (2)  | 0.0630 (16) | -0.0150 (12) | 0.0034 (11) | -0.0097 (13) |
| O2   | 0.062 (9)   | 0.085 (11) | 0.107 (13)  | 0.011 (7)    | -0.033 (9)  | -0.051 (10)  |
| O3   | 0.115 (16)  | 0.087 (11) | 0.23 (2)    | -0.061 (11)  | -0.042 (14) | 0.062 (13)   |
| O4   | 0.093 (11)  | 0.152 (15) | 0.046 (6)   | -0.002 (9)   | 0.000 (6)   | 0.013 (7)    |
| O5   | 0.072 (8)   | 0.088 (11) | 0.110 (11)  | 0.018 (7)    | -0.007 (7)  | -0.028 (8)   |
| C111 | 0.0544 (13) | 0.040 (2)  | 0.0630 (16) | -0.0150 (12) | 0.0034 (11) | -0.0097 (13) |
| O21  | 0.111 (17)  | 0.047 (7)  | 0.090 (12)  | -0.027 (9)   | -0.039 (10) | 0.003 (7)    |
| O31  | 0.044 (7)   | 0.117 (14) | 0.085 (9)   | -0.034 (8)   | 0.006 (6)   | -0.028 (9)   |
| O41  | 0.18 (3)    | 0.31 (4)   | 0.17 (3)    | -0.16 (3)    | 0.12 (2)    | -0.15 (3)    |
| O51  | 0.19 (3)    | 0.041 (8)  | 0.26 (3)    | -0.028 (10)  | -0.11 (2)   | 0.040 (11)   |

*Geometric parameters (Å, °)*

|                                      |             |              |            |
|--------------------------------------|-------------|--------------|------------|
| Cu1—O1                               | 1.950 (5)   | C2—H2C       | 0.9600     |
| Cu1—N1                               | 1.960 (5)   | C11—O5       | 1.390 (14) |
| Cu1—O2                               | 2.401 (15)  | C11—O4       | 1.391 (14) |
| O1—H11                               | 0.83 (7)    | C11—O3       | 1.414 (14) |
| O1—H12                               | 0.79 (9)    | C11—O2       | 1.418 (12) |
| N1—C1                                | 1.103 (9)   | C111—O51     | 1.386 (15) |
| C1—C2                                | 1.450 (9)   | C111—O41     | 1.387 (15) |
| C2—H2A                               | 0.9600      | C111—O31     | 1.409 (14) |
| C2—H2B                               | 0.9600      | C111—O21     | 1.413 (13) |
| O1—Cu1—O1 <sup>i</sup>               | 180.000 (1) | C1—C2—H2C    | 109.5      |
| O1—Cu1—N1                            | 90.4 (2)    | H2A—C2—H2C   | 109.5      |
| O1 <sup>i</sup> —Cu1—N1              | 89.6 (2)    | H2B—C2—H2C   | 109.5      |
| N1—Cu1—N1 <sup>i</sup>               | 180.0 (3)   | O5—C11—O4    | 110.2 (13) |
| O1—Cu1—O2 <sup>i</sup>               | 93.3 (4)    | O5—C11—O3    | 110.9 (15) |
| N1—Cu1—O2 <sup>i</sup>               | 97.7 (6)    | O4—C11—O3    | 110.2 (14) |
| N1 <sup>i</sup> —Cu1—O2 <sup>i</sup> | 82.3 (6)    | O5—C11—O2    | 112.1 (11) |
| O1—Cu1—O2                            | 86.7 (4)    | O4—C11—O2    | 110.3 (13) |
| O2 <sup>i</sup> —Cu1—O2              | 180.0 (7)   | O3—C11—O2    | 102.9 (14) |
| Cu1—O1—H11                           | 120 (4)     | C11—O2—Cu1   | 137.3 (11) |
| Cu1—O1—H12                           | 123 (6)     | O51—C111—O41 | 109 (2)    |

|            |           |              |            |
|------------|-----------|--------------|------------|
| H11—O1—H12 | 93 (7)    | O51—C111—O31 | 109.9 (15) |
| C1—N1—Cu1  | 176.0 (7) | O41—C111—O31 | 108.2 (15) |
| N1—C1—C2   | 178.6 (8) | O51—C111—O21 | 109.8 (13) |
| C1—C2—H2A  | 109.5     | O41—C111—O21 | 107.7 (16) |
| C1—C2—H2B  | 109.5     | O31—C111—O21 | 112.3 (14) |
| H2A—C2—H2B | 109.5     |              |            |

Symmetry code: (i)  $-x, -y+1, -z+1$ .

*Hydrogen-bond geometry (Å, °)*

| <i>D</i> —H $\cdots$ <i>A</i>     | <i>D</i> —H | H $\cdots$ <i>A</i> | <i>D</i> $\cdots$ <i>A</i> | <i>D</i> —H $\cdots$ <i>A</i> |
|-----------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| O1—H11 $\cdots$ O3 <sup>ii</sup>  | 0.83 (7)    | 2.03 (7)            | 2.758 (17)                 | 146 (6)                       |
| O1—H12 $\cdots$ O3 <sup>iii</sup> | 0.79 (9)    | 2.22 (10)           | 3.00 (3)                   | 165 (8)                       |

Symmetry codes: (ii)  $x, y-1, z$ ; (iii)  $-x+1, -y+1, -z+1$ .