

## 1,1,2,2-Tetrakis(1,3-benzoxazol-2-yl)-ethene

Tesfamariam K. Hagos,<sup>a</sup> Stefan D. Nogai,<sup>a</sup> Liliana Dobrzańska,<sup>a,b\*</sup> Stephanie Cronje<sup>a,c</sup> and Helgard G. Raubenheimer<sup>a</sup>

<sup>a</sup>Department of Chemistry, University of Stellenbosch, Private Bag X1, Matieland, South Africa, <sup>b</sup>Department of Chemistry, Katholieke Universiteit Leuven, Celestijnenlaan 200F - bus 2404, B-3001 Heverlee, Belgium, and <sup>c</sup>Institut für Anorganische und Analytische Chemie, Goethe-Universität Frankfurt, Max-von-Laue-Strasse 7, D-60348 Frankfurt am Main, Germany  
Correspondence e-mail: lianger@sun.ac.za


Received 1 July 2011; accepted 5 July 2011

Key indicators: single-crystal X-ray study;  $T = 273\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.049;  $wR$  factor = 0.121; data-to-parameter ratio = 15.8.

The title compound,  $\text{C}_{30}\text{H}_{16}\text{N}_4\text{O}_4$ , reveals  $\overline{1}$  crystallographic and molecular symmetry and accordingly the asymmetric unit comprises one half-molecule. The dihedral angle between the planes of the two geminal benzoxazole rings is  $74.39(5)^\circ$ . The packing features weak  $\text{C}-\text{H}\cdots\text{N}$  and  $\pi-\pi$  interactions [centroid–centroid distance =  $3.652(1)\text{ \AA}$ ].

### Related literature

For the chloroform solvate of 1,1,2,2-tetrakis(1,3-benzothiazol-2-yl)ethene, see: Hagos *et al.* (2010). For bond lengths in the benzoxazole moiety in related compounds, see: Jian *et al.* (2007); Lokaj *et al.* (1997); Muir *et al.* (1992). For details of the cut-off applied for  $\text{C}-\text{H}\cdots\text{N}$  interactions, see: Desiraju & Steiner (2006). For the synthesis of  $\text{AuCl}(\text{PPh}_3)$ , see: Bruce *et al.* (1989).



### Experimental

#### Crystal data

$\text{C}_{30}\text{H}_{16}\text{N}_4\text{O}_4$

$M_r = 496.47$

Monoclinic,  $P2_1/c$   
 $a = 9.2697(9)\text{ \AA}$   
 $b = 16.1943(16)\text{ \AA}$   
 $c = 8.0332(8)\text{ \AA}$   
 $\beta = 104.395(2)^\circ$   
 $V = 1168.1(2)\text{ \AA}^3$

$Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.10\text{ mm}^{-1}$   
 $T = 273\text{ K}$   
 $0.30 \times 0.25 \times 0.15\text{ mm}$

#### Data collection

Bruker APEX CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1997)  
 $T_{\min} = 0.972$ ,  $T_{\max} = 0.986$

6999 measured reflections  
2717 independent reflections  
2458 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.121$   
 $S = 1.06$   
2717 reflections

172 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.37\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.30\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C7—H7 $\cdots$ N12 <sup>i</sup>	0.93	2.71	3.348 (2)	127
C17—H17 $\cdots$ N3 <sup>ii</sup>	0.93	2.71	3.580 (2)	153
C14—H14 $\cdots$ N12 <sup>iii</sup>	0.93	2.75	3.387 (2)	127

Symmetry codes: (i)  $-x, -y + 1, -z + 2$ ; (ii)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (iii)  $-x + 1, -y + 1, -z + 1$ .

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2002); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL97*.

The authors thank the National Research Foundation of South Africa for financial support. LD also thanks the Research Foundation Flanders (FWO).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KP2340).

### References

- Bruce, M. I., Nicholson, B. K. & Shawkataly, O. B. (1989). *Inorg. Synth.* **26**, 324–326.
- Bruker (2001). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2002). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Desiraju, G. R. & Steiner, T. (2006). In *The Weak Hydrogen Bond in Structural Chemistry and Biology*. Oxford University Press.
- Hagos, T. K., Nogai, S. D., Dobrzańska, L., Cronje, S. & Raubenheimer, H. G. (2010). *Acta Cryst. E66*, o2378.
- Jian, F.-F., Yi, W., Wang, L.-M. & Wang, J. (2007). *Acta Cryst. E63*, o3887.
- Lokaj, J., Kettmann, V., Stetinová, J., Kada, R. & Vrábel, V. (1997). *Acta Cryst. C53*, 1358–1359.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Muir, M. M., Cox, O., Bernard, L. & Muir, J. A. (1992). *Acta Cryst. C48*, 583–585.
- Sheldrick, G. M. (1997). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

# supporting information

*Acta Cryst.* (2011). E67, o2014 [doi:10.1107/S1600536811026924]

## 1,1,2,2-Tetrakis(1,3-benzoxazol-2-yl)ethene

**Tesfamariam K. Hagos, Stefan D. Nogai, Liliana Dobrzańska, Stephanie Cronje and Helgard G. Raubenheimer**

### S1. Comment


The title compound is reminiscent of the crystal structure of 1,1,2,2-tetrakis(1,3-benzothiazol-2-yl)ethene chloroform solvate (Hagos *et al.*, 2010) and was isolated in a similar way. The asymmetric unit consists of half of the molecule with the other part being generated by an inversion centre (Fig. 1). The conformation of the molecule resembles that of the previous compound with a dihedral angle between the planes of the benzoxazole rings attached to the same carbon of 74.39 (5) $^{\circ}$ , whereas the corresponding value in the benzothiazole derivative was 85.74 (4) $^{\circ}$ . The bond length of 1.356 (3) Å for C1—C1<sup>i</sup> (symmetry operation (i):  $-x, 1 - y, 1 - z$ ) is almost identical to that in the earlier described structure (1.359 (3) Å). The bond lengths for the benzoxazole rings are comparable with previously reported values (Jian *et al.*, 2007; Lokaj *et al.*, 1997; Muir *et al.*, 1992). The packing is dominated by weak C—H···N (Table 1) and  $\pi$ – $\pi$  interactions. The hydrogen bonds between C7—H7···N12 and C17—H17···N3 generate corrugated layers which are stabilised by  $\pi$ – $\pi$  stacking interactions of the oxazole and phenyl rings with a distance of 3.652 (1) Å between the centroids of these rings (symmetry operation:  $-x, 1 - y, 2 - z$ ) and *ca.* 1.41 Å slippage. In addition to this, N12 acts as a bifurcated hydrogen bond acceptor and interlinks the layers through C14—H14···N12 weak interaction to form a three-dimensional assembly (Fig. 2).

### S2. Experimental


A solution of bis(2-benzoxazolyl)methane (0.075 g, 0.30 mmol) in THF (20 mL) at 253 K was treated with *n*-BuLi in *n*-hexane (0.25 mL, 1.4 M, 0.35 mmol) and stirred for 1 h. The temperature was then slowly raised to room temperature. A suspension of an excess of S<sub>8</sub> (*ca.* 2 mol equivalents) in 20 mL of THF was added to the mixture at 253 K and stirred for 1 h. After treating the resulting mixture with a solution of AuCl(PPh<sub>3</sub>) (0.15 g, 0.33 mmol) (Bruce *et al.*, 1989) in THF (20 mL), the mixture was allowed to slowly warm to ambient temperature while stirring. The solvent was removed under reduced pressure. Crystallisation of a dichloromethane solution of the residue layered with *n*-heptane at 253 K afforded light yellow crystals. Single crystal X-ray studies revealed that the unexpected oxidative dimerisation of bis(2-benzoxazolyl)methane had occurred to yield the title compound.

### S3. Refinement

All H atoms were positioned geometrically, with C—H = 0.93 and constrained to ride on their parent atoms with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

**Figure 1**

The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Unlabeled atoms are related to the labeled ones by the symmetry operation (i)  $-x, 1 - y, 1 - z$ .

**Figure 2**

Representation of the packing viewed down the  $c$  axis. The corrugated layers formed by weak C—H···N and  $\pi$ – $\pi$  interactions are indicated in blue and red; C14—H14···N12 contacts between these layers are shown as dotted light-blue lines.

### 1,1,2,2-Tetrakis(1,3-benzoxazol-2-yl)ethene

#### Crystal data

$C_{30}H_{16}N_4O_4$

$M_r = 496.47$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 9.2697 (9)$  Å

$b = 16.1943 (16)$  Å

$c = 8.0332 (8)$  Å

$\beta = 104.395 (2)^\circ$

$V = 1168.1 (2)$  Å<sup>3</sup>

$Z = 2$

$F(000) = 512$

$D_x = 1.412$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 3523 reflections

$\theta = 2.5\text{--}28.3^\circ$

$\mu = 0.10$  mm<sup>-1</sup>

$T = 273\text{ K}$   
Block, yellow

$0.30 \times 0.25 \times 0.15\text{ mm}$

#### Data collection

Bruker APEX CCD area-detector  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  scans  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1997)  
 $T_{\min} = 0.972$ ,  $T_{\max} = 0.986$

6999 measured reflections  
2717 independent reflections  
2458 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$   
 $\theta_{\max} = 28.3^\circ$ ,  $\theta_{\min} = 2.3^\circ$   
 $h = -12 \rightarrow 12$   
 $k = -17 \rightarrow 21$   
 $l = -10 \rightarrow 7$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.121$   
 $S = 1.06$   
2717 reflections  
172 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0513P)^2 + 0.8168P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.37\text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.30\text{ e } \text{\AA}^{-3}$

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.02783 (16)	0.46822 (9)	0.55330 (19)	0.0186 (3)
C2	-0.02821 (16)	0.44239 (9)	0.69934 (19)	0.0199 (3)
N3	0.01487 (14)	0.37588 (8)	0.78861 (16)	0.0197 (3)
C4	-0.06473 (16)	0.37680 (9)	0.91493 (19)	0.0196 (3)
C5	-0.06558 (18)	0.32189 (10)	1.0482 (2)	0.0240 (3)
H5	-0.0059	0.2749	1.0664	0.029*
C6	-0.15995 (18)	0.34088 (10)	1.1524 (2)	0.0258 (3)
H6	-0.1640	0.3055	1.2423	0.031*
C7	-0.24923 (18)	0.41171 (10)	1.1263 (2)	0.0262 (4)
H7	-0.3114	0.4218	1.1988	0.031*
C8	-0.24768 (18)	0.46738 (10)	0.9955 (2)	0.0246 (3)
H8	-0.3060	0.5148	0.9782	0.029*
C9	-0.15338 (16)	0.44734 (9)	0.89261 (19)	0.0199 (3)
O10	-0.12962 (12)	0.48989 (7)	0.75253 (14)	0.0217 (3)

C11	0.15956 (17)	0.42084 (10)	0.53413 (19)	0.0217 (3)
N12	0.29464 (13)	0.44984 (7)	0.58136 (16)	0.0179 (3)
C13	0.38086 (17)	0.38841 (9)	0.53122 (19)	0.0194 (3)
C14	0.53275 (17)	0.38556 (10)	0.5437 (2)	0.0251 (3)
H14	0.5959	0.4286	0.5920	0.030*
C15	0.58507 (18)	0.31551 (11)	0.4807 (2)	0.0284 (4)
H15	0.6863	0.3111	0.4872	0.034*
C16	0.4904 (2)	0.25091 (11)	0.4074 (2)	0.0309 (4)
H16	0.5307	0.2048	0.3666	0.037*
C17	0.33806 (19)	0.25342 (10)	0.3935 (2)	0.0284 (4)
H17	0.2746	0.2106	0.3450	0.034*
C18	0.28725 (16)	0.32445 (9)	0.45730 (19)	0.0205 (3)
O19	0.14322 (12)	0.34621 (7)	0.46062 (15)	0.0258 (3)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0167 (7)	0.0186 (7)	0.0206 (7)	-0.0016 (5)	0.0050 (5)	-0.0009 (5)
C2	0.0164 (7)	0.0216 (7)	0.0211 (7)	-0.0023 (5)	0.0037 (6)	-0.0013 (6)
N3	0.0192 (6)	0.0191 (6)	0.0211 (6)	-0.0002 (5)	0.0057 (5)	0.0020 (5)
C4	0.0190 (7)	0.0195 (7)	0.0197 (7)	-0.0035 (5)	0.0037 (6)	-0.0023 (5)
C5	0.0274 (8)	0.0202 (7)	0.0233 (8)	-0.0018 (6)	0.0044 (6)	0.0016 (6)
C6	0.0314 (8)	0.0260 (8)	0.0199 (7)	-0.0096 (6)	0.0062 (6)	-0.0002 (6)
C7	0.0272 (8)	0.0306 (8)	0.0236 (8)	-0.0072 (6)	0.0114 (6)	-0.0070 (6)
C8	0.0236 (8)	0.0249 (8)	0.0266 (8)	-0.0004 (6)	0.0088 (6)	-0.0039 (6)
C9	0.0202 (7)	0.0197 (7)	0.0189 (7)	-0.0033 (5)	0.0035 (5)	-0.0007 (6)
O10	0.0217 (5)	0.0216 (5)	0.0234 (5)	0.0014 (4)	0.0084 (4)	0.0019 (4)
C11	0.0239 (8)	0.0227 (7)	0.0197 (7)	0.0061 (6)	0.0080 (6)	0.0067 (6)
N12	0.0155 (6)	0.0149 (6)	0.0246 (6)	0.0013 (4)	0.0075 (5)	0.0011 (5)
C13	0.0210 (7)	0.0180 (7)	0.0202 (7)	0.0026 (5)	0.0069 (6)	0.0017 (5)
C14	0.0202 (7)	0.0283 (8)	0.0270 (8)	-0.0029 (6)	0.0062 (6)	0.0031 (6)
C15	0.0203 (8)	0.0376 (9)	0.0291 (8)	0.0079 (7)	0.0097 (6)	0.0068 (7)
C16	0.0343 (9)	0.0297 (9)	0.0308 (9)	0.0113 (7)	0.0123 (7)	-0.0028 (7)
C17	0.0311 (9)	0.0246 (8)	0.0291 (8)	-0.0005 (6)	0.0068 (7)	-0.0056 (6)
C18	0.0192 (7)	0.0223 (7)	0.0200 (7)	0.0019 (6)	0.0047 (6)	0.0012 (6)
O19	0.0220 (6)	0.0265 (6)	0.0283 (6)	0.0015 (4)	0.0051 (5)	-0.0013 (5)

*Geometric parameters ( $\text{\AA}$ ,  $\text{^\circ}$ )*

C1—C1 <sup>i</sup>	1.356 (3)	C9—O10	1.3830 (18)
C1—C2	1.458 (2)	C11—N12	1.302 (2)
C1—C11	1.482 (2)	C11—O19	1.3370 (19)
C2—N3	1.3006 (19)	N12—C13	1.3966 (18)
C2—O10	1.3628 (18)	C13—C18	1.386 (2)
N3—C4	1.3952 (19)	C13—C14	1.387 (2)
C4—C9	1.393 (2)	C14—C15	1.379 (2)
C4—C5	1.394 (2)	C14—H14	0.9300
C5—C6	1.387 (2)	C15—C16	1.398 (3)

C5—H5	0.9300	C15—H15	0.9300
C6—C7	1.399 (2)	C16—C17	1.389 (2)
C6—H6	0.9300	C16—H16	0.9300
C7—C8	1.387 (2)	C17—C18	1.388 (2)
C7—H7	0.9300	C17—H17	0.9300
C8—C9	1.382 (2)	C18—O19	1.3874 (18)
C8—H8	0.9300		
C1 <sup>i</sup> —C1—C2	124.77 (17)	C2—O10—C9	103.63 (11)
C1 <sup>i</sup> —C1—C11	121.42 (17)	N12—C11—O19	116.73 (13)
C2—C1—C11	113.75 (12)	N12—C11—C1	122.70 (14)
N3—C2—O10	115.83 (13)	O19—C11—C1	120.53 (13)
N3—C2—C1	124.01 (14)	C11—N12—C13	103.70 (12)
O10—C2—C1	120.14 (13)	C18—C13—C14	121.47 (14)
C2—N3—C4	104.29 (12)	C18—C13—N12	108.27 (13)
C9—C4—C5	120.43 (14)	C14—C13—N12	130.25 (14)
C9—C4—N3	108.48 (13)	C15—C14—C13	116.37 (15)
C5—C4—N3	131.09 (14)	C15—C14—H14	121.8
C6—C5—C4	116.56 (15)	C13—C14—H14	121.8
C6—C5—H5	121.7	C14—C15—C16	121.96 (15)
C4—C5—H5	121.7	C14—C15—H15	119.0
C5—C6—C7	121.96 (15)	C16—C15—H15	119.0
C5—C6—H6	119.0	C17—C16—C15	122.06 (15)
C7—C6—H6	119.0	C17—C16—H16	119.0
C8—C7—C6	121.95 (15)	C15—C16—H16	119.0
C8—C7—H7	119.0	C18—C17—C16	115.22 (15)
C6—C7—H7	119.0	C18—C17—H17	122.4
C9—C8—C7	115.30 (15)	C16—C17—H17	122.4
C9—C8—H8	122.4	C13—C18—O19	107.75 (13)
C7—C8—H8	122.4	C13—C18—C17	122.92 (14)
C8—C9—O10	128.44 (14)	O19—C18—C17	129.34 (14)
C8—C9—C4	123.79 (14)	C11—O19—C18	103.54 (12)
O10—C9—C4	107.77 (13)		
C1 <sup>i</sup> —C1—C2—N3	171.93 (18)	C1 <sup>i</sup> —C1—C11—N12	75.2 (2)
C11—C1—C2—N3	-11.0 (2)	C2—C1—C11—N12	-102.04 (16)
C1 <sup>i</sup> —C1—C2—O10	-9.6 (3)	C1 <sup>i</sup> —C1—C11—O19	-102.5 (2)
C11—C1—C2—O10	167.47 (13)	C2—C1—C11—O19	80.33 (17)
O10—C2—N3—C4	0.47 (17)	O19—C11—N12—C13	1.25 (17)
C1—C2—N3—C4	178.99 (13)	C1—C11—N12—C13	-176.47 (13)
C2—N3—C4—C9	-0.45 (15)	C11—N12—C13—C18	-0.89 (16)
C2—N3—C4—C5	-179.84 (16)	C11—N12—C13—C14	178.08 (16)
C9—C4—C5—C6	1.1 (2)	C18—C13—C14—C15	-0.7 (2)
N3—C4—C5—C6	-179.58 (15)	N12—C13—C14—C15	-179.59 (15)
C4—C5—C6—C7	-0.4 (2)	C13—C14—C15—C16	0.3 (2)
C5—C6—C7—C8	-0.5 (2)	C14—C15—C16—C17	0.0 (3)
C6—C7—C8—C9	0.7 (2)	C15—C16—C17—C18	0.1 (3)
C7—C8—C9—O10	179.17 (14)	C14—C13—C18—O19	-178.77 (13)

C7—C8—C9—C4	0.0 (2)	N12—C13—C18—O19	0.32 (16)
C5—C4—C9—C8	-1.0 (2)	C14—C13—C18—C17	0.9 (2)
N3—C4—C9—C8	179.59 (14)	N12—C13—C18—C17	180.00 (14)
C5—C4—C9—O10	179.77 (13)	C16—C17—C18—C13	-0.6 (2)
N3—C4—C9—O10	0.30 (16)	C16—C17—C18—O19	179.05 (15)
N3—C2—O10—C9	-0.29 (16)	N12—C11—O19—C18	-1.06 (17)
C1—C2—O10—C9	-178.87 (13)	C1—C11—O19—C18	176.71 (13)
C8—C9—O10—C2	-179.27 (15)	C13—C18—O19—C11	0.38 (15)
C4—C9—O10—C2	-0.03 (15)	C17—C18—O19—C11	-179.28 (16)

Symmetry code: (i)  $-x, -y+1, -z+1$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C7—H7 <sup>ii</sup> ···N12 <sup>ii</sup>	0.93	2.71	3.348 (2)	127
C17—H17 <sup>iii</sup> ···N3 <sup>iii</sup>	0.93	2.71	3.580 (2)	153
C14—H14 <sup>iv</sup> ···N12 <sup>iv</sup>	0.93	2.75	3.387 (2)	127

Symmetry codes: (ii)  $-x, -y+1, -z+2$ ; (iii)  $x, -y+1/2, z-1/2$ ; (iv)  $-x+1, -y+1, -z+1$ .