

3-Amino-N'-(2-oxoindolin-3-ylidene)-benzohydrazide

Rifat Ara Jamal,^{a*} Uzma Ashiq,^a Sammer Yousuf^b and Qurrat ul Ain^a

^aDepartment of Chemistry, University of Karachi, Karachi 75270, Pakistan, and ^bH.E.J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan

Correspondence e-mail: rifat_jamal@yahoo.com


Received 15 July 2011; accepted 19 July 2011

Key indicators: single-crystal X-ray study; $T = 273\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.052; wR factor = 0.122; data-to-parameter ratio = 12.6.

The title compound, $\text{C}_{15}\text{H}_{12}\text{N}_4\text{O}_2$, contains two substituted benzohydrazide and indole rings linked via a $\text{C}=\text{N}$ double bond. The dihedral angle between the benzene ring and the indole ring system is $11.38(10)^\circ$. The molecular structure is stabilized by an intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond, forming a six-membered ring. The crystal structure is consolidated by intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ interactions, which result in sheets.

Related literature

For the biological activity of related compounds, see: Ashiq *et al.* (2008); Maqsood *et al.* (2006); Sarangapani & Reddy (1994). For related structures, see: Bai *et al.* (2006); Yang & Pan (2004).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{12}\text{N}_4\text{O}_2$

$M_r = 280.29$

Monoclinic, $P2_1/c$

$a = 8.8036(8)\text{ \AA}$

$b = 8.9040(7)\text{ \AA}$

$c = 17.0732(14)\text{ \AA}$

$\beta = 92.335(2)^\circ$

$V = 1337.21(19)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.10\text{ mm}^{-1}$

$T = 273\text{ K}$

$0.17 \times 0.17 \times 0.12\text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.984$, $T_{\max} = 0.989$

7637 measured reflections
2491 independent reflections
1551 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.122$
 $S = 0.99$
2491 reflections
198 parameters
1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.18\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.17\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1A \cdots O2 ⁱ	0.86	2.02	2.785 (2)	148
N3—H3A \cdots O1	0.86	2.09	2.757 (2)	133
N4—H4C \cdots N2 ⁱⁱ	0.86 (2)	2.61 (2)	3.423 (3)	158 (2)
C12—H12A \cdots O1 ⁱⁱⁱ	0.93	2.59	3.424 (3)	149
Symmetry codes: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 2, -y + 1, -z$.				

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*, *PARST* (Nardelli, 1995) and *PLATON* (Spek, 2009).

The authors are thankful to the Higher Education Commission (HEC) Pakistan for financial support under the National Research Grants Program for Universities (grant No. 20-1862/R&D/10).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2431).

References

- Ashiq, U., Ara, R., Mahroof-Tahir, M., Maqsood, Z. T., Khan, K. M., Khan, S. N., Siddiqui, H. & Choudhary, M. I. (2008). *Chem. Biodivers.* **5**, 82–92.
- Bai, Z.-C., Li, H., Liu, Y. & Jing, Z.-L. (2006). *Acta Cryst. E62*, o2295–o2296.
- Bruker (2000). *SADABS*, *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Maqsood, Z. T., Khan, K. M., Ashiq, U., Ara, R., Chohan, Z. H., Mahroof-Tahir, M. & Supuran, C. T. (2006). *J. Enz. Inhib. Med. Chem.* **21**, 37–42.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Sarangapani, M. & Reddy, V. M. (1994). *Indian J. Pharm. Sci.* **56**, 174–177.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Yang, J.-G. & Pan, F.-Y. (2004). *Acta Cryst. E60*, o2009–o2010.

supporting information

Acta Cryst. (2011). E67, o2166 [doi:10.1107/S1600536811029242]

3-Amino-*N'*-(2-oxoindolin-3-ylidene)benzohydrazide

Rifat Ara Jamal, Uzma Ashiq, Sammer Yousuf and Qurrat ul Ain

S1. Comment

Compounds containing hydrazide moiety have been revealed to exhibit a wide variety of interesting biological properties (Ashiq *et al.* 2008; Maqsood *et al.* 2006). In a quest to expand and further explore the biological significance of hydrazides, we have undertaken a task to synthesize new Schiff bases of isatin with hydrazides. Isatins are very important compounds due to their antifungal properties (Sarangapani & Reddy, 1994). In this article we report the synthesis and crystal structure of the title compound.


The title molecule (Fig. 1) consists of substituted benzohydrazide and indole rings linked by C=N bond. The dihedral angle between the two substituted benzene (C10—C15) and indole ring (N1/C1—C8) is 11.38 (10) $^{\circ}$. The bond lengths and angles are in normal range as in other structurally related compounds (Bai *et al.* 2006; Yang & Pan, 2004). The geometry of the molecule is stabilized by N3—H3A…O1 intramolecular hydrogen bond resulting in a six membered ring. In the crystal structure, the molecules are linked to form two-dimensional sheets *via* N1—H1A…O2, N4—H4C…N2 and C12—H12A…O1 intermolecular hydrogen bonds (Tab. 1 & Fig. 2).

S2. Experimental


To a solution of 2,3-indolinedione (10 mmol, 1.47 g) in 15 ml of ethanol with a few drops of glacial acetic acid and 3-aminobenzohydrazide (10 mmol, 1.51 g) in 15 ml ethanol were added. The mixture was refluxed for 2 h and a solid was obtained upon removal of the solvent by rotary evaporation. The resulting solid was washed with ethanol to afford the title compound (yield 75%). The crystals of the title compound suitable for XRD analysis were grown from a mixture of ethanol and methanol (1:1) by slow evaporation at room temperature.

S3. Refinement

H atoms on the C atoms and N1 and N3 were positioned geometrically with C—H = 0.93 and N—H = 0.86 Å and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}/\text{N})$. The H atoms on N4 were located from a difference Fourier map and refined isotropically.

**Figure 1**

The molecular structure of the title molecule with displacement ellipsoids drawn at 50% probability level. The dashed lines indicate intramolecular hydrogen bond.

**Figure 2**

The crystal packing of the title compound. Hydrogen atoms not involved in hydrogen bonding have been excluded for clarity.

3-Amino-N'-(2-oxoindolin-3-ylidene)benzohydrazide*Crystal data*

$C_{15}H_{12}N_4O_2$
 $M_r = 280.29$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 8.8036 (8)$ Å
 $b = 8.9040 (7)$ Å
 $c = 17.0732 (14)$ Å
 $\beta = 92.335 (2)$ °
 $V = 1337.21 (19)$ Å³
 $Z = 4$

$F(000) = 584$
 $D_x = 1.392$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 1161 reflections
 $\theta = 2.4\text{--}27.8$ °
 $\mu = 0.10$ mm⁻¹
 $T = 273$ K
Block, colorless
 $0.17 \times 0.17 \times 0.12$ mm

Data collection

Bruker SMART APEX CCD area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2000)
 $T_{\min} = 0.984$, $T_{\max} = 0.989$

7637 measured reflections
2491 independent reflections
1551 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$
 $\theta_{\max} = 25.5$ °, $\theta_{\min} = 2.3$ °
 $h = -10 \rightarrow 10$
 $k = -10 \rightarrow 10$
 $l = -20 \rightarrow 20$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.122$
 $S = 0.99$
2491 reflections
198 parameters
1 restraint
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0584P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.18$ e Å⁻³
 $\Delta\rho_{\min} = -0.17$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.66942 (18)	0.35101 (18)	-0.07613 (8)	0.0531 (5)
O2	0.53184 (19)	0.37905 (18)	0.20442 (8)	0.0568 (5)
N1	0.5172 (2)	0.1671 (2)	-0.13475 (10)	0.0470 (5)

H1A	0.5578	0.1631	-0.1797	0.056*
N2	0.4730 (2)	0.27880 (19)	0.06115 (9)	0.0407 (5)
N3	0.5848 (2)	0.38136 (19)	0.07676 (9)	0.0415 (5)
H3A	0.6377	0.4169	0.0398	0.050*
N4	0.8354 (3)	0.8078 (3)	0.32398 (15)	0.0676 (7)
C1	0.2412 (3)	0.0351 (3)	-0.00233 (13)	0.0488 (6)
H1B	0.2175	0.0545	0.0493	0.059*
C2	0.1601 (3)	-0.0703 (3)	-0.04635 (15)	0.0557 (7)
H2A	0.0815	-0.1229	-0.0241	0.067*
C3	0.1950 (3)	-0.0980 (3)	-0.12350 (14)	0.0569 (7)
H3B	0.1379	-0.1682	-0.1524	0.068*
C4	0.3119 (3)	-0.0245 (3)	-0.15845 (13)	0.0523 (7)
H4A	0.3354	-0.0442	-0.2101	0.063*
C5	0.3923 (2)	0.0790 (2)	-0.11405 (12)	0.0417 (6)
C6	0.5659 (3)	0.2585 (3)	-0.07583 (12)	0.0409 (6)
C7	0.4654 (2)	0.2238 (2)	-0.00862 (11)	0.0374 (5)
C8	0.3583 (2)	0.1110 (2)	-0.03664 (12)	0.0391 (5)
C9	0.6110 (2)	0.4264 (2)	0.15255 (12)	0.0406 (6)
C10	0.7362 (2)	0.5353 (2)	0.16743 (12)	0.0400 (6)
C11	0.8540 (3)	0.5576 (3)	0.11661 (13)	0.0518 (6)
H11A	0.8582	0.5027	0.0704	0.062*
C12	0.9645 (3)	0.6629 (3)	0.13619 (15)	0.0606 (7)
H12A	1.0442	0.6778	0.1030	0.073*
C13	0.9587 (3)	0.7457 (3)	0.20365 (16)	0.0586 (7)
H13A	1.0342	0.8161	0.2154	0.070*
C14	0.8416 (3)	0.7259 (3)	0.25467 (14)	0.0490 (6)
C15	0.7329 (3)	0.6181 (2)	0.23602 (13)	0.0440 (6)
H15A	0.6555	0.6009	0.2704	0.053*
H4C	0.7478 (19)	0.819 (3)	0.3435 (17)	0.099 (12)*
H4B	0.889 (4)	0.894 (3)	0.3239 (17)	0.101 (11)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0485 (10)	0.0635 (11)	0.0480 (9)	-0.0019 (9)	0.0114 (8)	0.0040 (8)
O2	0.0650 (11)	0.0715 (12)	0.0346 (8)	-0.0188 (9)	0.0111 (8)	-0.0022 (8)
N1	0.0485 (12)	0.0630 (12)	0.0299 (9)	0.0067 (11)	0.0055 (9)	0.0003 (9)
N2	0.0402 (11)	0.0452 (11)	0.0366 (10)	0.0030 (9)	0.0024 (9)	0.0015 (8)
N3	0.0439 (11)	0.0476 (11)	0.0334 (10)	-0.0005 (10)	0.0080 (9)	0.0016 (8)
N4	0.0598 (18)	0.0655 (16)	0.0765 (16)	-0.0067 (14)	-0.0089 (14)	-0.0142 (14)
C1	0.0431 (14)	0.0557 (15)	0.0473 (13)	0.0064 (13)	0.0012 (12)	0.0011 (12)
C2	0.0413 (15)	0.0549 (16)	0.0704 (17)	0.0005 (13)	-0.0028 (13)	0.0059 (14)
C3	0.0515 (16)	0.0529 (15)	0.0644 (17)	0.0076 (13)	-0.0192 (14)	-0.0053 (13)
C4	0.0566 (17)	0.0566 (15)	0.0429 (13)	0.0126 (14)	-0.0088 (13)	-0.0042 (12)
C5	0.0392 (13)	0.0476 (13)	0.0377 (12)	0.0125 (12)	-0.0042 (11)	0.0000 (11)
C6	0.0366 (14)	0.0488 (14)	0.0372 (12)	0.0113 (12)	0.0017 (11)	0.0042 (11)
C7	0.0364 (13)	0.0432 (13)	0.0326 (11)	0.0115 (11)	0.0013 (10)	0.0028 (10)
C8	0.0354 (12)	0.0437 (13)	0.0379 (11)	0.0089 (11)	-0.0004 (10)	0.0025 (10)

C9	0.0436 (14)	0.0440 (13)	0.0345 (12)	0.0075 (11)	0.0040 (11)	0.0023 (10)
C10	0.0361 (13)	0.0412 (13)	0.0426 (12)	0.0055 (11)	0.0021 (10)	0.0070 (11)
C11	0.0452 (15)	0.0603 (16)	0.0504 (13)	0.0042 (13)	0.0079 (12)	0.0053 (12)
C12	0.0403 (15)	0.0737 (18)	0.0685 (17)	-0.0012 (14)	0.0084 (14)	0.0180 (15)
C13	0.0447 (16)	0.0558 (16)	0.0743 (17)	-0.0074 (13)	-0.0084 (14)	0.0100 (14)
C14	0.0428 (15)	0.0479 (14)	0.0557 (14)	0.0047 (12)	-0.0055 (13)	0.0030 (12)
C15	0.0377 (13)	0.0472 (14)	0.0470 (13)	0.0014 (11)	-0.0012 (11)	0.0047 (11)

Geometric parameters (\AA , $^{\circ}$)

O1—C6	1.228 (2)	C3—H3B	0.9300
O2—C9	1.224 (2)	C4—C5	1.372 (3)
N1—C6	1.351 (3)	C4—H4A	0.9300
N1—C5	1.407 (3)	C5—C8	1.396 (3)
N1—H1A	0.8600	C6—C7	1.509 (3)
N2—C7	1.287 (2)	C7—C8	1.446 (3)
N2—N3	1.361 (2)	C9—C10	1.482 (3)
N3—C9	1.365 (2)	C10—C15	1.385 (3)
N3—H3A	0.8600	C10—C11	1.393 (3)
N4—C14	1.393 (3)	C11—C12	1.382 (3)
N4—H4C	0.858 (10)	C11—H11A	0.9300
N4—H4B	0.90 (3)	C12—C13	1.370 (3)
C1—C2	1.382 (3)	C12—H12A	0.9300
C1—C8	1.383 (3)	C13—C14	1.388 (3)
C1—H1B	0.9300	C13—H13A	0.9300
C2—C3	1.387 (3)	C14—C15	1.383 (3)
C2—H2A	0.9300	C15—H15A	0.9300
C3—C4	1.376 (3)		
C6—N1—C5	112.13 (18)	N2—C7—C8	125.39 (19)
C6—N1—H1A	123.9	N2—C7—C6	128.0 (2)
C5—N1—H1A	123.9	C8—C7—C6	106.59 (17)
C7—N2—N3	116.52 (17)	C1—C8—C5	119.6 (2)
N2—N3—C9	118.39 (17)	C1—C8—C7	133.39 (19)
N2—N3—H3A	120.8	C5—C8—C7	107.01 (18)
C9—N3—H3A	120.8	O2—C9—N3	120.4 (2)
C14—N4—H4C	117 (2)	O2—C9—C10	122.86 (19)
C14—N4—H4B	113.9 (19)	N3—C9—C10	116.76 (18)
H4C—N4—H4B	113 (3)	C15—C10—C11	119.5 (2)
C2—C1—C8	118.6 (2)	C15—C10—C9	116.88 (19)
C2—C1—H1B	120.7	C11—C10—C9	123.6 (2)
C8—C1—H1B	120.7	C12—C11—C10	118.7 (2)
C1—C2—C3	120.5 (2)	C12—C11—H11A	120.6
C1—C2—H2A	119.7	C10—C11—H11A	120.6
C3—C2—H2A	119.7	C13—C12—C11	121.2 (2)
C4—C3—C2	121.7 (2)	C13—C12—H12A	119.4
C4—C3—H3B	119.2	C11—C12—H12A	119.4
C2—C3—H3B	119.2	C12—C13—C14	120.9 (2)

C5—C4—C3	117.3 (2)	C12—C13—H13A	119.6
C5—C4—H4A	121.4	C14—C13—H13A	119.6
C3—C4—H4A	121.4	C15—C14—C13	117.9 (2)
C4—C5—C8	122.3 (2)	C15—C14—N4	120.5 (2)
C4—C5—N1	128.8 (2)	C13—C14—N4	121.5 (2)
C8—C5—N1	108.91 (19)	C14—C15—C10	121.7 (2)
O1—C6—N1	127.9 (2)	C14—C15—H15A	119.1
O1—C6—C7	126.8 (2)	C10—C15—H15A	119.1
N1—C6—C7	105.3 (2)		
C7—N2—N3—C9	-170.74 (18)	N1—C5—C8—C7	0.4 (2)
C8—C1—C2—C3	-0.5 (3)	N2—C7—C8—C1	0.8 (4)
C1—C2—C3—C4	1.1 (4)	C6—C7—C8—C1	179.3 (2)
C2—C3—C4—C5	-0.6 (3)	N2—C7—C8—C5	-177.82 (19)
C3—C4—C5—C8	-0.4 (3)	C6—C7—C8—C5	0.6 (2)
C3—C4—C5—N1	178.9 (2)	N2—N3—C9—O2	-2.5 (3)
C6—N1—C5—C4	179.2 (2)	N2—N3—C9—C10	178.53 (17)
C6—N1—C5—C8	-1.4 (2)	O2—C9—C10—C15	-19.6 (3)
C5—N1—C6—O1	-178.1 (2)	N3—C9—C10—C15	159.37 (19)
C5—N1—C6—C7	1.7 (2)	O2—C9—C10—C11	160.6 (2)
N3—N2—C7—C8	178.30 (18)	N3—C9—C10—C11	-20.5 (3)
N3—N2—C7—C6	0.2 (3)	C15—C10—C11—C12	-0.2 (3)
O1—C6—C7—N2	-3.2 (4)	C9—C10—C11—C12	179.6 (2)
N1—C6—C7—N2	177.0 (2)	C10—C11—C12—C13	-0.7 (4)
O1—C6—C7—C8	178.4 (2)	C11—C12—C13—C14	0.2 (4)
N1—C6—C7—C8	-1.4 (2)	C12—C13—C14—C15	1.2 (4)
C2—C1—C8—C5	-0.5 (3)	C12—C13—C14—N4	179.1 (2)
C2—C1—C8—C7	-179.0 (2)	C13—C14—C15—C10	-2.2 (3)
C4—C5—C8—C1	1.0 (3)	N4—C14—C15—C10	179.9 (2)
N1—C5—C8—C1	-178.47 (19)	C11—C10—C15—C14	1.7 (3)
C4—C5—C8—C7	179.85 (19)	C9—C10—C15—C14	-178.15 (19)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1A···O2 ⁱ	0.86	2.02	2.785 (2)	148
N3—H3A···O1	0.86	2.09	2.757 (2)	133
N4—H4C···N2 ⁱⁱ	0.86 (2)	2.61 (2)	3.423 (3)	158 (2)
C12—H12A···O1 ⁱⁱⁱ	0.93	2.59	3.424 (3)	149

Symmetry codes: (i) $x, -y+1/2, z-1/2$; (ii) $-x+1, y+1/2, -z+1/2$; (iii) $-x+2, -y+1, -z$.