

9-[3-(Dimethylamino)propyl]-2-trifluoro-methyl-9*H*-thioxanthen-9-ol

Jerry P. Jasinski,^{a*} James A. Golen,^a M. S. Siddegowda,^b H. S. Yathirajan^b and M. T. Swamy^c

^aDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, ^bDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, and ^cDepartment of Chemistry, Sambhram Institute of Technology, Bangalore 560 097, India
Correspondence e-mail: jjjasinski@keene.edu


Received 23 June 2011; accepted 27 June 2011

Key indicators: single-crystal X-ray study; $T = 170\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; disorder in main residue; R factor = 0.043; wR factor = 0.116; data-to-parameter ratio = 15.3.

In the title compound, $\text{C}_{19}\text{H}_{20}\text{F}_3\text{NOS}$, the dihedral angle between the mean planes of the two benzene rings attached to the thioxanthene ring is $41.8(7)^\circ$; the latter has a slightly distorted boat conformation. The F atoms are disordered over three sets of sites [occupancy ratio = 0.564 (10):0.287 (10):0.148 (5)] and the methyl groups are disordered over two sets of sites [occupancy ratio = 0.72 (4):0.28 (4)]. The crystal packing is stabilized by $\text{O}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{S}$ hydrogen bonds and weak $\text{C}-\text{H}\cdots\text{Cg}$ interactions.

Related literature

For photo-initiators with excellent capabilities in UV-curing materials, see: Fouassier *et al.* (1995); Roffey (1997). For related structures, see: Post *et al.* (1975a,b); Liu, (2009). For puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{19}\text{H}_{20}\text{F}_3\text{NOS}$

$M_r = 367.42$

Monoclinic, $P2_1/n$

$a = 7.6183(3)\text{ \AA}$

$b = 13.9605(4)\text{ \AA}$

$c = 17.4172(7)\text{ \AA}$

$\beta = 101.053(4)^\circ$
 $V = 1818.05(11)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.21\text{ mm}^{-1}$
 $T = 170\text{ K}$
 $0.35 \times 0.33 \times 0.30\text{ mm}$

Data collection

Oxford Diffraction Xcalibur Eos Gemini diffractometer
Absorption correction: multi-scan (*CrysAlis RED*; Oxford Diffraction, 2010)
 $T_{\min} = 0.929$, $T_{\max} = 0.939$

17330 measured reflections
4697 independent reflections
3901 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.116$
 $S = 1.05$
4697 reflections
306 parameters
238 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.28\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.29\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg2$ and $Cg3$ are the centroids of the C2–C7 and C8–C13 rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1–H1A…N1	0.87 (2)	1.84 (2)	2.7141 (17)	176 (2)
C15–H15A…S1	0.99	2.76	3.4165 (15)	124
C5–H5A… $Cg3^i$	0.95	2.96	3.798 (3)	148
C17–H17A… $Cg2^{ii}$	0.99	2.97	3.949 (3)	170
C17–H17B… $Cg3^{iii}$	0.99	2.83	3.659 (3)	142

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $x - 1, y, z$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2010); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

MSS thanks the University of Mysore and R. L. Fine Chem, Bangalore, for access to their research facilities. JPJ acknowledges the NSF–MRI program (grant No. CHE1039027) for funds to purchase the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2761).

References

- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Fouassier, J. P., Ruhmann, D., Graff, D., Morlet-Savary, F. & Wieder, F. (1995). *Prog. Org. Coat.* **25**, 235–271.
- Liu, G. (2009). *Acta Cryst. E65*, o613.
- Oxford Diffraction (2010). *CrysAlis PRO* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
- Post, M. L., Kennard, O. & Horn, A. S. (1975a). *Acta Cryst. B31*, 2724–2726.
- Post, M. L., Kennard, O., Sheldrick, G. M. & Horn, A. S. (1975b). *Acta Cryst. B31*, 2366–2368.
- Roffey, C. (1997). In *Photogeneration of Reactive Species for UV-Curing*. Sussex, England: Wiley.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

supporting information

Acta Cryst. (2011). E67, o1883 [doi:10.1107/S1600536811025311]

9-[3-(Dimethylamino)propyl]-2-trifluoromethyl-9H-thioxanthen-9-ol

Jerry P. Jasinski, James A. Golen, M. S. Siddegowda, H. S. Yathirajan and M. T. Swamy

S1. Comment

The title compound is a flupenthixol impurity with systematic IUPAC name: (*RS*)-9-[3-(dimethylamino)propyl]-2-(trifluoromethyl)-9H-thioxanthen-9-ol. Flupenthixol is a thioxanthene derivative that may exist in two isomeric forms, α and β . Flupenthixol contains 45–55 % α -flupenthixol. The pharmacological effects of flupenthixol, α - and β -flupenthixol have been compared with those of clopenthixol, chlorprothixene, fluphenazine, perphenazine, chlorpromazine and haloperidol. In most pharmacological screening tests α -flupenthixol was equipotent with fluphenazine. β -Flupenthixol showed very low pharmacological activity. As expected the potency of flupenthixol was about one half that of α -flupenthixol. Thioxanthone derivatives are good photoinitiators with excellent capabilities in UV-curing materials (Fouassier *et al.*, 1995; Roffey, 1997). The crystal structures of α -flupenthixol (Post *et al.*, 1975*b*), β -flupenthixol (Post *et al.*, 1975*a*) and 2,4-diethylthioxanthen-9-one (Liu, 2009) have been reported. In view of the importance of the title compound the crystal structure is herein reported.


In the title compound, (I), the dihedral angle between the mean planes of the two benzene rings in the thioxanthene ring is 41.8 (7) ° (Fig. 1). The thioxanthene ring is in a slightly distorted boat conformation (Cremer & Pople, 1975) with puckering parameters Q, θ , and φ = 0.591 (2) Å, 92.72 (19) ° and 359.8 (2) °, respectively). Crystal packing is stabilized by O1—H1A···N1, C15—H15A···S1 hydrogen bonds and weak C—H···Cg intermolecular interactions (Fig. 2, Table 1).

S2. Experimental


The title compound was obtained as a gift sample from R. L. Fine Chem. Ltd., Bangalore, India. The compound was recrystallized from dichloromethane (*M.pt.*: 389–391 K).

S3. Refinement

The fluorine atoms on C14 are disordered over three positions [occupancy ratio 0.564 (10); 0.287 (10); 0.148 (5)] and the methyl groups on N1 are disordered over two positions [occupancy ratio 0.72 (4); 0.28 (4)]. The O—H hydrogen atom was located by Fourier analysis and refined isotropically. All of the remaining H atoms were placed in their calculated positions and then refined using the riding model with atom—H lengths of 0.95 Å (CH), 0.99 Å (CH₂) or 0.98 Å (CH₃). Isotropic displacement parameters for these atoms were set to 1.18–1.20 (CH) or 1.20 (CH₂) times U_{eq} of the parent atom.

**Figure 1**

Molecular structure of (I) showing the atom labeling scheme and 50% probability displacement ellipsoids.

**Figure 2**

Packing diagram of the title compound viewed down the a axis. Dashed lines represent disordered C—F atoms.

9-[3-(Dimethylamino)propyl]-2-trifluoromethyl-9*H*-thioxanthen-9-ol*Crystal data*

$C_{19}H_{20}F_3NOS$
 $M_r = 367.42$
Monoclinic, $P2_1/n$
Hall symbol: -P 2yn
 $a = 7.6183 (3)$ Å
 $b = 13.9605 (4)$ Å
 $c = 17.4172 (7)$ Å
 $\beta = 101.053 (4)$ °
 $V = 1818.05 (11)$ Å³
 $Z = 4$

$F(000) = 768$
 $D_x = 1.342$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 9234 reflections
 $\theta = 3.3\text{--}32.3$ °
 $\mu = 0.21$ mm⁻¹
 $T = 170$ K
Block, pale yellow
 $0.35 \times 0.33 \times 0.30$ mm

Data collection

Oxford Diffraction Xcalibur Eos Gemini
diffractometer
Radiation source: Enhance (Mo) X-ray Source
Graphite monochromator
Detector resolution: 16.1500 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(CrysAlis RED; Oxford Diffraction, 2010)
 $T_{\min} = 0.929$, $T_{\max} = 0.939$

17330 measured reflections
4697 independent reflections
3901 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$
 $\theta_{\max} = 28.7$ °, $\theta_{\min} = 3.5$ °
 $h = -10 \rightarrow 10$
 $k = -18 \rightarrow 15$
 $l = -23 \rightarrow 23$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.116$
 $S = 1.05$
4697 reflections
306 parameters
238 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0506P)^2 + 0.6042P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.037$
 $\Delta\rho_{\max} = 0.28$ e Å⁻³
 $\Delta\rho_{\min} = -0.29$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.74858 (6)	0.83313 (3)	0.17618 (2)	0.04558 (12)	
F1	0.6729 (8)	0.6443 (4)	0.5165 (2)	0.0783 (14)	0.564 (10)
F2	0.9343 (6)	0.5995 (5)	0.5119 (4)	0.0815 (15)	0.564 (10)

F3	0.7169 (10)	0.5137 (4)	0.4572 (4)	0.0765 (16)	0.564 (10)
F1A	0.6033 (13)	0.6017 (10)	0.4888 (7)	0.078 (2)	0.287 (10)
F2A	0.8891 (19)	0.6238 (7)	0.5244 (5)	0.072 (3)	0.287 (10)
F3A	0.7863 (16)	0.5067 (7)	0.4533 (7)	0.059 (2)	0.287 (10)
F1B	0.5941 (16)	0.5562 (12)	0.4648 (6)	0.070 (3)	0.148 (5)
F2B	0.785 (3)	0.6465 (10)	0.5293 (6)	0.075 (3)	0.148 (5)
F3B	0.864 (2)	0.5254 (12)	0.4705 (10)	0.091 (4)	0.148 (5)
O1	0.50685 (14)	0.54112 (7)	0.18448 (6)	0.0369 (2)	
H1A	0.408 (3)	0.5332 (16)	0.2017 (13)	0.070 (6)*	
N1	0.19359 (19)	0.52532 (10)	0.23591 (10)	0.0543 (4)	
C1	0.54324 (17)	0.63994 (9)	0.17934 (8)	0.0316 (3)	
C2	0.65553 (17)	0.67633 (9)	0.25587 (8)	0.0319 (3)	
C3	0.66939 (18)	0.62297 (10)	0.32396 (8)	0.0346 (3)	
H3A	0.6162	0.5612	0.3220	0.041*	
C4	0.7599 (2)	0.65865 (11)	0.39484 (9)	0.0410 (3)	
C5	0.8404 (2)	0.74815 (12)	0.39867 (10)	0.0501 (4)	
H5A	0.8989	0.7732	0.4476	0.060*	
C6	0.8350 (2)	0.80039 (12)	0.33140 (10)	0.0473 (4)	
H6A	0.8935	0.8607	0.3334	0.057*	
C7	0.74356 (18)	0.76466 (10)	0.26013 (9)	0.0363 (3)	
C8	0.7353 (2)	0.74071 (11)	0.10656 (9)	0.0406 (3)	
C9	0.8206 (3)	0.75615 (14)	0.04376 (11)	0.0562 (4)	
H9A	0.8784	0.8155	0.0388	0.067*	
C10	0.8210 (3)	0.68539 (17)	-0.01102 (12)	0.0683 (6)	
H10A	0.8778	0.6962	-0.0543	0.082*	
C11	0.7391 (3)	0.59858 (16)	-0.00345 (11)	0.0641 (5)	
H11A	0.7412	0.5494	-0.0410	0.077*	
C12	0.6541 (2)	0.58328 (13)	0.05899 (9)	0.0478 (4)	
H12A	0.5992	0.5231	0.0642	0.057*	
C13	0.64777 (18)	0.65429 (10)	0.11405 (8)	0.0363 (3)	
C14	0.7655 (3)	0.60126 (14)	0.46712 (10)	0.0534 (4)	
C15	0.36815 (18)	0.69875 (11)	0.15859 (9)	0.0411 (3)	
H15A	0.4009	0.7658	0.1494	0.049*	
H15B	0.2994	0.6740	0.1085	0.049*	
C16	0.2439 (2)	0.69987 (11)	0.21787 (11)	0.0489 (4)	
H16A	0.3181	0.6992	0.2712	0.059*	
H16B	0.1758	0.7607	0.2119	0.059*	
C17	0.1118 (2)	0.61725 (12)	0.21095 (12)	0.0525 (4)	
H17A	0.0208	0.6320	0.2429	0.063*	
H17B	0.0494	0.6119	0.1558	0.063*	
C18	0.2274 (12)	0.5222 (12)	0.3225 (4)	0.095 (2)	0.72 (4)
H18A	0.3144	0.5718	0.3435	0.142*	0.72 (4)
H18B	0.2749	0.4591	0.3404	0.142*	0.72 (4)
H18C	0.1153	0.5336	0.3407	0.142*	0.72 (4)
C19	0.0843 (17)	0.4459 (7)	0.2028 (8)	0.100 (3)	0.72 (4)
H19A	0.0712	0.4470	0.1457	0.150*	0.72 (4)
H19B	-0.0339	0.4509	0.2169	0.150*	0.72 (4)
H19C	0.1412	0.3857	0.2232	0.150*	0.72 (4)

C18A	0.224 (3)	0.5017 (14)	0.3172 (8)	0.067 (4)	0.28 (4)
H18D	0.2296	0.4320	0.3234	0.101*	0.28 (4)
H18E	0.1268	0.5272	0.3406	0.101*	0.28 (4)
H18F	0.3380	0.5298	0.3435	0.101*	0.28 (4)
C19A	0.061 (2)	0.4530 (13)	0.1931 (10)	0.051 (4)	0.28 (4)
H19D	0.0576	0.4581	0.1367	0.076*	0.28 (4)
H19E	-0.0578	0.4659	0.2041	0.076*	0.28 (4)
H19F	0.0989	0.3882	0.2108	0.076*	0.28 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0509 (2)	0.03275 (19)	0.0553 (2)	-0.00550 (15)	0.01586 (18)	0.00201 (15)
F1	0.089 (3)	0.098 (3)	0.055 (2)	0.031 (2)	0.032 (2)	0.0041 (17)
F2	0.0630 (17)	0.098 (4)	0.072 (3)	0.0147 (17)	-0.0148 (16)	0.020 (2)
F3	0.112 (4)	0.062 (2)	0.0513 (16)	-0.017 (3)	0.005 (3)	0.0106 (15)
F1A	0.084 (4)	0.102 (6)	0.061 (5)	0.024 (4)	0.041 (3)	0.013 (4)
F2A	0.098 (7)	0.070 (4)	0.034 (3)	0.003 (5)	-0.020 (4)	-0.012 (2)
F3A	0.081 (5)	0.045 (2)	0.045 (3)	0.007 (3)	-0.004 (4)	-0.0009 (19)
F1B	0.092 (5)	0.079 (7)	0.040 (5)	-0.015 (5)	0.015 (4)	0.007 (4)
F2B	0.090 (7)	0.093 (6)	0.040 (4)	-0.004 (6)	0.009 (6)	-0.006 (4)
F3B	0.095 (7)	0.088 (7)	0.083 (7)	0.036 (6)	-0.005 (6)	0.020 (5)
O1	0.0366 (5)	0.0292 (5)	0.0457 (6)	-0.0035 (4)	0.0102 (4)	-0.0017 (4)
N1	0.0462 (8)	0.0456 (7)	0.0770 (10)	-0.0004 (6)	0.0265 (7)	0.0158 (7)
C1	0.0277 (6)	0.0294 (6)	0.0373 (7)	-0.0005 (5)	0.0057 (5)	0.0019 (5)
C2	0.0257 (6)	0.0314 (6)	0.0391 (7)	0.0030 (5)	0.0071 (5)	-0.0029 (5)
C3	0.0318 (6)	0.0339 (6)	0.0383 (7)	0.0031 (5)	0.0077 (5)	-0.0024 (5)
C4	0.0403 (7)	0.0446 (8)	0.0378 (7)	0.0080 (6)	0.0063 (6)	-0.0045 (6)
C5	0.0520 (9)	0.0505 (9)	0.0451 (9)	-0.0004 (7)	0.0027 (7)	-0.0155 (7)
C6	0.0462 (8)	0.0374 (7)	0.0575 (10)	-0.0054 (6)	0.0083 (7)	-0.0130 (7)
C7	0.0323 (6)	0.0316 (6)	0.0458 (8)	0.0015 (5)	0.0097 (6)	-0.0028 (5)
C8	0.0369 (7)	0.0427 (8)	0.0428 (8)	-0.0026 (6)	0.0093 (6)	0.0024 (6)
C9	0.0581 (10)	0.0584 (10)	0.0568 (10)	-0.0140 (8)	0.0227 (8)	0.0040 (8)
C10	0.0771 (13)	0.0818 (14)	0.0547 (11)	-0.0166 (11)	0.0343 (10)	-0.0042 (10)
C11	0.0743 (13)	0.0736 (13)	0.0495 (10)	-0.0156 (10)	0.0252 (9)	-0.0170 (9)
C12	0.0485 (9)	0.0528 (9)	0.0429 (8)	-0.0109 (7)	0.0111 (7)	-0.0079 (7)
C13	0.0311 (6)	0.0413 (7)	0.0357 (7)	-0.0030 (5)	0.0045 (5)	0.0006 (5)
C14	0.0584 (10)	0.0599 (10)	0.0412 (8)	0.0141 (8)	0.0076 (7)	-0.0022 (7)
C15	0.0296 (6)	0.0377 (7)	0.0538 (9)	0.0010 (5)	0.0028 (6)	0.0079 (6)
C16	0.0329 (7)	0.0379 (8)	0.0779 (12)	0.0043 (6)	0.0158 (7)	0.0000 (7)
C17	0.0317 (7)	0.0475 (9)	0.0802 (12)	0.0003 (6)	0.0158 (8)	0.0068 (8)
C18	0.072 (4)	0.133 (6)	0.089 (3)	0.010 (4)	0.037 (3)	0.057 (3)
C19	0.111 (6)	0.055 (3)	0.147 (6)	-0.028 (4)	0.055 (4)	0.011 (3)
C18A	0.077 (9)	0.056 (6)	0.063 (6)	-0.027 (6)	0.001 (6)	0.000 (5)
C19A	0.039 (5)	0.042 (6)	0.072 (7)	-0.007 (4)	0.014 (4)	-0.007 (5)

Geometric parameters (\AA , \circ)

S1—C7	1.7535 (15)	C6—H6A	0.9500
S1—C8	1.7602 (16)	C8—C9	1.392 (2)
F1—C14	1.354 (3)	C8—C13	1.397 (2)
F2—C14	1.370 (5)	C9—C10	1.374 (3)
F3—C14	1.280 (5)	C9—H9A	0.9500
F1A—C14	1.360 (7)	C10—C11	1.381 (3)
F1A—F3A	2.102 (9)	C10—H10A	0.9500
F2A—C14	1.273 (8)	C11—C12	1.385 (2)
F3A—C14	1.357 (9)	C11—H11A	0.9500
F1B—C14	1.443 (10)	C12—C13	1.386 (2)
F2B—C14	1.237 (9)	C12—H12A	0.9500
F3B—C14	1.292 (10)	C15—C16	1.529 (2)
O1—C1	1.4134 (15)	C15—H15A	0.9900
O1—H1A	0.87 (2)	C15—H15B	0.9900
N1—C18A	1.429 (14)	C16—C17	1.520 (2)
N1—C19	1.439 (8)	C16—H16A	0.9900
N1—C17	1.456 (2)	C16—H16B	0.9900
N1—C18	1.481 (7)	C17—H17A	0.9900
N1—C19A	1.517 (12)	C17—H17B	0.9900
C1—C13	1.5213 (19)	C18—H18A	0.9800
C1—C2	1.5263 (19)	C18—H18B	0.9800
C1—C15	1.5489 (19)	C18—H18C	0.9800
C2—C3	1.3873 (19)	C19—H19A	0.9800
C2—C7	1.3989 (19)	C19—H19B	0.9800
C3—C4	1.386 (2)	C19—H19C	0.9800
C3—H3A	0.9500	C18A—H18D	0.9800
C4—C5	1.388 (2)	C18A—H18E	0.9800
C4—C14	1.486 (2)	C18A—H18F	0.9800
C5—C6	1.374 (2)	C19A—H19D	0.9800
C5—H5A	0.9500	C19A—H19E	0.9800
C6—C7	1.394 (2)	C19A—H19F	0.9800
C7—S1—C8	99.63 (7)	F3—C14—F1A	77.7 (5)
C1—O1—H1A	109.8 (15)	F3B—C14—F1A	122.8 (9)
C18A—N1—C19	101.1 (8)	F2B—C14—F2	64.3 (9)
C18A—N1—C17	118.6 (10)	F3A—C14—F2	87.4 (5)
C19—N1—C17	112.2 (6)	F3—C14—F2	106.0 (4)
C19—N1—C18	111.1 (5)	F1—C14—F2	101.1 (3)
C17—N1—C18	108.1 (6)	F3B—C14—F2	59.2 (9)
C18A—N1—C19A	107.5 (9)	F1A—C14—F2	130.2 (5)
C17—N1—C19A	103.5 (8)	F2B—C14—F1B	101.8 (7)
C18—N1—C19A	116.8 (9)	F3A—C14—F1B	73.0 (7)
O1—C1—C13	108.24 (11)	F2A—C14—F1B	131.2 (7)
O1—C1—C2	110.70 (11)	F3—C14—F1B	48.9 (7)
C13—C1—C2	108.84 (10)	F1—C14—F1B	68.4 (7)
O1—C1—C15	111.19 (11)	F3B—C14—F1B	99.1 (8)

C13—C1—C15	107.85 (11)	F2—C14—F1B	138.9 (6)
C2—C1—C15	109.92 (11)	F2B—C14—C4	116.4 (6)
C3—C2—C7	118.09 (13)	F3A—C14—C4	111.1 (6)
C3—C2—C1	120.40 (12)	F2A—C14—C4	115.3 (6)
C7—C2—C1	121.50 (12)	F3—C14—C4	116.1 (3)
C4—C3—C2	120.78 (13)	F1—C14—C4	111.4 (2)
C4—C3—H3A	119.6	F3B—C14—C4	113.8 (7)
C2—C3—H3A	119.6	F1A—C14—C4	110.6 (3)
C3—C4—C5	120.41 (15)	F2—C14—C4	111.3 (3)
C3—C4—C14	119.30 (15)	F1B—C14—C4	109.4 (4)
C5—C4—C14	120.27 (15)	C16—C15—C1	117.70 (13)
C6—C5—C4	119.75 (15)	C16—C15—H15A	107.9
C6—C5—H5A	120.1	C1—C15—H15A	107.9
C4—C5—H5A	120.1	C16—C15—H15B	107.9
C5—C6—C7	119.83 (15)	C1—C15—H15B	107.9
C5—C6—H6A	120.1	H15A—C15—H15B	107.2
C7—C6—H6A	120.1	C17—C16—C15	115.14 (14)
C6—C7—C2	121.02 (14)	C17—C16—H16A	108.5
C6—C7—S1	117.41 (11)	C15—C16—H16A	108.5
C2—C7—S1	121.55 (11)	C17—C16—H16B	108.5
C9—C8—C13	120.68 (15)	C15—C16—H16B	108.5
C9—C8—S1	116.99 (12)	H16A—C16—H16B	107.5
C13—C8—S1	122.30 (11)	N1—C17—C16	113.97 (13)
C10—C9—C8	119.86 (17)	N1—C17—H17A	108.8
C10—C9—H9A	120.1	C16—C17—H17A	108.8
C8—C9—H9A	120.1	N1—C17—H17B	108.8
C9—C10—C11	120.25 (17)	C16—C17—H17B	108.8
C9—C10—H10A	119.9	H17A—C17—H17B	107.7
C11—C10—H10A	119.9	N1—C18—H18A	109.5
C10—C11—C12	119.87 (17)	N1—C18—H18B	109.5
C10—C11—H11A	120.1	H18A—C18—H18B	109.5
C12—C11—H11A	120.1	N1—C18—H18C	109.5
C13—C12—C11	121.09 (16)	H18A—C18—H18C	109.5
C13—C12—H12A	119.5	H18B—C18—H18C	109.5
C11—C12—H12A	119.5	N1—C19—H19A	109.5
C12—C13—C8	118.20 (14)	N1—C19—H19B	109.5
C12—C13—C1	121.02 (13)	H19A—C19—H19B	109.5
C8—C13—C1	120.74 (13)	N1—C19—H19C	109.5
F2B—C14—F3A	130.8 (9)	H19A—C19—H19C	109.5
F3A—C14—F2A	106.6 (6)	H19B—C19—H19C	109.5
F2B—C14—F3	126.1 (7)	N1—C18A—H18D	109.5
F2A—C14—F3	119.6 (6)	N1—C18A—H18E	109.5
F3A—C14—F1	129.6 (6)	H18D—C18A—H18E	109.5
F2A—C14—F1	77.8 (5)	N1—C18A—H18F	109.5
F3—C14—F1	109.7 (3)	H18D—C18A—H18F	109.5
F2B—C14—F3B	113.9 (9)	H18E—C18A—H18F	109.5
F2A—C14—F3B	80.3 (9)	N1—C19A—H19D	109.5
F3—C14—F3B	51.2 (9)	N1—C19A—H19E	109.5

F1—C14—F3B	134.7 (7)	H19D—C19A—H19E	109.5
F2B—C14—F1A	73.7 (9)	N1—C19A—H19F	109.5
F3A—C14—F1A	101.4 (6)	H19D—C19A—H19F	109.5
F2A—C14—F1A	110.8 (6)	H19E—C19A—H19F	109.5
O1—C1—C2—C3	-16.56 (16)	F1A—F3A—C14—F3	-10 (2)
C13—C1—C2—C3	-135.40 (12)	F1A—F3A—C14—F1	28.2 (6)
C15—C1—C2—C3	106.68 (14)	F1A—F3A—C14—F3B	141.1 (17)
O1—C1—C2—C7	164.96 (11)	F1A—F3A—C14—F2	130.6 (5)
C13—C1—C2—C7	46.11 (16)	F1A—F3A—C14—F1B	-12.8 (7)
C15—C1—C2—C7	-71.81 (15)	F1A—F3A—C14—C4	-117.6 (5)
C7—C2—C3—C4	3.40 (19)	F3A—F1A—C14—F2B	-129.4 (9)
C1—C2—C3—C4	-175.14 (12)	F3A—F1A—C14—F2A	-112.9 (7)
C2—C3—C4—C5	-0.8 (2)	F3A—F1A—C14—F3	4.3 (8)
C2—C3—C4—C14	177.44 (13)	F3A—F1A—C14—F1	-143.5 (7)
C3—C4—C5—C6	-2.0 (2)	F3A—F1A—C14—F3B	-21.1 (10)
C14—C4—C5—C6	179.71 (15)	F3A—F1A—C14—F2	-96.4 (8)
C4—C5—C6—C7	2.2 (2)	F3A—F1A—C14—F1B	24.2 (13)
C5—C6—C7—C2	0.4 (2)	F3A—F1A—C14—C4	117.9 (6)
C5—C6—C7—S1	-177.87 (13)	C3—C4—C14—F2B	-153.9 (11)
C3—C2—C7—C6	-3.2 (2)	C5—C4—C14—F2B	24.4 (11)
C1—C2—C7—C6	175.29 (13)	C3—C4—C14—F3A	39.4 (6)
C3—C2—C7—S1	175.02 (10)	C5—C4—C14—F3A	-142.3 (6)
C1—C2—C7—S1	-6.46 (17)	C3—C4—C14—F2A	160.8 (7)
C8—S1—C7—C6	149.19 (12)	C5—C4—C14—F2A	-20.9 (7)
C8—S1—C7—C2	-29.12 (12)	C3—C4—C14—F3	13.7 (5)
C7—S1—C8—C9	-148.84 (14)	C5—C4—C14—F3	-168.1 (4)
C7—S1—C8—C13	29.48 (14)	C3—C4—C14—F1	-112.9 (4)
C13—C8—C9—C10	-0.8 (3)	C5—C4—C14—F1	65.4 (4)
S1—C8—C9—C10	177.53 (17)	C3—C4—C14—F3B	70.5 (11)
C8—C9—C10—C11	-0.9 (3)	C5—C4—C14—F3B	-111.2 (11)
C9—C10—C11—C12	0.9 (4)	C3—C4—C14—F1A	-72.4 (8)
C10—C11—C12—C13	0.7 (3)	C5—C4—C14—F1A	105.9 (8)
C11—C12—C13—C8	-2.3 (3)	C3—C4—C14—F2	135.1 (4)
C11—C12—C13—C1	175.65 (16)	C5—C4—C14—F2	-46.7 (4)
C9—C8—C13—C12	2.4 (2)	C3—C4—C14—F1B	-39.2 (8)
S1—C8—C13—C12	-175.88 (12)	C5—C4—C14—F1B	139.0 (8)
C9—C8—C13—C1	-175.60 (15)	O1—C1—C15—C16	64.51 (17)
S1—C8—C13—C1	6.1 (2)	C13—C1—C15—C16	-176.96 (12)
O1—C1—C13—C12	16.06 (18)	C2—C1—C15—C16	-58.43 (16)
C2—C1—C13—C12	136.43 (14)	C1—C15—C16—C17	-86.15 (18)
C15—C1—C13—C12	-104.34 (16)	C18A—N1—C17—C16	84.4 (9)
O1—C1—C13—C8	-166.03 (13)	C19—N1—C17—C16	-158.3 (5)
C2—C1—C13—C8	-45.65 (17)	C18—N1—C17—C16	78.8 (4)
C15—C1—C13—C8	73.57 (16)	C19A—N1—C17—C16	-156.7 (7)
F1A—F3A—C14—F2B	78.2 (12)	C15—C16—C17—N1	71.3 (2)
F1A—F3A—C14—F2A	116.0 (6)		

Hydrogen-bond geometry (Å, °)

Cg2 and Cg3 are the centroids of the C2–C7 and C8–C13 rings, respectively.

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1—H1 <i>A</i> ···N1	0.87 (2)	1.84 (2)	2.7141 (17)	176 (2)
C15—H15 <i>A</i> ···S1	0.99	2.76	3.4165 (15)	124
C5—H5 <i>A</i> ···Cg3 ⁱ	0.95	2.96	3.798 (3)	148
C17—H17 <i>A</i> ···Cg2 ⁱⁱ	0.99	2.97	3.949 (3)	170
C17—H17 <i>B</i> ···Cg3 ⁱⁱ	0.99	2.83	3.659 (3)	142

Symmetry codes: (i) $x-1/2, -y+1/2, z-1/2$; (ii) $x-1, y, z$.