

N-(3-Chlorophenyl)-*N'*-(3-methylphenyl)succinamide

 B. S. Saraswathi,^a Sabine Foro^b and B. Thimme Gowda^{a*}
^aDepartment of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, and ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany

Correspondence e-mail: gowdabt@yahoo.com


Received 29 June 2011; accepted 13 July 2011

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in main residue; R factor = 0.050; wR factor = 0.130; data-to-parameter ratio = 14.2.

The asymmetric unit of the title compound, $\text{C}_{17}\text{H}_{17}\text{ClN}_2\text{O}_2$, contains one half-molecule with a center of inversion at the mid-point of the central C—C bond. The amide N—H group is *anti* to the *meta*-chloro/methyl groups in the adjacent benzene rings. The dihedral angle between the benzene ring and the NH—C(O)—CH₂ segment is 43.5 (1)°. In the crystal, intermolecular N—H...O hydrogen bonds link the molecules into chains along the *a* axis. The methyl group and the Cl atom occupy the same position and were treated in a disorder model with site-occupation factors of 0.5 each.

Related literature

For our studies on the effects of substituents on the structures of *N*-(aryl)-amides, see: Bhat & Gowda (2000); Gowda *et al.* (2007); Saraswathi *et al.* (2011*a,b*) and on the structures of *N*-(aryl)-methanesulfonamides, see: Jayalakshmi & Gowda (2004). For similar structures, see: Pierrot *et al.* (1984). For restrained geometry, see: Nardelli (1999).



Experimental

Crystal data

 $\text{C}_{17}\text{H}_{17}\text{ClN}_2\text{O}_2$
 $M_r = 316.78$

 Triclinic, $P\bar{1}$
 $a = 4.840$ (1) Å
 $b = 5.560$ (1) Å
 $c = 14.752$ (3) Å
 $\alpha = 93.47$ (2)°
 $\beta = 91.39$ (2)°
 $\gamma = 97.71$ (2)°

 $V = 392.46$ (13) Å³
 $Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 0.25$ mm⁻¹
 $T = 293$ K
 $0.44 \times 0.20 \times 0.08$ mm

Data collection

 Oxford Diffraction Xcalibur diffractometer with Sapphire CCD detector
 Absorption correction: multi-scan (*CrysAlis RED*; Oxford)

 Diffraction, 2009)
 $T_{\min} = 0.897$, $T_{\max} = 0.980$
 2451 measured reflections
 1567 independent reflections
 1249 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.010$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.130$
 $S = 0.99$
 1567 reflections
 110 parameters

 14 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.17$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.17$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O1 ⁱ	0.86	2.05	2.894 (2)	168

 Symmetry code: (i) $x - 1, y, z$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

BSS thanks the University Grants Commission, Government of India, New Delhi, for the award of a research fellowship under its faculty improvement program.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2508).

References

- Bhat, D. K. & Gowda, B. T. (2000). *J. Indian Chem. Soc.* **77**, 279–284.
 Gowda, B. T., Foro, S. & Fuess, H. (2007). *Acta Cryst.* **E63**, o1975–o1976.
 Jayalakshmi, K. L. & Gowda, B. T. (2004). *Z. Naturforsch. Teil A*, **55**, 491–500.
 Nardelli, M. (1999). *J. Appl. Cryst.* **32**, 563–571.
 Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
 Pierrot, M., Baldy, A., Maire, J. C., Mehrotra, R. C., Kapoor, T. S. & Bachlas, B. P. (1984). *Acta Cryst.* **C40**, 1931–1934.
 Saraswathi, B. S., Foro, S. & Gowda, B. T. (2011*a*). *Acta Cryst.* **E67**, o966.
 Saraswathi, B. S., Foro, S. & Gowda, B. T. (2011*b*). *Acta Cryst.* **E67**, o1591.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2011). E67, o2077 [doi:10.1107/S1600536811028121]

N-(3-Chlorophenyl)-*N'*-(3-methylphenyl)succinamide

B. S. Saraswathi, Sabine Foro and B. Thimme Gowda

S1. Comment

The amide and sulfonamide moieties are important constituents of many biologically significant compounds. As part of our studies on the substituent effects on the structures of this class of compounds (Bhat & Gowda, 2000; Gowda *et al.*, 2007; Jayalakshmi & Gowda, 2004; Saraswathi *et al.*, 2011*a,b*), in the present work, the structure of *N*-(3-chlorophenyl),*N'*-(3-methylphenyl)-succinamide, (I), has been determined (Fig.1). The asymmetric unit of (I) contains half a molecule with a center of inversion at the mid-point of the central C—C bond, similar to that observed in bis(2-chlorophenylaminocarbonylmethyl)disulfide, (II), (Pierrot *et al.*, 1984), *N,N*-bis(3-chlorophenyl)-succinamide, (III), (Saraswathi *et al.*, 2011*a*) and *N,N*-bis(3-methylphenyl)-succinamide dihydrate, (IV), (Saraswathi *et al.*, 2011*b*).

The conformations of the amide O atoms are *anti* to the H atoms attached to the adjacent C atoms. Further, the conformations of the N—H bonds in the amide fragments are *anti* to the *meta*-chloro/methyl groups in the adjacent benzene rings, similar to the *anti* conformations observed with respect to the *meta*-chloro groups in (III) and *meta*-methyl groups in (IV).

Further, the C1—N1—C7—C8 and C1a—N1a—C7a—C8a segments in (I) are nearly planar and so also the C1—N1—C7—O1 and C1a—N1a—C7a—O1a segments, similar to those observed in (III) and (IV). The torsion angles of C2—C1—N1—C7 and C6—C1—N1—C7 are $-43.2(4)^\circ$ and $138.6(3)^\circ$, in contrast to the values of $-35.0(3)^\circ$ and $147.5(2)^\circ$ in (III), and $5.4(9)^\circ$ and $-173.6(6)^\circ$ in (IV).

The dihedral angle between the benzene ring and the NH—C(O)—CH₂ segment is $43.5(1)^\circ$, compared to the values of $62.1(2)^\circ$ in (III) and $5.6(4)^\circ$ in (IV).

The packing of the molecules in the crystal is accomplished by N—H⋯O hydrogen bonds (Table 1) that lead is shown in Fig. 2.

S2. Experimental

Succinic anhydride (0.01 mol) in toluene (25 ml) was treated dropwise with 3-chloroaniline (0.01 mol) also in toluene (20 ml) with constant stirring. The resulting mixture was stirred for one hour and set aside for an additional hour at room temperature for completion of the reaction. The mixture was then treated with dilute hydrochloric acid to remove unreacted 3-chloroaniline. The resultant solid *N*-(3-chlorophenyl)-succinamic acid was filtered under suction and washed thoroughly with water to remove the unreacted succinic anhydride and succinic acid. The compound was recrystallized to a constant melting point from ethanol. The purity of the compound was checked by elemental analysis and characterized by its infrared and NMR spectra.

The *N*-(3-chlorophenyl)succinamic acid obtained was then treated with phosphorous oxychloride and excess of 3-methylaniline at room temperature with constant stirring. The resultant mixture was stirred for 4 h, kept aside for additional 6 h for completion of the reaction and poured slowly into crushed ice with constant stirring. It was kept aside for a day. The resultant solid, *N*-(3-chlorophenyl), *N'*-(3-methylphenyl)-succinamide was filtered under suction, washed

thoroughly with water, dilute sodium hydroxide solution and finally with water. It was recrystallized to a constant melting point from a mixture of acetone and toluene (3:1 v/v). The compound was characterized by its infrared and NMR spectra.

Prism-like colorless single crystals used in X-ray diffraction studies were grown in a mixture of acetone and toluene (3:1 v/v) at room temperature.

S3. Refinement

The H atoms were positioned with idealized geometry using a riding model with C—H = 0.93 Å for aromatic, C—H = 0.97 Å for methylene and N—H = 0.86 Å for amide H atoms and were refined with isotropic displacement parameters, set to $1.2 \times U_{eq}$ of the parent atom. Atoms C9 and C11 occupy the same position. The disorder was treated by using a split-atom model. The corresponding site-occupation factors were fixed to 0.50:0.50. The U^{ij} components of these atoms were restrained to approximate isotropic behavior (Nardelli, 1999), the bond lengths C3—C9 and C3—C11 were restrained.





Figure 1

Molecular structure of the title compound, showing the atom labelling scheme and displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

Molecular packing of the title compound with hydrogen bonding shown as dashed lines.

***N*-(3-Chlorophenyl)-*N'*-(3-methylphenyl)succinamide**

Crystal data

$C_{17}H_{17}ClN_2O_2$

$M_r = 316.78$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 4.840\ (1)\ \text{\AA}$

$b = 5.560\ (1)\ \text{\AA}$

$c = 14.752\ (3)\ \text{\AA}$

$\alpha = 93.47\ (2)^\circ$

$\beta = 91.39\ (2)^\circ$

$\gamma = 97.71\ (2)^\circ$

$V = 392.46\ (13)\ \text{\AA}^3$

$Z = 1$

$F(000) = 166$

$D_x = 1.340\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1094 reflections

$\theta = 2.8\text{--}27.7^\circ$

$\mu = 0.25\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Prism, colourless

$0.44 \times 0.20 \times 0.08\ \text{mm}$

Data collection

Oxford Diffraction Xcalibur

diffractometer with Sapphire CCD detector

Radiation source: fine-focus sealed tube

Graphite monochromator

Rotation method data acquisition using ω scans

Absorption correction: multi-scan

(*CrysAlis RED*; Oxford Diffraction, 2009)

$T_{\min} = 0.897$, $T_{\max} = 0.980$

2451 measured reflections

1567 independent reflections

1249 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.010$

$\theta_{\max} = 26.3^\circ$, $\theta_{\min} = 2.8^\circ$

$h = -6 \rightarrow 5$

$k = -6 \rightarrow 6$

$l = -18 \rightarrow 17$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.050$

$wR(F^2) = 0.130$

$S = 0.99$

1567 reflections

110 parameters

14 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0501P)^2 + 0.2642P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.17 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.2418 (3)	0.7102 (3)	-0.09593 (13)	0.0651 (6)	
N1	-0.2109 (3)	0.5789 (3)	-0.13211 (12)	0.0417 (5)	
H1	-0.3807	0.5949	-0.1204	0.050*	
C1	-0.1722 (4)	0.4107 (4)	-0.20578 (14)	0.0384 (5)	
C2	0.0217 (4)	0.4712 (4)	-0.27063 (14)	0.0445 (5)	
H2	0.1310	0.6226	-0.2657	0.053*	
C3	0.0541 (5)	0.3084 (5)	-0.34254 (15)	0.0520 (6)	
C4	-0.1059 (6)	0.0831 (5)	-0.34971 (18)	0.0630 (7)	
H4A	-0.0831	-0.0283	-0.3977	0.076*	
C5	-0.2988 (6)	0.0250 (5)	-0.2854 (2)	0.0659 (7)	
H5A	-0.4070	-0.1269	-0.2904	0.079*	
C6	-0.3364 (5)	0.1866 (4)	-0.21353 (16)	0.0502 (6)	
H6	-0.4703	0.1453	-0.1709	0.060*	
C7	-0.0058 (4)	0.7190 (4)	-0.08329 (14)	0.0414 (5)	
C8	-0.0999 (4)	0.8842 (4)	-0.00851 (15)	0.0441 (5)	
H8A	-0.1160	0.7990	0.0470	0.053*	
H8B	-0.2828	0.9240	-0.0249	0.053*	
C9	0.251 (3)	0.364 (4)	-0.4221 (10)	0.155 (8)	0.50
H9A	0.3567	0.5224	-0.4105	0.185*	0.50
H9B	0.1426	0.3602	-0.4776	0.185*	0.50
H9C	0.3763	0.2442	-0.4274	0.185*	0.50
Cl1	0.2883 (5)	0.3904 (5)	-0.42162 (12)	0.0742 (6)	0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0252 (8)	0.0862 (13)	0.0794 (12)	0.0103 (7)	0.0031 (7)	-0.0380 (10)
N1	0.0249 (8)	0.0507 (11)	0.0484 (10)	0.0059 (7)	0.0052 (7)	-0.0101 (8)
C1	0.0295 (9)	0.0447 (12)	0.0415 (11)	0.0094 (8)	-0.0013 (8)	-0.0030 (9)
C2	0.0366 (11)	0.0497 (13)	0.0457 (12)	0.0028 (9)	0.0031 (9)	-0.0039 (9)
C3	0.0458 (12)	0.0695 (16)	0.0415 (12)	0.0152 (11)	0.0031 (10)	-0.0051 (11)
C4	0.0707 (17)	0.0637 (17)	0.0532 (14)	0.0147 (13)	0.0009 (12)	-0.0188 (12)

C5	0.0738 (18)	0.0468 (14)	0.0717 (17)	-0.0039 (12)	-0.0025 (14)	-0.0111 (12)
C6	0.0462 (12)	0.0493 (13)	0.0526 (13)	-0.0017 (10)	0.0059 (10)	-0.0003 (10)
C7	0.0283 (10)	0.0485 (12)	0.0471 (12)	0.0073 (8)	0.0046 (8)	-0.0065 (10)
C8	0.0302 (10)	0.0550 (13)	0.0458 (11)	0.0071 (9)	0.0048 (8)	-0.0114 (10)
C9	0.149 (9)	0.162 (9)	0.152 (9)	0.019 (5)	0.009 (5)	0.006 (5)
Cl1	0.0639 (9)	0.1109 (15)	0.0466 (7)	0.0093 (9)	0.0242 (7)	-0.0080 (8)

Geometric parameters (Å, °)

O1—C7	1.224 (2)	C4—H4A	0.9300
N1—C7	1.342 (3)	C5—C6	1.380 (3)
N1—C1	1.423 (3)	C5—H5A	0.9300
N1—H1	0.8591	C6—H6	0.9300
C1—C6	1.382 (3)	C7—C8	1.510 (3)
C1—C2	1.381 (3)	C8—C8 ⁱ	1.507 (4)
C2—C3	1.378 (3)	C8—H8A	0.9700
C2—H2	0.9300	C8—H8B	0.9700
C3—C4	1.379 (4)	C9—H9A	0.9600
C3—Cl1	1.686 (3)	C9—H9B	0.9600
C3—C9	1.550 (9)	C9—H9C	0.9600
C4—C5	1.369 (4)		
C7—N1—C1	125.39 (16)	C6—C5—H5A	119.2
C7—N1—H1	118.5	C5—C6—C1	119.0 (2)
C1—N1—H1	116.1	C5—C6—H6	120.5
C6—C1—C2	119.8 (2)	C1—C6—H6	120.5
C6—C1—N1	119.33 (19)	O1—C7—N1	122.90 (18)
C2—C1—N1	120.85 (19)	O1—C7—C8	121.54 (18)
C3—C2—C1	120.4 (2)	N1—C7—C8	115.51 (16)
C3—C2—H2	119.8	C7—C8—C8 ⁱ	112.1 (2)
C1—C2—H2	119.8	C7—C8—H8A	109.2
C2—C3—C4	120.0 (2)	C8 ⁱ —C8—H8A	109.2
C2—C3—Cl1	119.1 (2)	C7—C8—H8B	109.2
C4—C3—Cl1	120.9 (2)	C8 ⁱ —C8—H8B	109.2
C2—C3—C9	124.4 (7)	H8A—C8—H8B	107.9
C4—C3—C9	115.4 (7)	C3—C9—H9A	109.5
Cl1—C3—C9	5.9 (7)	C3—C9—H9B	109.5
C5—C4—C3	119.2 (2)	H9A—C9—H9B	109.5
C5—C4—H4A	120.4	C3—C9—H9C	109.5
C3—C4—H4A	120.4	H9A—C9—H9C	109.5
C4—C5—C6	121.6 (2)	H9B—C9—H9C	109.5
C4—C5—H5A	119.2		
C7—N1—C1—C6	138.7 (2)	C9—C3—C4—C5	-175.8 (9)
C7—N1—C1—C2	-43.0 (3)	C3—C4—C5—C6	-0.1 (4)
C6—C1—C2—C3	-0.6 (3)	C4—C5—C6—C1	-1.0 (4)
N1—C1—C2—C3	-178.9 (2)	C2—C1—C6—C5	1.3 (3)
C1—C2—C3—C4	-0.5 (4)	N1—C1—C6—C5	179.6 (2)

C1—C2—C3—C11	178.8 (2)	C1—N1—C7—O1	-2.4 (4)
C1—C2—C3—C9	175.8 (9)	C1—N1—C7—C8	179.8 (2)
C2—C3—C4—C5	0.8 (4)	O1—C7—C8—C8 ⁱ	32.9 (4)
C11—C3—C4—C5	-178.4 (2)	N1—C7—C8—C8 ⁱ	-149.2 (2)

Symmetry code: (i) $-x, -y+2, -z$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1...O1 ⁱⁱ	0.86	2.05	2.894 (2)	168

Symmetry code: (ii) $x-1, y, z$.