

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# catena-Poly[copper(II)-bis( $\mu$ -2-ethyl-5-methylimidazole-4-sulfonato- $\kappa^3N^3, O^4:O^4'$ )]

 Andrew P. Purdy<sup>a\*</sup> and Ray J. Butcher<sup>b</sup>

<sup>a</sup>Chemistry Division, Code 6120 Naval Research Laboratory, 4555 Overlook Avenue SW, Washington, DC 20375, USA, and <sup>b</sup>Department of Chemistry, Howard University, 525 College Street NW, Washington, DC 20059, USA  
Correspondence e-mail: andrew.purdy@nrl.navy.mil


Received 19 July 2011; accepted 19 August 2011

Key indicators: single-crystal X-ray study;  $T = 295$  K; mean  $\sigma(C-C) = 0.005$  Å;  $R$  factor = 0.064;  $wR$  factor = 0.181; data-to-parameter ratio = 14.6.

In the title compound,  $[Cu(C_6H_9N_2O_3S)_2]_n$ , the copper(II) ion sits on an inversion center and is chelated by the imidazole N and sulfonate O atoms of two ligands in equatorial positions. O atoms of adjacent molecules coordinate in the axial positions. Jahn–Teller tetragonal distortion is evident in the coordination geometry [Cu–N and Cu–O equatorial distances of 1.971 (3) and 2.045 (2) Å, respectively, with a Cu–O axial distance of 2.433 (3) Å]. The structure is propagated by an infinite chain of eight-membered (Cu–O–S–O)<sub>2</sub> ring systems along the  $a$  axis. Only N–H...O hydrogen bonding exists between the chains.

## Related literature

For literature related to the 2-ethyl-4-methylimidazole-5-sulfonic acid ligand, see: Purdy *et al.* (2007). For sulfonate-bridged Cu complexes with Cu–sulfonate chains, see: van Albada *et al.* (2001); Cai *et al.* (2004); Doyle *et al.* (1983); Han *et al.* (2006); He *et al.* (2009); Hubig *et al.* (2000); Sreenivasulu *et al.* (2005); Timmermans *et al.* (1984). For geometric data, see: Jahn & Teller (1937).



## Experimental

## Crystal data

$[Cu(C_6H_9N_2O_3S)_2]$   
 $M_r = 441.96$   
Monoclinic,  $P2_1/c$   
 $a = 5.0732$  (4) Å  
 $b = 11.8367$  (10) Å  
 $c = 13.6810$  (11) Å  
 $\beta = 94.473$  (7)°

$V = 819.04$  (12) Å<sup>3</sup>  
 $Z = 2$   
Cu  $K\alpha$  radiation  
 $\mu = 4.64$  mm<sup>-1</sup>  
 $T = 295$  K  
 $0.44 \times 0.32 \times 0.24$  mm

## Data collection

Oxford Diffraction Xcalibur Ruby Gemini diffractometer  
Absorption correction: analytical [*CrysAlis PRO* (Oxford Diffraction, 2007) based on expressions derived by Clark &

Reid (1995)  
 $T_{\min} = 0.270$ ,  $T_{\max} = 0.445$   
3114 measured reflections  
1704 independent reflections  
1605 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.033$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.064$   
 $wR(F^2) = 0.181$   
 $S = 1.08$   
1704 reflections

117 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 1.15$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.84$  e Å<sup>-3</sup>

Table 1

Selected geometric parameters (Å, °).

Cu1–N1	1.971 (3)	Cu1–O1 <sup>i</sup>	2.045 (2)
Cu1–O1	2.045 (2)	Cu1–O3 <sup>ii</sup>	2.433 (3)

 Symmetry codes: (i)  $-x + 1, -y + 1, -z + 1$ ; (ii)  $x - 1, y, z$ .

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2–H2C...O2 <sup>iii</sup>	0.86	1.95	2.784 (4)	164

 Symmetry code: (iii)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2007); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

RJB wishes to acknowledge the NSF-MRI program (grant No. CHE-0619278) for funds to purchase the diffractometer, and we thank the Office of Naval Research for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2453).

## References

- Albada, G. A. van, Mutikainen, I., Turpeinen, U. & Reedijk, J. (2001). *Inorg. Chim. Acta*, **324**, 273–277.  
Cai, J.-H., Jiang, Y.-M., Wang, X.-J. & Liu, Z.-M. (2004). *Acta Cryst. E* **60**, m1659–m1661.  
Clark, R. C. & Reid, J. S. (1995). *Acta Cryst. A* **51**, 887–897.  
Doyle, G., Eriksen, K. A. & van Engen, D. (1983). *Inorg. Chem.* **22**, 2892–2895.

- Han, J., Wu, H., Zhi, S., Li, Y. & Pan, Y. (2006). *Anal. Sci. X-ray Struct. Anal. Online*, **22**, x299–x300.
- He, Q.-T., Li, X.-P., Liu, Y., Yu, Z.-Q., Wang, W. & Su, C.-Y. (2009). *Angew. Chem. Int. Ed.* **48**, 6156–6159.
- Hubig, S. M., Lindeman, S. V. & Kochi, J. K. (2000). *Coord. Chem. Rev.* **200**, 831–873.
- Jahn, H. & Teller, E. (1937). *Proc. R. Soc. London Ser. A*, 220–235.
- Oxford Diffraction (2007). *CrysAlis PRO*. Oxford Diffraction Ltd, Abingdon, England.
- Purdy, A. P., Gilardi, R., Luther, J. & Butcher, R. J. (2007). *Polyhedron*, **26**, 3930–3938.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sreenivasulu, B., Vetrichelvan, M., Zhao, F., Gao, S. & Vittal, J. J. (2005). *Eur. J. Inorg. Chem.* pp. 4635–4645.
- Timmermans, P. J. J. A., Mackor, A., Spek, A. L. & Kojic-Prodic, B. (1984). *J. Organomet. Chem.* **276**, 287–295.

## supporting information

*Acta Cryst.* (2011). E67, m1303–m1304 [doi:10.1107/S160053681103409X]

**catena-Poly[copper(II)-bis( $\mu$ -2-ethyl-5-methylimidazole-4-sulfonato- $\kappa^3N^3, O^4:O^4$ )]**

Andrew P. Purdy and Ray J. Butcher

**S1. Comment**

In the title compound, the copper(II) ion sits on an inversion center and is chelated by the imidazole N1 and sulfonate O1 of two ligands. The two chelate rings on a Cu are 5-membered and co-planar. Two O3 O atoms of adjacent molecules coordinate the axial positions with the usual Jahn-Teller tetragonal distortion (Jahn & Teller, 1937) ((Cu—N and Cu—O equatorial distances of 1.971 (3) and 2.045 (2) Å, respectively, with a Cu—O axial distance of 2.433 (3) Å) and link the Cu atoms in an infinite chain of 8-membered (Cu—O—S—O)<sub>2</sub> rings along the *a* axis. A number of examples exist for catenated 8-membered rings of sulfonate bridged copper(I) ions - Doyle *et al.* (1983), Han *et al.* (2006), Timmermans *et al.* (1984), and Hubig *et al.* (2000). All previous examples have 4 or 5 coordinate copper(I) and edge-shared catenation between the rings. This infinite chain of (Cu—O—S—O)<sub>2</sub> rings is unique for copper(II), and its rings are corner shared and linear. The Cu coordination is nearly octahedral, with adjacent angles ranging from 84.92 (10) to 93.10 (10)°. Our silver(I) complex of the same ligand (Purdy, *et al.* (2007)) has edge shared 8-membered rings connected by a tetrahedral Ag atom. The CuO<sub>4</sub> moieties are planar, and are nearly perpendicular (85.00 (8)°) to a plane composed of the linear N—Cu—N units within a chain. Likewise the plane formed by the 5-membered Cu—N—C—S—O chelate rings forms a dihedral angle of 86.61 (8)° with the plane formed by the Cu and O3 atoms within a chain.

As noted above, although all Cu—O distances are within the ranges normally observed in sulfonate complexes, the Cu—O3 distance is 0.4 Å longer than Cu—O1 as is seen in bis( $\mu$ -2-((2-oxybenzylidene)amino)ethyl)sulfonato-diaquadicopper(II) dihydrate and other similar chelated copper(II) sulfonates (Sreenivasulu *et al.*, 2005; Cai *et al.*, 2004). Copper(II) sulfonates where the sulfonate is not part of a chelate ring tend to have Cu—O distance of 2.3 Å or greater as for example in bis( $\mu$ -hydroxo)-bis( $\mu$ -trifluoromethanesulfonato-*O,O'*)-bis(4,4-dimethyl-2,2'-bipyridine)-di-copper(II) (van Albada, *et al.*, 2001) and in a sulfonate bridged complex (He, *et al.*, 2009).


The O2 atoms of the sulfonates are hydrogen bonded to the hydrogen on N2 of the imidazole ring of an adjacent chain, at a N—O distance of 2.784 (4) Å. This interaction bonds the chains into a fully three-dimensional structure.

**S2. Experimental**

Both 1:1 and 1:1.5 solutions of the potassium salts of the 2-ethyl-4-methyl-imidazole-5-sulfonic acid were prepared by combining 1 g (5.25 mmol) of the free acid with 1 and 1.5 equivalents of KOH solution respectively, and diluting the solutions to 1M based on K<sup>+</sup>. (All solutions were made with distilled water.) Two test reactions were done in vials with a 1M solution of CuCl<sub>2</sub>·2H<sub>2</sub>O, and a 0.2 ml metered pipet was used for the additions. In reaction #1, 0.2 ml of CuCl<sub>2</sub> solution was combined with 0.4 ml of the 1:1 solution. In reaction #2, 0.4 ml of the CuCl<sub>2</sub> solution was combined with 0.6 ml of the 1:1.5 solution. Both solutions were heated to a boil and allowed to cool. Both reactions produced a pale green precipitate, but green crystals of the title compound grow in #2 only, over several days.


### S3. Refinement

H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with an N—H distance of 0.86 Å and C—H distances of 0.97 Å  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  and 0.96 Å for  $\text{CH}_3$  [ $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ ].



**Figure 1**

Diagram of  $\text{C}_{12}\text{H}_{18}\text{CuN}_4\text{O}_6\text{S}_2$  illustrating the atom numbering scheme used. Thermal displacement parameters are at the 30% probability level. [symmetry codes for labelled atoms: O1A, N1A,  $1-x, 1-y, 1-z$ ; O3B,  $x-1, y, z$ ; O3C,  $2-x, 1-y, 1-z$ ]

**Figure 2**

The molecular packing for  $C_{12}H_{18}CuN_4O_6S_2$  viewed down the  $c$  axis.

**catena-Poly[copper(II)-bis( $\mu$ -2-ethyl-5-methylimidazole-4-sulfonato- $\kappa^3N^3,O^4:O^4$ )]**

*Crystal data*

$[Cu(C_6H_9N_2O_3S)_2]$

$M_r = 441.96$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2ybc$

$a = 5.0732$  (4) Å

$b = 11.8367$  (10) Å

$c = 13.6810$  (11) Å

$\beta = 94.473$  (7)°

$V = 819.04$  (12) Å<sup>3</sup>

$Z = 2$

$F(000) = 454$

$D_x = 1.792$  Mg m<sup>-3</sup>

Cu  $K\alpha$  radiation,  $\lambda = 1.54178$  Å

Cell parameters from 2637 reflections

$\theta = 5.0$ – $77.1$ °

$\mu = 4.64$  mm<sup>-1</sup>

$T = 295$  K

Chunk, pale green-blue

$0.44 \times 0.32 \times 0.24$  mm

*Data collection*

Oxford Diffraction Xcalibur Ruby Gemini diffractometer  
 Radiation source: Enhance (Cu) X-ray sealed tube  
 Graphite monochromator  
 Detector resolution: 10.51 pixels mm<sup>-1</sup>  
 $\omega$  scans  
 Absorption correction: analytical (CrysAlis PRO; Oxford Diffraction, 2007; Clark & Reid, 1995)

$T_{\min} = 0.270$ ,  $T_{\max} = 0.445$   
 3114 measured reflections  
 1704 independent reflections  
 1605 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.033$   
 $\theta_{\max} = 77.6^\circ$ ,  $\theta_{\min} = 5.0^\circ$   
 $h = -6 \rightarrow 6$   
 $k = -14 \rightarrow 14$   
 $l = -17 \rightarrow 10$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.064$   
 $wR(F^2) = 0.181$   
 $S = 1.08$   
 1704 reflections  
 117 parameters  
 0 restraints  
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map  
 Hydrogen site location: inferred from neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.1339P)^2 + 0.7529P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 1.15 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.84 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.5000	0.5000	0.5000	0.0255 (3)
S1	0.94199 (14)	0.32615 (6)	0.53104 (5)	0.0252 (3)
O1	0.8041 (5)	0.4192 (2)	0.57814 (17)	0.0327 (6)
O2	0.8628 (6)	0.2154 (2)	0.56252 (19)	0.0402 (7)
O3	1.2247 (5)	0.3403 (2)	0.53856 (19)	0.0345 (6)
N1	0.6262 (5)	0.4204 (2)	0.38608 (19)	0.0251 (6)
N2	0.7482 (6)	0.3464 (2)	0.2509 (2)	0.0283 (6)
H2C	0.7537	0.3326	0.1894	0.034*
C1	0.5837 (6)	0.4210 (3)	0.2898 (2)	0.0257 (6)
C2	0.3917 (8)	0.4933 (3)	0.2308 (3)	0.0316 (8)
H2A	0.4281	0.5717	0.2474	0.038*
H2B	0.2151	0.4763	0.2492	0.038*
C3	0.3959 (10)	0.4793 (4)	0.1198 (3)	0.0466 (10)
H3A	0.2724	0.5309	0.0872	0.070*
H3B	0.3475	0.4032	0.1018	0.070*

H3C	0.5704	0.4949	0.1008	0.070*
C4	0.9057 (7)	0.2957 (3)	0.3252 (2)	0.0272 (6)
C5	0.8264 (7)	0.3429 (3)	0.4086 (2)	0.0263 (6)
C6	1.1174 (8)	0.2112 (3)	0.3088 (3)	0.0375 (8)
H6A	1.2247	0.1995	0.3689	0.056*
H6B	1.2257	0.2389	0.2595	0.056*
H6C	1.0376	0.1410	0.2875	0.056*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cu1	0.0248 (4)	0.0294 (5)	0.0230 (4)	0.0050 (2)	0.0057 (3)	-0.0033 (2)
S1	0.0256 (5)	0.0264 (5)	0.0241 (5)	0.0014 (3)	0.0055 (3)	0.0015 (2)
O1	0.0322 (12)	0.0400 (13)	0.0261 (11)	0.0090 (10)	0.0036 (9)	-0.0048 (9)
O2	0.0507 (16)	0.0371 (15)	0.0333 (13)	-0.0043 (12)	0.0073 (11)	0.0076 (10)
O3	0.0269 (12)	0.0368 (13)	0.0401 (14)	0.0023 (9)	0.0048 (10)	0.0012 (10)
N1	0.0245 (12)	0.0257 (13)	0.0258 (13)	0.0007 (10)	0.0066 (10)	-0.0016 (9)
N2	0.0323 (15)	0.0299 (13)	0.0235 (12)	0.0021 (11)	0.0079 (11)	-0.0032 (10)
C1	0.0267 (14)	0.0240 (14)	0.0273 (15)	-0.0009 (11)	0.0070 (11)	-0.0017 (11)
C2	0.0343 (18)	0.0326 (17)	0.0282 (17)	0.0050 (12)	0.0045 (14)	0.0017 (12)
C3	0.059 (3)	0.053 (2)	0.0269 (17)	0.012 (2)	0.0002 (17)	0.0004 (16)
C4	0.0287 (15)	0.0269 (14)	0.0269 (14)	0.0012 (12)	0.0071 (12)	-0.0021 (12)
C5	0.0293 (15)	0.0245 (14)	0.0256 (14)	0.0011 (12)	0.0050 (12)	0.0002 (11)
C6	0.0361 (18)	0.0340 (18)	0.0433 (19)	0.0102 (15)	0.0081 (15)	-0.0062 (15)

*Geometric parameters (Å, °)*

Cu1—N1 <sup>i</sup>	1.971 (3)	N2—C4	1.380 (4)
Cu1—N1	1.971 (3)	N2—H2C	0.8600
Cu1—O1	2.045 (2)	C1—C2	1.487 (5)
Cu1—O1 <sup>i</sup>	2.045 (2)	C2—C3	1.529 (5)
Cu1—O3 <sup>ii</sup>	2.433 (3)	C2—H2A	0.9700
Cu1—O3 <sup>iii</sup>	2.433 (3)	C2—H2B	0.9700
S1—O3	1.439 (3)	C3—H3A	0.9600
S1—O2	1.447 (3)	C3—H3B	0.9600
S1—O1	1.479 (2)	C3—H3C	0.9600
S1—C5	1.743 (3)	C4—C5	1.359 (4)
O3—Cu1 <sup>iv</sup>	2.433 (3)	C4—C6	1.497 (5)
N1—C1	1.318 (4)	C6—H6A	0.9600
N1—C5	1.385 (4)	C6—H6B	0.9600
N2—C1	1.352 (4)	C6—H6C	0.9600
N1 <sup>i</sup> —Cu1—N1	180.00 (9)	C4—N2—H2C	125.2
N1 <sup>i</sup> —Cu1—O1	95.08 (10)	N1—C1—N2	109.4 (3)
N1—Cu1—O1	84.92 (10)	N1—C1—C2	126.5 (3)
N1 <sup>i</sup> —Cu1—O1 <sup>i</sup>	84.92 (10)	N2—C1—C2	124.1 (3)
N1—Cu1—O1 <sup>i</sup>	95.08 (10)	C1—C2—C3	114.7 (3)
O1—Cu1—O1 <sup>i</sup>	180.000 (1)	C1—C2—H2A	108.6

N1 <sup>i</sup> —Cu1—O3 <sup>ii</sup>	88.44 (10)	C3—C2—H2A	108.6
N1—Cu1—O3 <sup>ii</sup>	91.56 (10)	C1—C2—H2B	108.6
O1—Cu1—O3 <sup>ii</sup>	86.90 (10)	C3—C2—H2B	108.6
O1 <sup>i</sup> —Cu1—O3 <sup>ii</sup>	93.10 (10)	H2A—C2—H2B	107.6
N1 <sup>i</sup> —Cu1—O3 <sup>iii</sup>	91.56 (10)	C2—C3—H3A	109.5
N1—Cu1—O3 <sup>iii</sup>	88.44 (10)	C2—C3—H3B	109.5
O1—Cu1—O3 <sup>iii</sup>	93.10 (10)	H3A—C3—H3B	109.5
O1 <sup>i</sup> —Cu1—O3 <sup>iii</sup>	86.90 (10)	C2—C3—H3C	109.5
O3 <sup>ii</sup> —Cu1—O3 <sup>iii</sup>	180.0	H3A—C3—H3C	109.5
O3—S1—O2	112.46 (16)	H3B—C3—H3C	109.5
O3—S1—O1	112.61 (15)	C5—C4—N2	104.3 (3)
O2—S1—O1	113.19 (16)	C5—C4—C6	131.6 (3)
O3—S1—C5	108.32 (16)	N2—C4—C6	124.1 (3)
O2—S1—C5	108.00 (16)	C4—C5—N1	110.2 (3)
O1—S1—C5	101.46 (15)	C4—C5—S1	131.3 (3)
S1—O1—Cu1	118.92 (14)	N1—C5—S1	118.4 (2)
S1—O3—Cu1 <sup>iv</sup>	131.33 (15)	C4—C6—H6A	109.5
C1—N1—C5	106.6 (3)	C4—C6—H6B	109.5
C1—N1—Cu1	138.6 (2)	H6A—C6—H6B	109.5
C5—N1—Cu1	114.7 (2)	C4—C6—H6C	109.5
C1—N2—C4	109.5 (3)	H6A—C6—H6C	109.5
C1—N2—H2C	125.2	H6B—C6—H6C	109.5
O3—S1—O1—Cu1	128.60 (17)	Cu1—N1—C1—C2	-1.4 (5)
O2—S1—O1—Cu1	-102.44 (19)	C4—N2—C1—N1	-0.5 (4)
C5—S1—O1—Cu1	13.0 (2)	C4—N2—C1—C2	178.2 (3)
N1 <sup>i</sup> —Cu1—O1—S1	167.66 (17)	N1—C1—C2—C3	177.2 (4)
N1—Cu1—O1—S1	-12.34 (17)	N2—C1—C2—C3	-1.2 (5)
O3 <sup>ii</sup> —Cu1—O1—S1	79.50 (17)	C1—N2—C4—C5	0.3 (4)
O3 <sup>iii</sup> —Cu1—O1—S1	-100.50 (17)	C1—N2—C4—C6	-177.9 (3)
O2—S1—O3—Cu1 <sup>iv</sup>	180.00 (18)	N2—C4—C5—N1	0.0 (4)
O1—S1—O3—Cu1 <sup>iv</sup>	-50.7 (2)	C6—C4—C5—N1	178.0 (3)
C5—S1—O3—Cu1 <sup>iv</sup>	60.7 (2)	N2—C4—C5—S1	-176.9 (3)
O1—Cu1—N1—C1	-170.2 (3)	C6—C4—C5—S1	1.1 (6)
O1 <sup>i</sup> —Cu1—N1—C1	9.8 (3)	C1—N1—C5—C4	-0.2 (4)
O3 <sup>ii</sup> —Cu1—N1—C1	103.0 (3)	Cu1—N1—C5—C4	-177.8 (2)
O3 <sup>iii</sup> —Cu1—N1—C1	-77.0 (3)	C1—N1—C5—S1	177.1 (2)
O1—Cu1—N1—C5	6.3 (2)	Cu1—N1—C5—S1	-0.5 (3)
O1 <sup>i</sup> —Cu1—N1—C5	-173.7 (2)	O3—S1—C5—C4	50.0 (4)
O3 <sup>ii</sup> —Cu1—N1—C5	-80.5 (2)	O2—S1—C5—C4	-72.1 (4)
O3 <sup>iii</sup> —Cu1—N1—C5	99.5 (2)	O1—S1—C5—C4	168.7 (3)
C5—N1—C1—N2	0.4 (4)	O3—S1—C5—N1	-126.7 (3)
Cu1—N1—C1—N2	177.1 (2)	O2—S1—C5—N1	111.3 (3)
C5—N1—C1—C2	-178.2 (3)	O1—S1—C5—N1	-8.0 (3)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $x-1, y, z$ ; (iii)  $-x+2, -y+1, -z+1$ ; (iv)  $x+1, y, z$ .



*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—H2C···O2 <sup>v</sup>	0.86	1.95	2.784 (4)	164

Symmetry code: (v)  $x, -y+1/2, z-1/2$ .