

## 4-(1,3-Benzothiazol-2-yl)-1,5-dimethyl-2-phenyl-1*H*-pyrazol-3(2*H*)-one

Imane Chakib,<sup>a</sup> Abdelfettah Zerzouf,<sup>a</sup> Youssef Kandri Rodi,<sup>b</sup> El Mokhtar Essassi<sup>a</sup> and Seik Weng Ng<sup>c,d\*</sup>

<sup>a</sup>Laboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, <sup>b</sup>Laboratoire de Chimie Organique Appliquée, Faculté des Sciences et Techniques, Université Sidi Mohamed Ben Abdallah, Fès, Morocco, <sup>c</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and <sup>d</sup>Chemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: seikweng@um.edu.my


Received 14 September 2011; accepted 15 September 2011

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.047;  $wR$  factor = 0.131; data-to-parameter ratio = 17.0.

The central five-membered ring of the title compound,  $\text{C}_{18}\text{H}_{15}\text{N}_3\text{OS}$ , is almost planar (r.m.s. deviation =  $0.028\text{ \AA}$ ) and the benzothiazole fused-ring system is close to coplanar with this ring [dihedral angle =  $6.1(1)^\circ$ ]. The phenyl substituent is twisted by  $62.5(1)^\circ$ .

### Related literature

For the structure of the reactant 4-(2,3-dihydro-1,3-benzothiazol-2-ylidene)-3-methyl-1-phenyl-1*H*-pyrazol-5(4*H*)-one, see: Chakibe *et al.* (2010).



### Experimental

#### Crystal data

$\text{C}_{18}\text{H}_{15}\text{N}_3\text{OS}$   
 $M_r = 321.39$   
Monoclinic,  $P2_1/c$   
 $a = 8.7428(2)\text{ \AA}$   
 $b = 25.7551(5)\text{ \AA}$   
 $c = 6.9660(1)\text{ \AA}$   
 $\beta = 97.460(1)^\circ$

$V = 1555.27(5)\text{ \AA}^3$   
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.22\text{ mm}^{-1}$   
 $T = 293\text{ K}$   
 $0.50 \times 0.10 \times 0.10\text{ mm}$

#### Data collection

Bruker APEXII diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.900$ ,  $T_{\max} = 0.979$

18953 measured reflections  
3569 independent reflections  
2418 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.053$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$   
 $wR(F^2) = 0.131$   
 $S = 1.01$   
3569 reflections

210 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.26\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.27\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors thank Université Sidi Mohamed Ben Abdallah, Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2331).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2005). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chakibe, I., Zerzouf, A., Essassi, E. M., Reichelt, M. & Reuter, H. (2010). *Acta Cryst. E66*, o1096.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

# supporting information

*Acta Cryst.* (2011). E67, o2700 [https://doi.org/10.1107/S1600536811037652]

## 4-(1,3-Benzothiazol-2-yl)-1,5-dimethyl-2-phenyl-1*H*-pyrazol-3(2*H*)-one

Imane Chakib, Abdelfettah Zerzouf, Youssef Kandri Rodi, El Mokhtar Essassi and Seik Weng Ng

### S1. Comment

In the study, the tertiary nitrogen atom of the five-membered ring of 4-(2,3-dihydro-1,3-benzothiazol-2-ylidene)-3-methyl-1-phenyl-1*H*-pyrazol-5(4*H*)-one (Chakibe *et al.*, 2010) is used to displace iodine from methyl iodide to give the title compound; the carbon–carbon double-bond in the reactant is consequently converted to a double bond (Scheme I, Fig. 1). The central five-membered ring and the benzothiazolyl fused-ring is nearly co-planar (dihedral angle 6.1 (1)°). The phenyl substituent is twisted by 62.5 (1)° with respect to the five-membered ring.

### S2. Experimental

To a solution of (*E*)-4-(2,3-dihydro-1,3-benzothiazol-2-ylidene)-3-methyl-1-phenyl-1*H*-pyrazol-5(4*H*)-one (1 g, 3.25 mmol) in DMF (50 ml) was added sodium carbonate (2.5 g, 23 mmol), tetra-*n*-butylammonium bromide (0.15 g, 1 mmol) and methyl iodide (7.1 g, 50 mmol). The mixture was stirred for 24 h. The solid material was removed by filtration and the solution was evaporated. The residue was washed with dichloromethane and hexane, and was recrystallized from ethanol to afford the title compound as colorless crystals.

### S3. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93–0.96 Å) and were included in the refinement in the riding model approximation, with *U*(H) set to 1.2–1.5*U*(C). Omitted from the refinement was the (0 2 0) reflection.




Figure 1

Thermal ellipsoid plot (Barbour, 2001) of C<sub>18</sub>H<sub>15</sub>N<sub>3</sub>OS at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

**4-(1,3-Benzothiazol-2-yl)-1,5-dimethyl-2-phenyl-1*H*-pyrazol- 3(2*H*)-one***Crystal data*

$C_{18}H_{15}N_3OS$   
 $M_r = 321.39$   
Monoclinic,  $P2_1/c$   
Hall symbol: -P 2ybc  
 $a = 8.7428 (2)$  Å  
 $b = 25.7551 (5)$  Å  
 $c = 6.9660 (1)$  Å  
 $\beta = 97.460 (1)^\circ$   
 $V = 1555.27 (5)$  Å<sup>3</sup>  
 $Z = 4$

$F(000) = 672$   
 $D_x = 1.373$  Mg m<sup>-3</sup>  
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 3894 reflections  
 $\theta = 2.5\text{--}24.6^\circ$   
 $\mu = 0.22$  mm<sup>-1</sup>  
 $T = 293$  K  
Prism, colorless  
 $0.50 \times 0.10 \times 0.10$  mm

*Data collection*

Bruker APEXII  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.900$ ,  $T_{\max} = 0.979$

18953 measured reflections  
3569 independent reflections  
2418 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.053$   
 $\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 2.4^\circ$   
 $h = -11 \rightarrow 11$   
 $k = -29 \rightarrow 33$   
 $l = -9 \rightarrow 9$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.047$   
 $wR(F^2) = 0.131$   
 $S = 1.01$   
3569 reflections  
210 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0644P)^2 + 0.296P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.26$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.27$  e Å<sup>-3</sup>

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.92125 (7)	0.46686 (2)	0.19736 (8)	0.04602 (18)
N1	0.8107 (2)	0.43170 (7)	0.5013 (3)	0.0485 (5)
N2	0.6696 (2)	0.60559 (6)	0.3813 (2)	0.0450 (4)
N3	0.6087 (2)	0.58549 (6)	0.5419 (2)	0.0436 (4)
O1	0.80712 (19)	0.57089 (6)	0.1499 (2)	0.0555 (4)
C1	0.9552 (2)	0.40265 (8)	0.2603 (3)	0.0435 (5)
C2	1.0365 (3)	0.36554 (8)	0.1694 (3)	0.0531 (6)
H2	1.0832	0.3740	0.0609	0.064*
C3	1.0459 (3)	0.31630 (9)	0.2442 (4)	0.0611 (7)
H3	1.1003	0.2910	0.1862	0.073*
C4	0.9756 (3)	0.30371 (9)	0.4045 (4)	0.0682 (7)
H4	0.9818	0.2698	0.4510	0.082*
C5	0.8967 (3)	0.34025 (9)	0.4969 (4)	0.0657 (7)

H5	0.8506	0.3313	0.6053	0.079*
C6	0.8869 (2)	0.39090 (8)	0.4253 (3)	0.0462 (5)
C7	0.8201 (2)	0.47347 (7)	0.3978 (3)	0.0399 (5)
C8	0.7531 (2)	0.56620 (8)	0.3030 (3)	0.0417 (5)
C9	0.7516 (2)	0.52318 (7)	0.4329 (3)	0.0387 (5)
C10	0.6676 (2)	0.53751 (7)	0.5781 (3)	0.0398 (5)
C11	0.6421 (3)	0.50940 (9)	0.7570 (3)	0.0514 (6)
H11A	0.5334	0.5064	0.7630	0.077*
H11B	0.6897	0.5282	0.8679	0.077*
H11C	0.6868	0.4754	0.7559	0.077*
C12	0.5534 (3)	0.62109 (9)	0.6788 (3)	0.0527 (6)
H12A	0.5437	0.6031	0.7972	0.079*
H12B	0.4546	0.6346	0.6253	0.079*
H12C	0.6252	0.6492	0.7045	0.079*
C13	0.5892 (3)	0.64466 (7)	0.2625 (3)	0.0414 (5)
C14	0.4314 (3)	0.64253 (9)	0.2115 (3)	0.0532 (6)
H14	0.3732	0.6167	0.2606	0.064*
C15	0.3615 (3)	0.67960 (10)	0.0859 (4)	0.0657 (7)
H15	0.2553	0.6787	0.0497	0.079*
C16	0.4484 (4)	0.71801 (10)	0.0139 (4)	0.0673 (8)
H16	0.4007	0.7428	-0.0705	0.081*
C17	0.6046 (4)	0.71958 (9)	0.0666 (4)	0.0627 (7)
H17	0.6629	0.7454	0.0175	0.075*
C18	0.6759 (3)	0.68310 (8)	0.1916 (3)	0.0508 (5)
H18	0.7820	0.6844	0.2281	0.061*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0543 (3)	0.0407 (3)	0.0427 (3)	0.0044 (2)	0.0050 (2)	0.0038 (2)
N1	0.0525 (11)	0.0414 (10)	0.0524 (11)	0.0046 (8)	0.0100 (9)	0.0068 (8)
N2	0.0576 (11)	0.0379 (9)	0.0397 (9)	0.0063 (8)	0.0079 (8)	0.0048 (7)
N3	0.0568 (11)	0.0403 (10)	0.0334 (9)	0.0041 (8)	0.0047 (8)	-0.0002 (7)
O1	0.0732 (11)	0.0503 (9)	0.0456 (9)	0.0109 (8)	0.0180 (8)	0.0067 (7)
C1	0.0397 (11)	0.0404 (11)	0.0480 (12)	-0.0006 (9)	-0.0033 (9)	0.0008 (9)
C2	0.0500 (13)	0.0493 (13)	0.0590 (14)	0.0057 (11)	0.0036 (11)	-0.0051 (11)
C3	0.0572 (15)	0.0471 (14)	0.0778 (17)	0.0078 (11)	0.0044 (13)	-0.0065 (12)
C4	0.0668 (17)	0.0390 (13)	0.099 (2)	0.0085 (12)	0.0103 (15)	0.0110 (13)
C5	0.0698 (16)	0.0469 (14)	0.0841 (18)	0.0074 (12)	0.0240 (14)	0.0183 (13)
C6	0.0429 (12)	0.0386 (11)	0.0560 (13)	0.0005 (9)	0.0026 (10)	0.0061 (9)
C7	0.0398 (11)	0.0391 (11)	0.0383 (10)	-0.0017 (9)	-0.0043 (8)	0.0018 (8)
C8	0.0485 (12)	0.0376 (11)	0.0381 (11)	0.0028 (9)	0.0016 (9)	-0.0011 (8)
C9	0.0424 (11)	0.0360 (10)	0.0353 (10)	0.0003 (9)	-0.0036 (8)	-0.0004 (8)
C10	0.0433 (11)	0.0389 (11)	0.0342 (10)	-0.0014 (9)	-0.0067 (8)	-0.0005 (8)
C11	0.0613 (14)	0.0524 (13)	0.0392 (11)	0.0003 (11)	0.0020 (10)	0.0049 (10)
C12	0.0629 (15)	0.0518 (13)	0.0438 (12)	0.0052 (11)	0.0077 (11)	-0.0078 (10)
C13	0.0561 (13)	0.0316 (10)	0.0365 (10)	0.0055 (9)	0.0052 (9)	-0.0033 (8)
C14	0.0567 (14)	0.0439 (12)	0.0582 (14)	0.0007 (11)	0.0043 (11)	0.0035 (10)

C15	0.0643 (16)	0.0681 (17)	0.0617 (16)	0.0190 (13)	-0.0025 (13)	0.0022 (13)
C16	0.098 (2)	0.0560 (15)	0.0487 (14)	0.0268 (15)	0.0123 (14)	0.0137 (11)
C17	0.094 (2)	0.0384 (13)	0.0596 (15)	0.0055 (13)	0.0241 (14)	0.0075 (11)
C18	0.0626 (14)	0.0399 (12)	0.0506 (13)	-0.0025 (11)	0.0100 (11)	-0.0024 (10)

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

S1—C1	1.727 (2)	C8—C9	1.432 (3)
S1—C7	1.755 (2)	C9—C10	1.376 (3)
N1—C7	1.304 (2)	C10—C11	1.483 (3)
N1—C6	1.386 (3)	C11—H11A	0.9600
N2—C8	1.401 (3)	C11—H11B	0.9600
N2—N3	1.399 (2)	C11—H11C	0.9600
N2—C13	1.428 (2)	C12—H12A	0.9600
N3—C10	1.350 (2)	C12—H12B	0.9600
N3—C12	1.451 (3)	C12—H12C	0.9600
O1—C8	1.227 (2)	C13—C18	1.377 (3)
C1—C2	1.392 (3)	C13—C14	1.381 (3)
C1—C6	1.395 (3)	C14—C15	1.382 (3)
C2—C3	1.369 (3)	C14—H14	0.9300
C2—H2	0.9300	C15—C16	1.381 (4)
C3—C4	1.382 (4)	C15—H15	0.9300
C3—H3	0.9300	C16—C17	1.368 (4)
C4—C5	1.375 (4)	C16—H16	0.9300
C4—H4	0.9300	C17—C18	1.374 (3)
C5—C6	1.395 (3)	C17—H17	0.9300
C5—H5	0.9300	C18—H18	0.9300
C7—C9	1.448 (3)		
C1—S1—C7	88.78 (10)	C8—C9—C7	122.65 (19)
C7—N1—C6	110.27 (18)	N3—C10—C9	109.58 (17)
C8—N2—N3	108.35 (15)	N3—C10—C11	120.60 (19)
C8—N2—C13	121.79 (16)	C9—C10—C11	129.77 (19)
N3—N2—C13	120.94 (17)	C10—C11—H11A	109.5
C10—N3—N2	108.24 (16)	C10—C11—H11B	109.5
C10—N3—C12	127.40 (17)	H11A—C11—H11B	109.5
N2—N3—C12	119.07 (16)	C10—C11—H11C	109.5
C2—C1—C6	121.6 (2)	H11A—C11—H11C	109.5
C2—C1—S1	128.56 (18)	H11B—C11—H11C	109.5
C6—C1—S1	109.83 (16)	N3—C12—H12A	109.5
C3—C2—C1	118.1 (2)	N3—C12—H12B	109.5
C3—C2—H2	120.9	H12A—C12—H12B	109.5
C1—C2—H2	120.9	N3—C12—H12C	109.5
C2—C3—C4	120.9 (2)	H12A—C12—H12C	109.5
C2—C3—H3	119.5	H12B—C12—H12C	109.5
C4—C3—H3	119.5	C18—C13—C14	121.0 (2)
C5—C4—C3	121.4 (2)	C18—C13—N2	117.5 (2)
C5—C4—H4	119.3	C14—C13—N2	121.43 (19)

C3—C4—H4	119.3	C15—C14—C13	118.7 (2)
C4—C5—C6	119.0 (2)	C15—C14—H14	120.7
C4—C5—H5	120.5	C13—C14—H14	120.7
C6—C5—H5	120.5	C14—C15—C16	120.4 (3)
N1—C6—C5	125.6 (2)	C14—C15—H15	119.8
N1—C6—C1	115.37 (18)	C16—C15—H15	119.8
C5—C6—C1	119.0 (2)	C17—C16—C15	120.1 (2)
N1—C7—C9	125.38 (19)	C17—C16—H16	120.0
N1—C7—S1	115.75 (15)	C15—C16—H16	120.0
C9—C7—S1	118.85 (15)	C16—C17—C18	120.3 (2)
O1—C8—N2	123.08 (18)	C16—C17—H17	119.9
O1—C8—C9	131.43 (19)	C18—C17—H17	119.9
N2—C8—C9	105.45 (17)	C17—C18—C13	119.6 (2)
C10—C9—C8	107.86 (17)	C17—C18—H18	120.2
C10—C9—C7	129.34 (18)	C13—C18—H18	120.2
C8—N2—N3—C10	7.4 (2)	N2—C8—C9—C10	1.0 (2)
C13—N2—N3—C10	155.18 (17)	O1—C8—C9—C7	-0.5 (4)
C8—N2—N3—C12	163.59 (18)	N2—C8—C9—C7	176.98 (18)
C13—N2—N3—C12	-48.6 (3)	N1—C7—C9—C10	1.4 (3)
C7—S1—C1—C2	-179.4 (2)	S1—C7—C9—C10	179.71 (16)
C7—S1—C1—C6	0.61 (15)	N1—C7—C9—C8	-173.65 (19)
C6—C1—C2—C3	1.2 (3)	S1—C7—C9—C8	4.7 (3)
S1—C1—C2—C3	-178.77 (17)	N2—N3—C10—C9	-6.8 (2)
C1—C2—C3—C4	0.4 (4)	C12—N3—C10—C9	-160.4 (2)
C2—C3—C4—C5	-1.3 (4)	N2—N3—C10—C11	171.05 (18)
C3—C4—C5—C6	0.5 (4)	C12—N3—C10—C11	17.4 (3)
C7—N1—C6—C5	-178.3 (2)	C8—C9—C10—N3	3.6 (2)
C7—N1—C6—C1	0.0 (3)	C7—C9—C10—N3	-172.02 (19)
C4—C5—C6—N1	179.4 (2)	C8—C9—C10—C11	-174.0 (2)
C4—C5—C6—C1	1.1 (4)	C7—C9—C10—C11	10.4 (4)
C2—C1—C6—N1	179.51 (19)	C8—N2—C13—C18	-74.5 (2)
S1—C1—C6—N1	-0.5 (2)	N3—N2—C13—C18	142.02 (19)
C2—C1—C6—C5	-2.0 (3)	C8—N2—C13—C14	101.9 (2)
S1—C1—C6—C5	177.98 (18)	N3—N2—C13—C14	-41.5 (3)
C6—N1—C7—C9	178.88 (19)	C18—C13—C14—C15	0.5 (3)
C6—N1—C7—S1	0.5 (2)	N2—C13—C14—C15	-175.8 (2)
C1—S1—C7—N1	-0.67 (16)	C13—C14—C15—C16	-0.2 (4)
C1—S1—C7—C9	-179.16 (17)	C14—C15—C16—C17	0.0 (4)
N3—N2—C8—O1	172.67 (19)	C15—C16—C17—C18	-0.2 (4)
C13—N2—C8—O1	25.2 (3)	C16—C17—C18—C13	0.5 (3)
N3—N2—C8—C9	-5.1 (2)	C14—C13—C18—C17	-0.7 (3)
C13—N2—C8—C9	-152.51 (18)	N2—C13—C18—C17	175.78 (19)
O1—C8—C9—C10	-176.5 (2)		