

4-(6-Chloroimidazo[1,2-*b*]pyridazin-3-yl)benzonitrile

Yiliang Zhao* and Clarissa K. L. Ng

Faculty of Pharmacy, Science Road, The University of Sydney, Sydney, Australia
Correspondence e-mail: yiliang.zhao@sydney.edu.au


Received 8 September 2011; accepted 16 September 2011

Key indicators: single-crystal X-ray study; $T = 150\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.001\text{ \AA}$; R factor = 0.043; wR factor = 0.141; data-to-parameter ratio = 94.8.

In the title compound, $\text{C}_{13}\text{H}_7\text{ClN}_4$, the imidazopyridazine ring system is essentially planar [maximum deviation 0.015 (1) \AA]. It is inclined to the benzene ring of the benzonitrile group by 11.31 (2) $^\circ$. In the crystal, molecules are linked via $\text{C}-\text{H}\cdots\text{Cl}$ and $\text{C}-\text{H}\cdots\text{N}$ interactions.

Related literature

For related structures, see Kia *et al.* (2009); Khan *et al.* (2010); Xue (2010); Zhao *et al.* (2009).



Experimental

Crystal data

$\text{C}_{13}\text{H}_7\text{ClN}_4$	$Z = 16$
$M_r = 254.68$	
Tetragonal, $I4_1/a$	Mo $K\alpha$ radiation
$a = 13.5513 (12)\text{ \AA}$	$\mu = 0.32\text{ mm}^{-1}$
$c = 24.566 (3)\text{ \AA}$	$T = 150\text{ K}$
$V = 4511.3 (7)\text{ \AA}^3$	$0.25 \times 0.15 \times 0.10\text{ mm}$

Data collection

Bruker SMART APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2008)
 $T_{\min} = 0.659$, $T_{\max} = 0.746$

140841 measured reflections
18107 independent reflections
11833 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.141$
 $S = 1.08$
18107 reflections
191 parameters

7 restraints
All H-atom parameters refined
 $\Delta\rho_{\max} = 0.85\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.34\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C2—H2 \cdots N1 ⁱ	1.08 (1)	2.60 (1)	3.3502 (7)	126 (1)
C3—H3 \cdots Cl1 ⁱⁱ	1.08 (1)	2.70 (1)	3.7389 (6)	161 (1)
C5—H5 \cdots N4 ⁱⁱⁱ	1.08 (1)	2.31 (1)	3.3341 (7)	157 (1)
C8—H8 \cdots N4 ⁱⁱⁱ	1.08 (1)	2.52 (1)	3.6049 (8)	177 (1)
C12—H12 \cdots N2	1.08 (1)	2.25 (1)	2.9975 (6)	125 (1)

Symmetry codes: (i) $x, y + 1, z$; (ii) $-y + \frac{3}{4}, x + \frac{1}{4}, z + \frac{1}{4}$; (iii) $-y + \frac{3}{4}, x - \frac{1}{4}, -z + \frac{3}{4}$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1999); software used to prepare material for publication: *SHELXL97* and *publCIF* (Westrip, 2010).

The authors would like to thank Professor Dai Hibbs from the Faculty of Pharmacy, the University of Sydney, for his kind and generous support of this research project.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5227).

References

- Bruker (2008). *SADABS*, *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Khan, F. N., Manivel, P., Prabakaran, K., Hathwar, V. R. & Akkurt, M. (2010). *Acta Cryst.* **E66**, o1081.
- Kia, R., Fun, H.-K. & Kargar, H. (2009). *Acta Cryst.* **E65**, o660–o661.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Xue, M. W. (2010). *Acta Cryst.* **E66**, o2006.
- Zhao, M. M., Li, Y. H., Wu, D. H. & Wan, Q. (2009). *Acta Cryst.* **E65**, o1261.

supporting information

Acta Cryst. (2011). E67, o2710 [https://doi.org/10.1107/S1600536811037901]

4-(6-Chloroimidazo[1,2-*b*]pyridazin-3-yl)benzonitrile

Yiliang Zhao and Clarissa K. L. Ng

S1. Comment

The title compound, C₁₃H₇C₁₁N₄, was newly synthesized, crystallized and analysed at high resolution sin theta/lambda 1.1. The bond lengths are as expected. The crystal packing reveals that in one unit cell, every four molecules stack in parallel position by a number of non-classical hydrogen bonds. Hydrogen atoms are fixed at 1.083 Å as determined from neutron diffraction, and the temperature factors were refined isotropically.

S2. Experimental

To a thick wall borosilicate glass vial were added 6-Chloro-3-iodo-imidazo[1,2-*b*]pyridazine (0.05 g, 1.0 equiv.), 4-cyanophenylboronic acid (0.032 g, 1.2 equiv.), Pd(PPh₃)₄ (0.02 g, 10 mol%), K₂CO₃ (0.037 g, 1.5 equiv.), and EtOH/H₂O (4 : 1, 5 ml). The vial was sealed with a silicon septum and the reaction mixture was pre-stirred for 1 min. The reaction mixture was subjected to microwave irradiation at 90 °C for 14 min at the maximum power of 300 W with sufficient stirring. The reaction mixture was cooled to room temperature and solvent was removed under reduced pressure. The crude product was purified by flash column chromatography, eluting with ethyl acetate/hexane (1 : 1) to give the titled compound as a bright yellow solid (0.038 g, 83 %).




Figure 1

The *ORTEP* view of the title compound with 50% probability displacement ellipsoids for non-H atoms.

4-(6-Chloroimidazo[1,2-*b*]pyridazin-3-yl)benzonitrile*Crystal data*

$C_{13}H_7ClN_4$
 $M_r = 254.68$
Tetragonal, $I4_1/a$
 $a = 13.5513 (12)$ Å
 $c = 24.566 (3)$ Å
 $V = 4511.3 (7)$ Å³
 $Z = 16$
 $F(000) = 2080$
none

$D_x = 1.500$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 500 reflections
 $\theta = 1.7\text{--}62.3^\circ$
 $\mu = 0.32$ mm⁻¹
 $T = 150$ K
Block, colourless
 $0.25 \times 0.15 \times 0.10$ mm

Data collection

Bruker SMART APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
Absorption correction: multi-scan
(SAINT; Bruker, 2008)
 $T_{\min} = 0.659$, $T_{\max} = 0.746$

140841 measured reflections
18107 independent reflections
11833 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$
 $\theta_{\max} = 62.3^\circ$, $\theta_{\min} = 1.7^\circ$
 $h = -33 \rightarrow 32$
 $k = -33 \rightarrow 31$
 $l = -58 \rightarrow 60$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.141$
 $S = 1.08$
18107 reflections
191 parameters
7 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
All H-atom parameters refined
 $w = 1/[\sigma^2(F_o^2) + (0.0756P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.85$ e Å⁻³
 $\Delta\rho_{\min} = -0.34$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Reflections -1 11 4, -2 4 0, -5 7 10, 4 4 0, 2 4 2, 1 8 13, -4 8 6, 1 3 16, -2 4 22, -1 10 1, -1 6 1, 0 0 24, -4 6 10, 4 5 11, -2 5 7 were omitted due to bad agreement statistics.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.39217 (3)	0.36640 (3)	0.159032 (17)	0.01678 (5)
C2	0.37908 (3)	0.45664 (3)	0.187490 (19)	0.01942 (6)
C3	0.36436 (3)	0.45225 (3)	0.242477 (19)	0.01919 (6)

C4	0.36372 (3)	0.35883 (3)	0.267513 (16)	0.01652 (5)
C5	0.35942 (3)	0.23262 (3)	0.319699 (16)	0.01807 (5)
C6	0.37353 (3)	0.19353 (3)	0.267716 (15)	0.01514 (5)
C7	0.37986 (3)	0.09040 (3)	0.251063 (15)	0.01514 (5)
C8	0.38792 (3)	0.01863 (3)	0.292379 (18)	0.01890 (6)
C9	0.38734 (4)	-0.08101 (3)	0.27984 (2)	0.02124 (7)
C10	0.38037 (3)	-0.11125 (3)	0.22543 (2)	0.01965 (6)
C11	0.37593 (3)	-0.04085 (3)	0.183904 (18)	0.01980 (6)
C12	0.37476 (3)	0.05927 (3)	0.196657 (16)	0.01772 (5)
C13	0.37580 (4)	-0.21460 (3)	0.21251 (2)	0.02413 (8)
N1	0.37014 (5)	-0.29780 (4)	0.20267 (3)	0.03142 (10)
N2	0.39085 (2)	0.27773 (2)	0.179828 (13)	0.01557 (4)
N3	0.37627 (2)	0.27594 (2)	0.234420 (13)	0.01445 (4)
N4	0.35305 (3)	0.33273 (3)	0.319431 (15)	0.01907 (5)
Cl1	0.413952 (9)	0.370824 (9)	0.089725 (5)	0.02143 (3)
H2	0.3846 (8)	0.5243 (4)	0.1643 (3)	0.031 (2)*
H3	0.3542 (7)	0.5204 (4)	0.2647 (4)	0.030 (2)*
H5	0.3569 (8)	0.1898 (7)	0.3569 (2)	0.034 (3)*
H8	0.3999 (8)	0.0435 (7)	0.33372 (15)	0.031 (2)*
H9	0.3914 (7)	-0.1341 (6)	0.3127 (3)	0.029 (2)*
H11	0.3662 (8)	-0.0628 (9)	0.14188 (16)	0.042 (3)*
H12	0.3662 (7)	0.1155 (5)	0.1657 (3)	0.030 (2)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.01593 (12)	0.01674 (12)	0.01767 (13)	0.00052 (9)	-0.00065 (9)	0.00152 (9)
C2	0.02065 (14)	0.01551 (12)	0.02209 (15)	0.00112 (10)	-0.00169 (11)	0.00123 (10)
C3	0.02072 (14)	0.01484 (12)	0.02199 (15)	0.00197 (10)	-0.00191 (11)	-0.00164 (10)
C4	0.01706 (12)	0.01544 (11)	0.01706 (12)	0.00196 (9)	-0.00169 (9)	-0.00230 (9)
C5	0.02051 (14)	0.01867 (13)	0.01504 (11)	0.00221 (10)	-0.00118 (10)	-0.00095 (9)
C6	0.01548 (11)	0.01474 (11)	0.01520 (11)	0.00104 (8)	-0.00108 (8)	-0.00048 (8)
C7	0.01486 (11)	0.01468 (11)	0.01587 (11)	0.00048 (8)	-0.00133 (8)	-0.00043 (8)
C8	0.02312 (15)	0.01589 (12)	0.01768 (13)	0.00058 (10)	-0.00269 (11)	0.00081 (10)
C9	0.02647 (18)	0.01571 (13)	0.02155 (15)	0.00072 (11)	-0.00397 (13)	0.00104 (11)
C10	0.02066 (14)	0.01475 (12)	0.02353 (16)	0.00046 (10)	-0.00355 (12)	-0.00161 (11)
C11	0.02297 (16)	0.01711 (13)	0.01932 (14)	-0.00005 (11)	-0.00236 (11)	-0.00292 (10)
C12	0.02068 (14)	0.01622 (12)	0.01625 (12)	-0.00001 (10)	-0.00164 (10)	-0.00097 (9)
C13	0.02702 (19)	0.01604 (14)	0.0293 (2)	0.00087 (12)	-0.00495 (16)	-0.00242 (13)
N1	0.0399 (3)	0.01696 (15)	0.0374 (3)	0.00031 (15)	-0.0067 (2)	-0.00451 (15)
N2	0.01569 (10)	0.01588 (10)	0.01514 (9)	0.00048 (7)	-0.00012 (7)	0.00018 (7)
N3	0.01422 (9)	0.01436 (9)	0.01476 (9)	0.00097 (7)	-0.00081 (7)	-0.00084 (7)
N4	0.02264 (13)	0.01842 (12)	0.01614 (11)	0.00297 (9)	-0.00162 (9)	-0.00302 (9)
Cl1	0.02372 (5)	0.02314 (5)	0.01744 (4)	0.00047 (3)	0.00083 (3)	0.00373 (3)

Geometric parameters (\AA , $\text{^{\circ}}$)

C1—N2	1.3058 (5)	C7—C12	1.4033 (5)
C1—C2	1.4197 (6)	C7—C8	1.4100 (6)
C1—Cl1	1.7291 (5)	C8—C9	1.3849 (6)
C2—C3	1.3668 (7)	C8—H8	1.0824 (10)
C2—H2	1.0819 (10)	C9—C10	1.4014 (7)
C3—C4	1.4074 (6)	C9—H9	1.0824 (10)
C3—H3	1.0816 (10)	C10—C11	1.3980 (7)
C4—N4	1.3315 (6)	C10—C13	1.4374 (6)
C4—N3	1.3970 (5)	C11—C12	1.3925 (6)
C5—N4	1.3593 (6)	C11—H11	1.0825 (10)
C5—C6	1.3957 (5)	C12—H12	1.0823 (10)
C5—H5	1.0827 (10)	C13—N1	1.1557 (7)
C6—N3	1.3847 (5)	N2—N3	1.3558 (5)
C6—C7	1.4588 (5)		
N2—C1—C2	126.74 (4)	C9—C8—C7	120.80 (4)
N2—C1—Cl1	114.80 (3)	C9—C8—H8	120.9 (6)
C2—C1—Cl1	118.45 (3)	C7—C8—H8	118.2 (6)
C3—C2—C1	117.87 (4)	C8—C9—C10	119.84 (4)
C3—C2—H2	124.6 (5)	C8—C9—H9	118.8 (6)
C1—C2—H2	117.5 (5)	C10—C9—H9	121.4 (6)
C2—C3—C4	118.16 (4)	C11—C10—C9	119.96 (4)
C2—C3—H3	118.7 (5)	C11—C10—C13	120.14 (4)
C4—C3—H3	123.2 (5)	C9—C10—C13	119.89 (4)
N4—C4—N3	110.92 (3)	C12—C11—C10	120.08 (4)
N4—C4—C3	131.17 (4)	C12—C11—H11	118.7 (7)
N3—C4—C3	117.91 (4)	C10—C11—H11	120.9 (7)
N4—C5—C6	112.52 (4)	C11—C12—C7	120.44 (4)
N4—C5—H5	122.4 (6)	C11—C12—H12	121.9 (5)
C6—C5—H5	125.0 (6)	C7—C12—H12	117.6 (5)
N3—C6—C5	103.77 (3)	N1—C13—C10	178.50 (7)
N3—C6—C7	127.26 (3)	C1—N2—N3	113.92 (3)
C5—C6—C7	128.92 (4)	N2—N3—C6	127.06 (3)
C12—C7—C8	118.82 (4)	N2—N3—C4	125.38 (3)
C12—C7—C6	123.52 (3)	C6—N3—C4	107.54 (3)
C8—C7—C6	117.62 (3)	C4—N4—C5	105.24 (3)
N2—C1—C2—C3	0.66 (7)	C8—C7—C12—C11	1.11 (6)
Cl1—C1—C2—C3	-178.41 (3)	C6—C7—C12—C11	-176.50 (4)
C1—C2—C3—C4	0.41 (6)	C11—C10—C13—N1	-112 (3)
C2—C3—C4—N4	178.65 (4)	C9—C10—C13—N1	67 (3)
C2—C3—C4—N3	-1.24 (6)	C2—C1—N2—N3	-0.74 (6)
N4—C5—C6—N3	0.34 (5)	C11—C1—N2—N3	178.35 (3)
N4—C5—C6—C7	-177.28 (4)	C1—N2—N3—C6	-178.44 (4)
N3—C6—C7—C12	-10.13 (6)	C1—N2—N3—C4	-0.23 (5)
C5—C6—C7—C12	166.96 (4)	C5—C6—N3—N2	178.38 (3)

N3—C6—C7—C8	172.24 (4)	C7—C6—N3—N2	−3.94 (6)
C5—C6—C7—C8	−10.67 (6)	C5—C6—N3—C4	−0.09 (4)
C12—C7—C8—C9	−2.26 (6)	C7—C6—N3—C4	177.58 (4)
C6—C7—C8—C9	175.48 (4)	N4—C4—N3—N2	−178.70 (4)
C7—C8—C9—C10	1.11 (7)	C3—C4—N3—N2	1.21 (6)
C8—C9—C10—C11	1.22 (7)	N4—C4—N3—C6	−0.19 (4)
C8—C9—C10—C13	−177.53 (5)	C3—C4—N3—C6	179.72 (4)
C9—C10—C11—C12	−2.37 (7)	N3—C4—N4—C5	0.39 (5)
C13—C10—C11—C12	176.38 (4)	C3—C4—N4—C5	−179.51 (4)
C10—C11—C12—C7	1.19 (7)	C6—C5—N4—C4	−0.46 (5)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C2—H2···N1 ⁱ	1.08 (1)	2.60 (1)	3.3502 (7)	126 (1)
C3—H3···Cl1 ⁱⁱ	1.08 (1)	2.70 (1)	3.7389 (6)	161 (1)
C5—H5···N4 ⁱⁱⁱ	1.08 (1)	2.31 (1)	3.3341 (7)	157 (1)
C8—H8···N4 ⁱⁱⁱ	1.08 (1)	2.52 (1)	3.6049 (8)	177 (1)
C12—H12···N2	1.08 (1)	2.25 (1)	2.9975 (6)	125 (1)

Symmetry codes: (i) $x, y+1, z$; (ii) $-y+3/4, x+1/4, z+1/4$; (iii) $-y+3/4, x-1/4, -z+3/4$.