

2-[*N*-(3-Amino-4-nitrophenyl)-carboximidoyl]phenol

Gholam Hossein Shahverdizadeh,^a‡ Seik Weng Ng,^{b,c}
Edward R. T. Tiekink^{b*} and Babak Mirtamizdoust^d

^aDepartment of Chemistry, Faculty of Science, Tabriz Branch, Islamic Azad University, Tabriz, PO Box 1655, Iran, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, ^cChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia, and ^dDepartment of Inorganic Chemistry, Faculty of Chemistry, University of Tabriz, Tabriz, Iran
Correspondence e-mail: edward.tiekink@gmail.com


Received 10 October 2011; accepted 11 October 2011

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.035; wR factor = 0.113; data-to-parameter ratio = 12.1.

The title compound, $C_{13}H_{11}N_3O_3$, is essentially planar (r.m.s. for the 19 non-H atoms = 0.031 Å), a conformation stabilized in part by intramolecular O—H···N and N—H···O hydrogen bonds. The configuration about the imine bond [1.2919 (12) Å] is *E*. The presence of N—H···O(nitro) hydrogen bonds leads to the formation of supramolecular tapes in the crystal structure. These are connected into layers by π – π interactions [centroid–centroid distance = 3.6046 (6) Å] occurring between the hydroxy- and amino-substituted benzene rings.

Related literature

For related work on Schiff bases, see: Prasath *et al.* (2010); Shahverdizadeh & Tiekink (2011). For specialized crystallization techniques, see: Harrowfield *et al.* (1996).



Experimental

Crystal data



$M_r = 257.25$

| | |
|------------------------------|--|
| Triclinic, $P\bar{1}$ | $V = 569.87 (5)\text{ \AA}^3$ |
| $a = 7.0961 (3)\text{ \AA}$ | $Z = 2$ |
| $b = 7.5168 (4)\text{ \AA}$ | Cu $K\alpha$ radiation |
| $c = 12.1627 (6)\text{ \AA}$ | $\mu = 0.92\text{ mm}^{-1}$ |
| $\alpha = 100.067 (4)^\circ$ | $T = 100\text{ K}$ |
| $\beta = 94.751 (4)^\circ$ | $0.25 \times 0.20 \times 0.15\text{ mm}$ |
| $\gamma = 115.011 (5)^\circ$ | |

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010)
 $T_{\min} = 0.748$, $T_{\max} = 1.000$

3654 measured reflections
2231 independent reflections
2105 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.008$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.113$
 $S = 1.07$
2231 reflections
184 parameters
3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.28\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.28\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|--|--------------|--------------------|-------------|----------------------|
| O1—H1 _o ···N1 | 0.86 (1) | 1.79 (1) | 2.5933 (10) | 154 (2) |
| N2—H1 _n ···O2 | 0.89 (1) | 2.06 (1) | 2.6542 (11) | 123 (1) |
| N2—H1 _n ···O2 ⁱ | 0.89 (1) | 2.42 (1) | 3.1479 (11) | 140 (1) |
| N2—H2 _n ···O3 ⁱⁱ | 0.86 (1) | 2.25 (1) | 3.0746 (10) | 161 (1) |

Symmetry codes: (i) $-x + 2$, $-y + 1$, $-z$; (ii) $x - 1$, y , z .

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors thank Tabriz Azad University and the University of Malaya for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG5110).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Harrowfield, J. M., Miyamae, H., Skelton, B. W., Soudi, A. A. & White, A. H. (1996). *Aust. J. Chem.* **49**, 1165–1169.
- Prasath, R., Bhavana, P., Ng, S. W. & Tiekink, E. R. T. (2010). *Acta Cryst. E66*, o3123.
- Shahverdizadeh, G. H. & Tiekink, E. R. T. (2011). *Acta Cryst. E67*, o798.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

‡ Additional correspondence author, e-mail: shahverdizadeh@iaut.ac.ir.

supporting information

Acta Cryst. (2011). E67, o2964 [doi:10.1107/S160053681104181X]

2-[*N*-(3-Amino-4-nitrophenyl)carboximidoyl]phenol

Gholam Hossein Shahverdizadeh, Seik Weng Ng, Edward R. T. Tiekkink and Babak Mirtamizdoust

S1. Comment

In continuation of structural studies of Schiff bases (Prasath *et al.*, 2010; Shahverdizadeh & Tiekkink, 2011), the title compound was synthesized and characterized crystallographically.

The molecule of (I), Fig. 1, is planar with the r.m.s. deviation of the 19 non-hydrogen atoms being 0.031 Å; the maximum and minimum deviations are 0.066 (1) Å for atom C12 and -0.062 (1) Å for atom O2. The observed planar conformation is stabilized in part by intramolecular O—H···N and N—H···O hydrogen bonds, Table 1. The amino group is planar with the sum of the angles about the N2 atom being approximately 359°. The configuration about the N1=C7 bond [1.2919 (12) Å] is *E*, and the hydroxy and amino groups are *syn*.


In the crystal structure, supramolecular tapes in the (1 1 1) plane are formed *via* N—H···O(nitro) hydrogen bonds, Fig. 2 and Table 1. The spine of the tape comprises alternating 12-membered rectangular {···HNC₂NO}₂ and square {···HNH···ONO}₂ synthons. The tapes are connected into layers *via* π–π interactions occurring between the hydroxy- and amino-benzene rings [centroid(C1–C6)···centroid(C8–C13)ⁱ distance = 3.6046 (6) Å for *i*: 1 - *x*, -*y*, 1 - *z*].

S2. Experimental


A solution of 4-nitrobenzene-1,3-diamine (10 mmol) in methanol (50 ml) was added drop wise to a solution of salicyl-aldehyde (10 mmol) in methanol (50 ml). The mixture was stirred for 5 h. The resulting solution was filtered to obtain a Schiff base, and dried. Single crystals of the title compound were obtained by using the branched tube method (Harrowfield *et al.*, 1996). Thus, the Schiff base (5 mmol) was placed in the arm to be heated. Methanol was added to fill both arms, and then the arm to be heated was placed in a bath at 333 K. After 2 days, orange crystals were deposited in the cooler arm, which were filtered, washed with water and air dried. Yield: 88%. *M.pt.*: 432 K.

S3. Refinement


Carbon-bound H-atoms were placed in calculated positions [C—H 0.95 Å, $U_{iso}(H) = 1.2U_{eq}(C)$] and were included in the refinement in the riding model approximation. The O—H and N—H H atoms were located from a difference map and refined with O—H = 0.84±0.01 Å and N—H = 0.88±0.01 Å, respectively, and with unconstrained $U_{iso}(H)$ values.

**Figure 1**

The molecular structure of (I) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.

**Figure 2**

A view of the supramolecular tape in the (1 1 1) plane and sustained by N—H···O hydrogen bonds (blue dashed lines) in the crystal structure of (I).

**Figure 3**

A view in projection down the a axis of the crystal packing in (I) highlighting the stacking of layers. The $\text{N}—\text{H}··\cdot\text{O}$ and $\pi—\pi$ interactions are shown as blue and purple dashed lines, respectively.

2-[*N*-(3-Amino-4-nitrophenyl)carboximidoyl]phenol

Crystal data

$\text{C}_{13}\text{H}_{11}\text{N}_3\text{O}_3$
 $M_r = 257.25$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 7.0961 (3)$ Å
 $b = 7.5168 (4)$ Å
 $c = 12.1627 (6)$ Å
 $\alpha = 100.067 (4)^\circ$
 $\beta = 94.751 (4)^\circ$
 $\gamma = 115.011 (5)^\circ$
 $V = 569.87 (5)$ Å³

$Z = 2$
 $F(000) = 268$
 $D_x = 1.499 \text{ Mg m}^{-3}$
Cu $K\alpha$ radiation, $\lambda = 1.54184$ Å
Cell parameters from 2525 reflections
 $\theta = 3.8—74.1^\circ$
 $\mu = 0.92 \text{ mm}^{-1}$
 $T = 100$ K
Prism, orange
 $0.25 \times 0.20 \times 0.15$ mm

Data collection

Agilent SuperNova Dual
diffractometer with an Atlas detector
Radiation source: SuperNova (Cu) X-ray
Source
Mirror monochromator
Detector resolution: 10.4041 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2010)

$T_{\min} = 0.748, T_{\max} = 1.000$
3654 measured reflections
2231 independent reflections
2105 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.008$
 $\theta_{\max} = 74.3^\circ, \theta_{\min} = 3.8^\circ$
 $h = -8 \rightarrow 8$
 $k = -9 \rightarrow 9$
 $l = -13 \rightarrow 15$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.035$$

$$wR(F^2) = 0.113$$

$$S = 1.07$$

2231 reflections

184 parameters

3 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0753P)^2 + 0.0839P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^* / U_{\text{eq}}$ |
|-----|--------------|--------------|-------------|------------------------------------|
| O1 | 0.40452 (11) | 0.27962 (11) | 0.59127 (6) | 0.0227 (2) |
| H1o | 0.495 (2) | 0.285 (3) | 0.5475 (13) | 0.062 (5)* |
| O2 | 1.09653 (11) | 0.36753 (12) | 0.05261 (6) | 0.0246 (2) |
| O3 | 1.31365 (10) | 0.29051 (11) | 0.14605 (6) | 0.0233 (2) |
| N1 | 0.71614 (12) | 0.25239 (11) | 0.50498 (7) | 0.0158 (2) |
| N2 | 0.78293 (13) | 0.40302 (13) | 0.14327 (7) | 0.0191 (2) |
| H1n | 0.840 (2) | 0.420 (2) | 0.0816 (9) | 0.039 (4)* |
| H2n | 0.6610 (15) | 0.4021 (18) | 0.1447 (10) | 0.026 (3)* |
| N3 | 1.15808 (12) | 0.32599 (12) | 0.13890 (7) | 0.0172 (2) |
| C1 | 0.47038 (15) | 0.22517 (14) | 0.68163 (8) | 0.0168 (2) |
| C2 | 0.36032 (15) | 0.21048 (14) | 0.77211 (8) | 0.0197 (2) |
| H2 | 0.2408 | 0.2372 | 0.7690 | 0.024* |
| C3 | 0.42483 (16) | 0.15713 (14) | 0.86643 (8) | 0.0209 (2) |
| H3 | 0.3500 | 0.1491 | 0.9281 | 0.025* |
| C4 | 0.59855 (15) | 0.11503 (14) | 0.87207 (8) | 0.0202 (2) |
| H4 | 0.6405 | 0.0763 | 0.9366 | 0.024* |
| C5 | 0.70904 (15) | 0.13020 (14) | 0.78289 (8) | 0.0182 (2) |
| H5 | 0.8283 | 0.1031 | 0.7870 | 0.022* |
| C6 | 0.64775 (14) | 0.18514 (13) | 0.68626 (8) | 0.0155 (2) |
| C7 | 0.76768 (14) | 0.20111 (13) | 0.59454 (8) | 0.0161 (2) |
| H7 | 0.8864 | 0.1733 | 0.6004 | 0.019* |
| C8 | 0.83502 (14) | 0.27202 (13) | 0.41602 (8) | 0.0146 (2) |
| C9 | 0.76291 (14) | 0.32620 (13) | 0.32488 (8) | 0.0151 (2) |
| H9 | 0.6412 | 0.3491 | 0.3265 | 0.018* |

| | | | | |
|-----|--------------|--------------|-------------|------------|
| C10 | 0.86365 (14) | 0.34913 (13) | 0.22862 (8) | 0.0150 (2) |
| C11 | 1.04545 (14) | 0.31485 (13) | 0.23212 (8) | 0.0150 (2) |
| C12 | 1.12150 (14) | 0.26527 (14) | 0.32711 (8) | 0.0168 (2) |
| H12 | 1.2463 | 0.2472 | 0.3281 | 0.020* |
| C13 | 1.01929 (15) | 0.24257 (14) | 0.41788 (8) | 0.0172 (2) |
| H13 | 1.0711 | 0.2076 | 0.4812 | 0.021* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|------------|------------|------------|------------|------------|------------|
| O1 | 0.0209 (4) | 0.0344 (4) | 0.0206 (4) | 0.0173 (3) | 0.0055 (3) | 0.0120 (3) |
| O2 | 0.0274 (4) | 0.0366 (4) | 0.0191 (4) | 0.0190 (3) | 0.0095 (3) | 0.0143 (3) |
| O3 | 0.0206 (4) | 0.0328 (4) | 0.0259 (4) | 0.0180 (3) | 0.0105 (3) | 0.0104 (3) |
| N1 | 0.0162 (4) | 0.0170 (4) | 0.0136 (4) | 0.0067 (3) | 0.0031 (3) | 0.0038 (3) |
| N2 | 0.0185 (4) | 0.0274 (4) | 0.0183 (4) | 0.0143 (3) | 0.0067 (3) | 0.0102 (3) |
| N3 | 0.0166 (4) | 0.0173 (4) | 0.0184 (4) | 0.0077 (3) | 0.0051 (3) | 0.0047 (3) |
| C1 | 0.0167 (5) | 0.0163 (4) | 0.0164 (5) | 0.0068 (4) | 0.0018 (4) | 0.0036 (3) |
| C2 | 0.0183 (5) | 0.0190 (4) | 0.0219 (5) | 0.0088 (4) | 0.0054 (4) | 0.0028 (4) |
| C3 | 0.0241 (5) | 0.0193 (5) | 0.0166 (5) | 0.0066 (4) | 0.0082 (4) | 0.0032 (4) |
| C4 | 0.0240 (5) | 0.0201 (4) | 0.0146 (5) | 0.0079 (4) | 0.0023 (4) | 0.0052 (3) |
| C5 | 0.0182 (5) | 0.0179 (5) | 0.0181 (5) | 0.0076 (4) | 0.0023 (4) | 0.0048 (3) |
| C6 | 0.0157 (4) | 0.0141 (4) | 0.0149 (5) | 0.0054 (3) | 0.0019 (3) | 0.0027 (3) |
| C7 | 0.0153 (5) | 0.0160 (4) | 0.0170 (5) | 0.0071 (4) | 0.0031 (4) | 0.0036 (4) |
| C8 | 0.0135 (4) | 0.0140 (4) | 0.0146 (5) | 0.0049 (3) | 0.0030 (3) | 0.0021 (3) |
| C9 | 0.0140 (4) | 0.0157 (4) | 0.0163 (5) | 0.0073 (3) | 0.0030 (4) | 0.0033 (4) |
| C10 | 0.0153 (4) | 0.0132 (4) | 0.0153 (5) | 0.0055 (3) | 0.0025 (3) | 0.0031 (3) |
| C11 | 0.0148 (4) | 0.0153 (4) | 0.0151 (5) | 0.0063 (4) | 0.0046 (4) | 0.0039 (3) |
| C12 | 0.0138 (4) | 0.0179 (4) | 0.0195 (5) | 0.0079 (4) | 0.0025 (3) | 0.0038 (3) |
| C13 | 0.0171 (4) | 0.0207 (5) | 0.0152 (5) | 0.0094 (4) | 0.0019 (3) | 0.0056 (3) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|--------|-------------|---------|-------------|
| O1—C1 | 1.3519 (11) | C4—C5 | 1.3830 (13) |
| O1—H1o | 0.858 (9) | C4—H4 | 0.9500 |
| O2—N3 | 1.2424 (10) | C5—C6 | 1.4064 (13) |
| O3—N3 | 1.2403 (10) | C5—H5 | 0.9500 |
| N1—C7 | 1.2919 (12) | C6—C7 | 1.4494 (13) |
| N1—C8 | 1.4163 (12) | C7—H7 | 0.9500 |
| N2—C10 | 1.3471 (12) | C8—C9 | 1.3772 (13) |
| N2—H1n | 0.885 (9) | C8—C13 | 1.4145 (13) |
| N2—H2n | 0.864 (8) | C9—C10 | 1.4200 (12) |
| N3—C11 | 1.4346 (12) | C9—H9 | 0.9500 |
| C1—C2 | 1.3947 (14) | C10—C11 | 1.4172 (13) |
| C1—C6 | 1.4115 (14) | C11—C12 | 1.4079 (13) |
| C2—C3 | 1.3840 (14) | C12—C13 | 1.3668 (13) |
| C2—H2 | 0.9500 | C12—H12 | 0.9500 |
| C3—C4 | 1.3957 (15) | C13—H13 | 0.9500 |
| C3—H3 | 0.9500 | | |

| | | | |
|---------------|-------------|-----------------|-------------|
| C1—O1—H1o | 104.2 (12) | C5—C6—C7 | 119.76 (9) |
| C7—N1—C8 | 121.52 (8) | C1—C6—C7 | 121.50 (9) |
| C10—N2—H1n | 122.2 (10) | N1—C7—C6 | 121.24 (9) |
| C10—N2—H2n | 117.9 (8) | N1—C7—H7 | 119.4 |
| H1n—N2—H2n | 119.2 (12) | C6—C7—H7 | 119.4 |
| O3—N3—O2 | 121.55 (8) | C9—C8—C13 | 120.23 (9) |
| O3—N3—C11 | 118.76 (8) | C9—C8—N1 | 115.89 (8) |
| O2—N3—C11 | 119.68 (8) | C13—C8—N1 | 123.87 (8) |
| O1—C1—C2 | 118.68 (9) | C8—C9—C10 | 122.45 (8) |
| O1—C1—C6 | 121.45 (9) | C8—C9—H9 | 118.8 |
| C2—C1—C6 | 119.87 (9) | C10—C9—H9 | 118.8 |
| C3—C2—C1 | 120.17 (9) | N2—C10—C11 | 125.47 (8) |
| C3—C2—H2 | 119.9 | N2—C10—C9 | 118.58 (8) |
| C1—C2—H2 | 119.9 | C11—C10—C9 | 115.95 (8) |
| C2—C3—C4 | 120.79 (9) | C12—C11—C10 | 121.18 (9) |
| C2—C3—H3 | 119.6 | C12—C11—N3 | 117.08 (8) |
| C4—C3—H3 | 119.6 | C10—C11—N3 | 121.73 (8) |
| C5—C4—C3 | 119.36 (9) | C13—C12—C11 | 121.27 (9) |
| C5—C4—H4 | 120.3 | C13—C12—H12 | 119.4 |
| C3—C4—H4 | 120.3 | C11—C12—H12 | 119.4 |
| C4—C5—C6 | 121.07 (9) | C12—C13—C8 | 118.89 (9) |
| C4—C5—H5 | 119.5 | C12—C13—H13 | 120.6 |
| C6—C5—H5 | 119.5 | C8—C13—H13 | 120.6 |
| C5—C6—C1 | 118.73 (9) | | |
| | | | |
| O1—C1—C2—C3 | -179.34 (8) | N1—C8—C9—C10 | -179.14 (7) |
| C6—C1—C2—C3 | 0.12 (15) | C8—C9—C10—N2 | -179.84 (8) |
| C1—C2—C3—C4 | -0.76 (15) | C8—C9—C10—C11 | -0.63 (14) |
| C2—C3—C4—C5 | 1.05 (15) | N2—C10—C11—C12 | 177.90 (8) |
| C3—C4—C5—C6 | -0.70 (15) | C9—C10—C11—C12 | -1.25 (13) |
| C4—C5—C6—C1 | 0.07 (14) | N2—C10—C11—N3 | -3.06 (15) |
| C4—C5—C6—C7 | 179.57 (8) | C9—C10—C11—N3 | 177.79 (7) |
| O1—C1—C6—C5 | 179.66 (7) | O3—N3—C11—C12 | 0.51 (13) |
| C2—C1—C6—C5 | 0.22 (14) | O2—N3—C11—C12 | 179.41 (7) |
| O1—C1—C6—C7 | 0.17 (15) | O3—N3—C11—C10 | -178.57 (8) |
| C2—C1—C6—C7 | -179.27 (8) | O2—N3—C11—C10 | 0.32 (13) |
| C8—N1—C7—C6 | 178.78 (7) | C10—C11—C12—C13 | 1.95 (14) |
| C5—C6—C7—N1 | -179.59 (7) | N3—C11—C12—C13 | -177.14 (8) |
| C1—C6—C7—N1 | -0.10 (15) | C11—C12—C13—C8 | -0.71 (14) |
| C7—N1—C8—C9 | 179.58 (7) | C9—C8—C13—C12 | -1.16 (14) |
| C7—N1—C8—C13 | -1.46 (15) | N1—C8—C13—C12 | 179.92 (8) |
| C13—C8—C9—C10 | 1.85 (14) | | |

Hydrogen-bond geometry (Å, °)

| D—H···A | D—H | H···A | D···A | D—H···A |
|-------------|----------|----------|-------------|---------|
| O1—H1o···N1 | 0.86 (1) | 1.79 (1) | 2.5933 (10) | 154 (2) |

| | | | | |
|------------------------------------|----------|----------|-------------|---------|
| N2—H1 <i>n</i> ···O2 | 0.89 (1) | 2.06 (1) | 2.6542 (11) | 123 (1) |
| N2—H1 <i>n</i> ···O2 ⁱ | 0.89 (1) | 2.42 (1) | 3.1479 (11) | 140 (1) |
| N2—H2 <i>n</i> ···O3 ⁱⁱ | 0.86 (1) | 2.25 (1) | 3.0746 (10) | 161 (1) |

Symmetry codes: (i) $-x+2, -y+1, -z$; (ii) $x-1, y, z$.