

1-(3,4-Dichlorobenzyl)pyridinium bis(2-sulfanylidene-1,3-dithiole-4,5-dithiolato- $\kappa^2 S,S'$)nickelate(III)

Guang-Xiang Liu

School of Biochemical and Environmental Engineering, Nanjing Xiaozhuang University, Nanjing 211171, People's Republic of China
 Correspondence e-mail: njuliugx@gmail.com


Received 8 October 2011; accepted 14 October 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.030; wR factor = 0.082; data-to-parameter ratio = 16.2.

The title compound, $(\text{C}_{12}\text{H}_{10}\text{Cl}_2\text{N})[\text{Ni}(\text{C}_3\text{S}_5)_2]$, is an ion-pair complex consisting of 1-(3,4-dichlorobenzyl)pyridinium cations and $[\text{Ni}(\text{dmit})_2]$ anions ($\text{dmit} = 2\text{-sulfanylidene-1,3-dithiole-4,5-dithiolate}$). In the anion, the Ni^{III} ion exhibits a square-planar coordination involving four S atoms from two dmit ligands. In the crystal, weak $\text{S}\cdots\text{S}$ [3.368 (2) and 3.482 (3) \AA], $\text{Ni}\cdots\text{S}$ [3.680 (2) \AA] and $\text{Cl}\cdots\text{S}$ [3.491 (2) \AA] interactions and $\text{C}-\text{H}\cdots\text{S}$ hydrogen bonds lead to a three-dimensional supramolecular network.

Related literature

For general background to the network topologies and applications of bis(dithiolate)-metal complexes, see: Cassoux (1999). For the synthesis, structures and properties of related complexes containing dmit ligands, see: Akutagawa & Nakamura (2000); Liu *et al.* (2010); Li *et al.* (2006); Zang *et al.* (2006, 2009). For the synthesis of a starting material, see: Wang *et al.* (1998).



Experimental

Crystal data

$(\text{C}_{12}\text{H}_{10}\text{Cl}_2\text{N})[\text{Ni}(\text{C}_3\text{S}_5)_2]$

$M_r = 690.48$

Triclinic, $P\bar{1}$

$a = 9.3711 (11)\text{ \AA}$

$b = 11.7210 (14)\text{ \AA}$

$c = 11.9640 (14)\text{ \AA}$

$\alpha = 82.814 (1)^\circ$

$\beta = 88.854 (1)^\circ$

$\gamma = 76.644 (1)^\circ$

$V = 1268.5 (3)\text{ \AA}^3$

$Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.81\text{ mm}^{-1}$

$T = 293\text{ K}$
 $0.22 \times 0.20 \times 0.16\text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.692$, $T_{\max} = 0.761$

9520 measured reflections
 4692 independent reflections
 4083 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.082$
 $S = 1.04$
 4692 reflections

290 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.38\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.34\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C14—H14 \cdots S10 ⁱ	0.93	2.82	3.622 (3)	145
C18—H18 \cdots S1 ⁱⁱ	0.93	2.79	3.708 (3)	168

Symmetry codes: (i) $x + 1, y, z$; (ii) $x, y + 1, z$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the National Natural Science Foundation of China (grant No. 20971004), the Key Project of the Chinese Ministry of Education (grant No. 210102) and the Natural Science Foundation of Anhui Province (grant No. 11040606M45).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2651).

References

- Akutagawa, T. & Nakamura, T. (2000). *Coord. Chem. Rev.* **198**, 297–311.
- Bruker (2000). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cassoux, P. (1999). *Coord. Chem. Rev.* **185–186**, 213–232.
- Li, J., Yao, L., Su, Y. & Tao, R. (2006). *Acta Cryst. E* **62**, m1990–m1991.
- Liu, G.-X., Yang, H., Guo, W., Liu, Y., Huang, R.-Y., Nishihara, S. & Ren, X.-M. (2010). *Polyhedron*, **29**, 2916–2923.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Wang, C., Batsanov, A. S., Bryce, M. R. & Howard, J. A. K. (1998). *Synthesis*, pp. 1615–1618.
- Zang, S.-Q., Ren, X.-M., Su, Y., Song, Y., Tong, W.-J., Ni, Z.-P., Zhao, H.-H., Gao, S. & Meng, Q.-J. (2009). *Inorg. Chem.* **48**, 9623–9630.
- Zang, S.-Q., Su, Y. & Tao, R.-J. (2006). *Acta Cryst. E* **62**, m1004–m1005.

supporting information

Acta Cryst. (2011). E67, m1570 [doi:10.1107/S160053681104267X]

1-(3,4-Dichlorobenzyl)pyridinium bis(2-sulfanylidene-1,3-dithiole-4,5-dithiolato- κ^2S,S')nickelate(III)

Guang-Xiang Liu

S1. Comment

Extensive research has been focused on the synthesis and characterization of bis(dithiolate)-metal complexes and their analogues, due to their properties and potential applications as conducting, magnetic and non-linear optical (NLO) materials (Cassoux, 1999). 2-Thioxo-1,3-dithiole-4,5-dithiolate (dmit) metal complexes are in fact excellent building blocks employed for the construction of molecular magnetic materials (Li *et al.*, 2006; Liu *et al.*, 2010; Zang *et al.*, 2006, 2009) apart from their well known electric conductivity as molecular conductors (Akutagawa & Nakamura, 2000).

Herein the crystal structure of the title compound, a new ion-pair complex, is reported.


The title compound comprises $[\text{Ni}(\text{dmit})_2]^-$ anions and 1-(3,4-dichlorobenzyl)pyridinium cations (Fig. 1). The Ni ion adopts a square-planar geometry coordinated by four S atoms from two dmit ligands, with Ni—S bond lengths ranging from 2.1518 (7) to 2.1714 (7) Å. The $[\text{Ni}(\text{dmit})_2]^-$ anions are in a parallel arrangement, with S···S interactions ranging from 3.474 (3) to 3.547 (3) Å. Two neighbouring anions are parallel in a face-to-face fashion with the shortest Ni···S distance of 3.680 (2) Å (Ni1—S2ⁱ) [symmetry code: (i) -x, -y, -z], indicating the existence of the Ni···S interactions. Adjacent $[\text{Ni}(\text{dmit})_2]^-$ anions are associated together through such Ni···S interactions resulting in a dimer. The dimers are linked together through S9···S3ⁱⁱ and S9···S5ⁱⁱ [symmetry code: (ii) x, 1 + y, z] interactions forming a one-dimensional chain structure, as depicted in Fig. 2. The $(\text{C}_{12}\text{H}_{10}\text{Cl}_2\text{N})^+$ cation has a Λ-shaped conformation, and the dihedral angles formed by the C12/C13/N1 plane with the benzene and pyridinium rings are 85.29 (2) and 77.84 (2)^o, respectively. Cations and the anions are linked by S···Cl interactions and C—H···S hydrogen bonds to generate a three-dimensional supramolecular structure (Fig. 3).

S2. Experimental

4,5-Di(thiobenzoyl)-1,3-dithiole-2-thione (812 mg, 2 mmol; Wang *et al.*, 1998) was suspended in methanol (10 ml). Sodium methoxide in methanol (prepared form 184 mg of sodium in 10 ml of methanol) was added to the above mixture under argon atmosphere at room temperature from 30 min to give a dark red solution. To this solution, $\text{NiCl}_2 \sim 2 \sim 6\text{H}_2\tilde{\text{O}}$ (238 mg, 1 mmol) was added. After 30 min, a solution of I $\sim 2 \sim$ (127 mg, 1 mmol) and NaI (150 mg, 1 mmol) in methanol (20 ml) was added (the monoanionic $[\text{Ni}(\text{dmit})_2]^-$ are obtained from the dianionic $[\text{Ni}(\text{dmit})_2]^{2-}$ by I $\sim 3 \sim$ oxidation). After another 10 min, a solution of 1-(3,4-dichlorobenzyl)pyridinium bromide [(DiClPy)Br] (317 mg, 1 mmol) in methanol (20 ml) was added to the reaction mixture. The solution was stirred for 30 min and cooled in a refrigerator overnight. The resultant dark green crystalline solid was collected by filtration, and purified by recrystallization using a mixed solution of acetonitrile and benzene (1:1 *v/v*).

S3. Refinement


H atoms were positioned geometrically, with C—H = 0.93 and 0.97 Å for aromatic and methylene H atoms, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.

**Figure 1**

The cation and anion in $[\text{DiClPy}][\text{Ni}(\text{dmit})\sim 2\sim]$, showing thermal ellipsoids drawn at the 30% probability level. Hydrogen atoms have been omitted for clarity.

**Figure 2**

The one-dimensional chain structure of $[\text{Ni}(\text{dmit})\sim 2\sim]$ - anions through S···S and Ni···S contacts. Dashed lines indicate weak interactions.

**Figure 3**

Packing of $[\text{DiClPy}][\text{Ni}(\text{dmit})_{\sim 2 \sim}]$ viewed along the b axis.

1-(3,4-Dichlorobenzyl)pyridinium bis(2-sulfanylidene-1,3-dithiole-4,5-dithiolato- κ^2S,S')nickelate(III)

Crystal data



$M_r = 690.48$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.3711 (11) \text{ \AA}$

$b = 11.7210 (14) \text{ \AA}$

$c = 11.9640 (14) \text{ \AA}$

$\alpha = 82.814 (1)^\circ$

$\beta = 88.854 (1)^\circ$

$\gamma = 76.644 (1)^\circ$

$V = 1268.5 (3) \text{ \AA}^3$

$Z = 2$

$F(000) = 694$

$D_x = 1.808 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5426 reflections

$\theta = 2.2\text{--}27.4^\circ$

$\mu = 1.81 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, black

$0.22 \times 0.20 \times 0.16 \text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector
diffractometer

Radiation source: sealed tube

Graphite monochromator

phi and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2000)

$T_{\min} = 0.692$, $T_{\max} = 0.761$

9520 measured reflections

4692 independent reflections

4083 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\max} = 25.5^\circ$, $\theta_{\min} = 1.7^\circ$

$h = -11 \rightarrow 11$

$k = -14 \rightarrow 14$

$l = -14 \rightarrow 14$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.030$$

$$wR(F^2) = 0.082$$

$$S = 1.04$$

4692 reflections

290 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.037P)^2 + 0.5172P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.38 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.34 \text{ e \AA}^{-3}$$

Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0118 (8)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.17029 (3)	0.03411 (2)	0.11584 (2)	0.03211 (11)
S1	0.32210 (7)	-0.05513 (5)	-0.00222 (5)	0.03971 (16)
S2	0.05550 (7)	-0.10606 (5)	0.14274 (5)	0.04056 (16)
S3	0.09306 (7)	-0.33353 (5)	0.04221 (5)	0.03969 (16)
S4	0.34360 (8)	-0.29296 (6)	-0.08879 (6)	0.04871 (18)
S5	0.24451 (11)	-0.51768 (7)	-0.09332 (7)	0.0698 (3)
S6	0.28357 (7)	0.17691 (5)	0.09083 (6)	0.03971 (16)
S7	0.01554 (7)	0.11932 (5)	0.23353 (6)	0.04310 (17)
S8	-0.00090 (8)	0.34602 (6)	0.33578 (6)	0.04481 (17)
S9	0.23172 (7)	0.40488 (5)	0.19675 (6)	0.03994 (16)
S10	0.09387 (8)	0.56899 (6)	0.35872 (6)	0.04947 (19)
C1	0.2605 (3)	-0.1815 (2)	-0.00797 (19)	0.0345 (5)
C2	0.1445 (3)	-0.20223 (19)	0.05421 (19)	0.0325 (5)
C3	0.2283 (3)	-0.3877 (2)	-0.0501 (2)	0.0436 (6)
C4	0.1888 (3)	0.27107 (19)	0.1800 (2)	0.0333 (5)
C5	0.0760 (3)	0.2457 (2)	0.2430 (2)	0.0355 (5)
C6	0.1081 (3)	0.4452 (2)	0.3007 (2)	0.0363 (5)
C7	0.6079 (3)	0.9937 (2)	0.3192 (2)	0.0501 (7)
H7	0.5323	1.0062	0.2667	0.060*
C8	0.6073 (3)	1.0755 (2)	0.3933 (2)	0.0485 (6)
C9	0.7202 (3)	1.0565 (2)	0.4706 (2)	0.0498 (7)
C10	0.8302 (3)	0.9562 (3)	0.4754 (3)	0.0578 (8)
H10	0.9047	0.9426	0.5290	0.069*

C11	0.8304 (3)	0.8754 (2)	0.4010 (2)	0.0512 (7)
H11	0.9062	0.8080	0.4036	0.061*
C12	0.7197 (3)	0.8940 (2)	0.3231 (2)	0.0445 (6)
C13	0.7260 (4)	0.8061 (3)	0.2393 (2)	0.0553 (8)
H13A	0.6883	0.8484	0.1671	0.066*
H13B	0.8274	0.7663	0.2292	0.066*
C14	0.6972 (3)	0.6255 (2)	0.3536 (2)	0.0453 (6)
H14	0.7872	0.6219	0.3868	0.054*
C15	0.6241 (3)	0.5380 (2)	0.3842 (2)	0.0483 (6)
H15	0.6653	0.4742	0.4371	0.058*
C16	0.4906 (3)	0.5447 (3)	0.3368 (2)	0.0522 (7)
H16	0.4399	0.4858	0.3569	0.063*
C17	0.4328 (3)	0.6394 (3)	0.2594 (3)	0.0576 (8)
H17	0.3413	0.6460	0.2272	0.069*
C18	0.5084 (3)	0.7237 (2)	0.2294 (2)	0.0532 (7)
H18	0.4691	0.7873	0.1760	0.064*
Cl1	0.72732 (13)	1.15957 (9)	0.56077 (8)	0.0904 (3)
Cl2	0.46624 (11)	1.19991 (8)	0.38728 (9)	0.0858 (3)
N1	0.6396 (2)	0.71609 (18)	0.27623 (16)	0.0396 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.03753 (19)	0.02357 (17)	0.03463 (18)	-0.00516 (12)	0.00213 (13)	-0.00512 (12)
S1	0.0444 (4)	0.0296 (3)	0.0457 (4)	-0.0097 (3)	0.0117 (3)	-0.0064 (3)
S2	0.0457 (4)	0.0316 (3)	0.0478 (4)	-0.0123 (3)	0.0151 (3)	-0.0138 (3)
S3	0.0490 (4)	0.0300 (3)	0.0430 (4)	-0.0122 (3)	0.0052 (3)	-0.0103 (3)
S4	0.0604 (4)	0.0377 (4)	0.0490 (4)	-0.0094 (3)	0.0197 (3)	-0.0155 (3)
S5	0.1137 (7)	0.0433 (4)	0.0602 (5)	-0.0236 (4)	0.0240 (5)	-0.0297 (4)
S6	0.0433 (4)	0.0307 (3)	0.0478 (4)	-0.0105 (3)	0.0124 (3)	-0.0134 (3)
S7	0.0522 (4)	0.0326 (3)	0.0487 (4)	-0.0155 (3)	0.0160 (3)	-0.0123 (3)
S8	0.0539 (4)	0.0346 (3)	0.0484 (4)	-0.0112 (3)	0.0161 (3)	-0.0153 (3)
S9	0.0424 (4)	0.0286 (3)	0.0513 (4)	-0.0096 (3)	0.0080 (3)	-0.0127 (3)
S10	0.0481 (4)	0.0387 (4)	0.0655 (5)	-0.0078 (3)	0.0055 (3)	-0.0265 (3)
C1	0.0414 (13)	0.0279 (11)	0.0322 (12)	-0.0032 (10)	0.0038 (10)	-0.0049 (9)
C2	0.0395 (13)	0.0240 (11)	0.0327 (12)	-0.0041 (9)	-0.0026 (10)	-0.0041 (9)
C3	0.0615 (17)	0.0322 (13)	0.0366 (13)	-0.0067 (12)	0.0040 (12)	-0.0107 (10)
C4	0.0379 (13)	0.0244 (11)	0.0369 (12)	-0.0051 (9)	-0.0010 (10)	-0.0045 (9)
C5	0.0416 (13)	0.0273 (11)	0.0365 (13)	-0.0036 (10)	0.0017 (10)	-0.0085 (10)
C6	0.0377 (13)	0.0286 (12)	0.0406 (13)	-0.0012 (10)	-0.0040 (10)	-0.0084 (10)
C7	0.0596 (17)	0.0500 (16)	0.0416 (15)	-0.0162 (14)	-0.0084 (13)	-0.0006 (12)
C8	0.0566 (17)	0.0423 (15)	0.0439 (15)	-0.0076 (13)	0.0006 (13)	-0.0019 (12)
C9	0.0634 (18)	0.0460 (15)	0.0425 (15)	-0.0130 (13)	0.0014 (13)	-0.0142 (12)
C10	0.0571 (18)	0.0591 (18)	0.0574 (18)	-0.0101 (14)	-0.0133 (14)	-0.0115 (15)
C11	0.0505 (16)	0.0456 (15)	0.0566 (17)	-0.0074 (13)	0.0037 (13)	-0.0099 (13)
C12	0.0595 (17)	0.0406 (14)	0.0381 (14)	-0.0205 (13)	0.0094 (12)	-0.0071 (11)
C13	0.083 (2)	0.0518 (16)	0.0410 (15)	-0.0342 (16)	0.0201 (14)	-0.0110 (12)
C14	0.0390 (14)	0.0527 (16)	0.0428 (15)	-0.0092 (12)	-0.0013 (11)	-0.0024 (12)

C15	0.0506 (16)	0.0469 (15)	0.0432 (15)	-0.0080 (12)	0.0037 (12)	0.0039 (12)
C16	0.0549 (17)	0.0564 (17)	0.0527 (17)	-0.0257 (14)	0.0091 (14)	-0.0125 (14)
C17	0.0458 (16)	0.069 (2)	0.0609 (19)	-0.0165 (15)	-0.0098 (14)	-0.0116 (16)
C18	0.0634 (19)	0.0434 (15)	0.0472 (16)	-0.0029 (13)	-0.0158 (14)	0.0007 (12)
Cl1	0.1180 (8)	0.0762 (6)	0.0815 (6)	-0.0117 (5)	-0.0170 (6)	-0.0447 (5)
Cl2	0.0858 (6)	0.0596 (5)	0.0972 (7)	0.0171 (5)	-0.0164 (5)	-0.0142 (5)
N1	0.0488 (12)	0.0408 (12)	0.0321 (11)	-0.0139 (10)	0.0058 (9)	-0.0106 (9)

Geometric parameters (\AA , $^{\circ}$)

Ni1—S2	2.1518 (7)	C8—C9	1.380 (4)
Ni1—S7	2.1643 (7)	C8—Cl2	1.722 (3)
Ni1—S6	2.1681 (7)	C9—C10	1.369 (4)
Ni1—S1	2.1714 (7)	C9—Cl1	1.731 (3)
S1—C1	1.719 (2)	C10—C11	1.378 (4)
S2—C2	1.708 (2)	C10—H10	0.9300
S3—C3	1.725 (3)	C11—C12	1.370 (4)
S3—C2	1.739 (2)	C11—H11	0.9300
S4—C3	1.738 (3)	C12—C13	1.515 (4)
S4—C1	1.750 (2)	C13—N1	1.493 (3)
S5—C3	1.642 (2)	C13—H13A	0.9700
S6—C4	1.717 (2)	C13—H13B	0.9700
S7—C5	1.721 (2)	C14—N1	1.336 (3)
S8—C6	1.727 (2)	C14—C15	1.369 (4)
S8—C5	1.742 (2)	C14—H14	0.9300
S9—C6	1.717 (3)	C15—C16	1.365 (4)
S9—C4	1.742 (2)	C15—H15	0.9300
S10—C6	1.662 (2)	C16—C17	1.368 (4)
C1—C2	1.356 (3)	C16—H16	0.9300
C4—C5	1.353 (3)	C17—C18	1.353 (4)
C7—C12	1.374 (4)	C17—H17	0.9300
C7—C8	1.383 (4)	C18—N1	1.340 (3)
C7—H7	0.9300	C18—H18	0.9300
S2—Ni1—S7	85.25 (3)	C7—C8—Cl2	119.6 (2)
S2—Ni1—S6	179.02 (3)	C10—C9—C8	120.1 (3)
S7—Ni1—S6	93.77 (2)	C10—C9—Cl1	119.0 (2)
S2—Ni1—S1	93.18 (3)	C8—C9—Cl1	120.9 (2)
S7—Ni1—S1	178.41 (3)	C9—C10—C11	120.0 (3)
S6—Ni1—S1	87.80 (3)	C9—C10—H10	120.0
C1—S1—Ni1	101.80 (8)	C11—C10—H10	120.0
C2—S2—Ni1	102.21 (8)	C12—C11—C10	120.3 (3)
C3—S3—C2	97.20 (12)	C12—C11—H11	119.8
C3—S4—C1	97.08 (12)	C10—C11—H11	119.8
C4—S6—Ni1	101.19 (8)	C11—C12—C7	119.8 (2)
C5—S7—Ni1	101.47 (9)	C11—C12—C13	119.1 (3)
C6—S8—C5	97.19 (11)	C7—C12—C13	121.0 (3)
C6—S9—C4	97.89 (11)	N1—C13—C12	112.5 (2)

C2—C1—S1	120.90 (18)	N1—C13—H13A	109.1
C2—C1—S4	115.57 (18)	C12—C13—H13A	109.1
S1—C1—S4	123.50 (14)	N1—C13—H13B	109.1
C1—C2—S2	121.87 (17)	C12—C13—H13B	109.1
C1—C2—S3	116.76 (18)	H13A—C13—H13B	107.8
S2—C2—S3	121.35 (14)	N1—C14—C15	120.3 (2)
S5—C3—S3	121.91 (17)	N1—C14—H14	119.9
S5—C3—S4	124.71 (17)	C15—C14—H14	119.9
S3—C3—S4	113.37 (13)	C16—C15—C14	119.7 (3)
C5—C4—S6	122.13 (17)	C16—C15—H15	120.1
C5—C4—S9	115.30 (17)	C14—C15—H15	120.1
S6—C4—S9	122.56 (14)	C15—C16—C17	118.9 (3)
C4—C5—S7	121.35 (18)	C15—C16—H16	120.6
C4—C5—S8	116.38 (17)	C17—C16—H16	120.6
S7—C5—S8	122.25 (15)	C18—C17—C16	120.1 (3)
S10—C6—S9	122.86 (15)	C18—C17—H17	119.9
S10—C6—S8	124.01 (15)	C16—C17—H17	119.9
S9—C6—S8	113.12 (13)	N1—C18—C17	120.5 (3)
C12—C7—C8	120.3 (3)	N1—C18—H18	119.7
C12—C7—H7	119.9	C17—C18—H18	119.7
C8—C7—H7	119.9	C14—N1—C18	120.5 (2)
C9—C8—C7	119.5 (3)	C14—N1—C13	119.1 (2)
C9—C8—Cl2	120.9 (2)	C18—N1—C13	120.4 (2)
S2—Ni1—S1—C1	1.40 (9)	Ni1—S7—C5—S8	-175.28 (13)
S6—Ni1—S1—C1	-178.64 (8)	C6—S8—C5—C4	1.5 (2)
S7—Ni1—S2—C2	177.99 (8)	C6—S8—C5—S7	-179.86 (15)
S1—Ni1—S2—C2	-1.80 (8)	C4—S9—C6—S10	178.91 (15)
S7—Ni1—S6—C4	1.33 (8)	C4—S9—C6—S8	-2.37 (15)
S1—Ni1—S6—C4	-178.88 (8)	C5—S8—C6—S10	179.61 (16)
S2—Ni1—S7—C5	177.64 (9)	C5—S8—C6—S9	0.91 (15)
S6—Ni1—S7—C5	-2.31 (9)	C12—C7—C8—C9	0.4 (4)
Ni1—S1—C1—C2	-0.6 (2)	C12—C7—C8—Cl2	-180.0 (2)
Ni1—S1—C1—S4	-178.50 (13)	C7—C8—C9—C10	-1.4 (4)
C3—S4—C1—C2	0.5 (2)	Cl2—C8—C9—C10	178.9 (2)
C3—S4—C1—S1	178.56 (16)	C7—C8—C9—Cl1	177.3 (2)
S1—C1—C2—S2	-1.1 (3)	Cl2—C8—C9—Cl1	-2.3 (4)
S4—C1—C2—S2	177.06 (12)	C8—C9—C10—C11	1.7 (5)
S1—C1—C2—S3	-179.55 (12)	Cl1—C9—C10—C11	-177.0 (2)
S4—C1—C2—S3	-1.4 (3)	C9—C10—C11—C12	-1.1 (5)
Ni1—S2—C2—C1	2.0 (2)	C10—C11—C12—C7	0.0 (4)
Ni1—S2—C2—S3	-179.53 (11)	C10—C11—C12—C13	177.6 (3)
C3—S3—C2—C1	1.6 (2)	C8—C7—C12—C11	0.3 (4)
C3—S3—C2—S2	-176.89 (15)	C8—C7—C12—C13	-177.2 (2)
C2—S3—C3—S5	177.82 (17)	C11—C12—C13—N1	95.9 (3)
C2—S3—C3—S4	-1.20 (17)	C7—C12—C13—N1	-86.5 (3)
C1—S4—C3—S5	-178.40 (18)	N1—C14—C15—C16	-1.2 (4)
C1—S4—C3—S3	0.59 (17)	C14—C15—C16—C17	0.0 (4)

Ni1—S6—C4—C5	0.4 (2)	C15—C16—C17—C18	1.0 (5)
Ni1—S6—C4—S9	179.73 (12)	C16—C17—C18—N1	-0.9 (5)
C6—S9—C4—C5	3.5 (2)	C15—C14—N1—C18	1.4 (4)
C6—S9—C4—S6	-175.90 (15)	C15—C14—N1—C13	-176.2 (2)
S6—C4—C5—S7	-2.6 (3)	C17—C18—N1—C14	-0.4 (4)
S9—C4—C5—S7	177.97 (12)	C17—C18—N1—C13	177.2 (3)
S6—C4—C5—S8	176.01 (13)	C12—C13—N1—C14	-78.8 (3)
S9—C4—C5—S8	-3.4 (3)	C12—C13—N1—C18	103.7 (3)
Ni1—S7—C5—C4	3.3 (2)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C14—H14···S10 ⁱ	0.93	2.82	3.622 (3)	145
C18—H18···S1 ⁱⁱ	0.93	2.79	3.708 (3)	168

Symmetry codes: (i) $x+1, y, z$; (ii) $x, y+1, z$.