

**μ_3 -Dodecatungsto(V,VI)aluminato-
 $\kappa^3O:O':O''$ -tris[aquabis(ethylene-
diamine- κ^2N,N')copper(II)]**

Yu-Kun Lu,^{a*} Yuan-Yuan Qu,^b Ming-Ming Tian,^a
Cheng-Lin Diao^b and Yun-Qi Liu^b

^aState Key Laboratory of Heavy Oil Processing, College of Science, China University of Petroleum (East China), Qingdao Shandong 266555, People's Republic of China, and ^bState Key Laboratory of Heavy Oil Processing, College of Chemical Engineering, China University of Petroleum (East China), Qingdao Shandong 266555, People's Republic of China

Correspondence e-mail: lyk@upc.edu.cn


Received 10 October 2011; accepted 14 November 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.019$ Å; disorder in main residue; R factor = 0.035; wR factor = 0.071; data-to-parameter ratio = 14.1.

The title compound, $[AlCu_3W_{12}O_{40}(C_2H_8N_2)_6(H_2O)_3]$, was prepared under hydrothermal conditions. The Cu^{2+} ion displays an elongated octahedral geometry defined by one bridging O atom from the polyoxidoanion and a coordinated water molecule in axial positions and four N atoms of the two chelating ethylenediamine (en) ligands in equatorial positions. The one-electron reduced $[AlW_{12}O_{40}]^{6-}$ anion coordinates three $[Cu(en)(H_2O)]^{2+}$ fragments, generating a neutral tri-supported Keggin-type polyoxidometalate (POM). This tri-supported POM is located in a special position of $\bar{3}$ symmetry and therefore O atoms from the central AlO_4 tetrahedron are disordered over two sets of sites. Disorder is also observed for three other bridging O atoms of the POM. In the crystal, molecules are connected via $N-H\cdots O$ and $O-H\cdots O$ hydrogen bonds, forming a three-dimensional framework.

Related literature

For the isotopic V^{IV} and Si^{IV} structures, see: Lu, Cui, Chen *et al.* (2009). For general background to polyoxidometalates, see: Pope & Müller (1991); Hill (1998); López *et al.* (2001). For modified Keggin-type structures with transition metal complexes, see: Xu *et al.* (2000); Yuan, Li *et al.* (2003). For the structure and chemistry of one-electron reduced heteropolytungstate, see: Lan *et al.* (2008); Meng *et al.* (2008). For other dodecatungstoaluminates, see: Wang *et al.* (2006); Yuan, Qin *et al.* (2009). For polyoxidometalates prepared with strongly reducing agents, see: Lu, Cui, Liu *et al.* (2009); Lu, Xu & Yu (2010); Lu, Xu, Cui *et al.* (2010).



Experimental

Crystal data

$[AlCu_3W_{12}O_{40}(C_2H_8N_2)_6(H_2O)_3]$	$Z = 6$
$M_r = 3478.47$	Mo $K\alpha$ radiation
Trigonal, $\bar{R}\bar{3}c$	$\mu = 26.38$ mm $^{-1}$
$a = 17.9719$ (14) Å	$T = 296$ K
$c = 29.335$ (5) Å	$0.11 \times 0.11 \times 0.10$ mm
$V = 8206$ (2) Å 3	

Data collection

Rigaku R-Axis RAPID diffractometer	22760 measured reflections
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	2220 independent reflections
$T_{min} = 0.159$, $T_{max} = 0.178$	1864 reflections with $I > 2\sigma(I)$
	$R_{int} = 0.071$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$	157 parameters
$wR(F^2) = 0.071$	H-atom parameters constrained
$S = 1.10$	$\Delta\rho_{max} = 1.85$ e Å $^{-3}$
2220 reflections	$\Delta\rho_{min} = -3.56$ e Å $^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

D-H···A	D-H	H···A	D···A	D-H···A
O1W—H1W···O2 ⁱ	0.85	2.25	2.856 (9)	128
N1—H1B···O5 ⁱ	0.90	2.26	3.138 (17)	163
N1—H1B···O5' ⁱ	0.90	2.30	3.185 (17)	170
N2—H2A···O7 ⁱⁱ	0.90	2.35	3.101 (17)	141
N2—H2B···O1 ⁱⁱⁱ	0.90	2.11	2.956 (12)	157

Symmetry codes: (i) $x + \frac{1}{3}, x - y + \frac{5}{3}, z + \frac{1}{6}$; (ii) $-x + \frac{2}{3}, -y + \frac{2}{3}, -z + \frac{1}{3}$; (iii) $x - y + 1, -y + 2, -z + \frac{1}{2}$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Natural Science Foundation of Shandong Province (ZR2011BQ004) and the Fundamental Research Funds for the Central Universities (09CX04045A).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2416).

References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Hill, C. L. (1998). *Chem. Rev.* **98**, 1–2.
- Lan, Y. Q., Li, S. L., Li, Y. G., Su, Z. M., Shao, K. Z. & Wang, X. L. (2008). *CrystEngComm*, **10**, 1129–1131.
- López, X., Maestre, J. M., Bo, C. & Poblet, J. M. (2001). *J. Am. Chem. Soc.* **123**, 9571–9576.
- Lu, Y. K., Cui, X. B., Chen, Y., Xu, J. N., Zhang, Q. B., Liu, Y. B., Xu, J. Q. & Wang, T. G. (2009). *J. Solid State Chem.* **182**, 2111–2117.
- Lu, Y. K., Cui, X. B., Liu, Y. B., Yang, Q. F., Shi, S. Y., Xu, J. Q. & Wang, T. G. (2009). *J. Solid State Chem.* **182**, 690–697.
- Lu, Y. K., Xu, J. N., Cui, X. B., Jin, J., Shi, S. Y. & Xu, J. Q. (2010). *Inorg. Chem. Commun.* **13**, 46–49.
- Lu, Y., Xu, J. & Yu, H. (2010). *Acta Cryst. E66*, m263–m264.
- Meng, F. X., Chen, Y. G., Pang, H. J., Shi, D. M. & Sun, Y. (2008). *J. Coord. Chem.* **61**, 1513–1524.
- Pope, M. T. & Müller, A. (1991). *Angew. Chem. Int. Ed. Engl.* **30**, 34–48.
- Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Wang, J. P., Shen, Y. & Niu, J. Y. (2006). *J. Coord. Chem.* **59**, 1007–1014.
- Xu, Y., Xu, J. Q., Zhang, K. L., Zhang, Y. & You, X. Z. (2000). *Chem. Commun.* pp. 153–154.
- Yuan, M., Li, Y. G., Wang, E. B., Tian, C. G., Wang, L., Hu, C. W., Hu, N. H. & Jia, H. Q. (2003). *Inorg. Chem.* **42**, 3670–3676.
- Yuan, L., Qin, C., Wang, X. L., Li, Y. G. & Wang, E. B. (2009). *Dalton Trans.* pp. 4169–4175.

supporting information

Acta Cryst. (2011). E67, m1778–m1779 [https://doi.org/10.1107/S1600536811048288]

μ_3 -Dodecatungsto(V,VI)aluminato- $\kappa^3O\cdot O'\cdot O''$ -tris[aquabis(ethylenediamine- κ^2N,N')copper(II)]

Yu-Kun Lu, Yuan-Yuan Qu, Ming-Ming Tian, Cheng-Lin Diao and Yun-Qi Liu

S1. Comment

There has been extensive interest in polyoxometalates (POMs), owing to their fascinating properties and great potential applications in many fields including catalysis, material science, medicine, and magnetochemistry, and their unusual structural diversities (Pope & Müller, 1991; Hill, 1998; López *et al.*, 2001). Especially, the modified POMs, which are the decoration of polyoxoanions with various transition metal ions, organic ligands, and/or their complex moieties, can be regarded as an ideal atomic level structural model for the determination of the mechanisms of oxide-supported catalysts (Xu *et al.*, 2000; Yuan, Li *et al.*, 2003). Therefore, we focussed our research on the preparation of modified POMs with strong reducing reagents (Lu, Cui, Liu *et al.*, 2009; Lu, Xu, Cui *et al.*, 2010, Lu, Xu, Yu *et al.*, 2010).

As shown in Fig. 1, the title compound shows a neutral tri-supported classical psedo-Keggin type structure where three $[Cu(en)_2(H_2O)]^{2+}$ fragments are decorating the one-electron reduced heteropolyanion $[AlW_{11}^{VI}W^V O_4]^{6-}$, which is isotropic with its V^{IV} and Si^{IV} analogue (Lu, Cui, Chen *et al.*, 2009). The tri-supported POM is located in a special position of $\bar{3}$ symmetry and therefore oxygen atoms from the central AlO_4 tetrahedron are disordered over two sites. The pseudo-Keggin unit $[AlW_{12}O_{40}]^{6-}$ may be viewed as a shell of $\{W_{12}O_{36}\}$ encapsulating a disordered $\{AlO_4\}$ moiety, present at its center and responsible for the local tetrahedral geometry (Wang *et al.*, 2006; Yuan, Qin *et al.*, 2009). The central Al atom is surrounded by a cube of eight oxygen (six O8 and two O9) atoms with each of them having half-occupancy due to the inversion symmetry at Al1, and the oxygens of the $\{AlO_4\}$ group are covalently bonded to three different tungsten centers of the shell. All W atoms possess similar distorted octahedral geometry WO_6 defined by one terminal oxygen atom, four doubly bridging oxo-groups and one central oxygen atom. Three doubly-bridging oxo-groups are disordered over two sets of sites each (O5, O5', O6, O6', O7 and O7') with the occupancy factor assigned as 0.5. Al—O8 and Al—O9 bond lengths are 1.714 (11) and 1.81 (2), respectively, with mean bond distance 1.74 Å, in good agreement with the literature (López *et al.*, 2001). The three classes of W—O average distances (being 1.688, 1.939 and 2.297 Å, respectively) are comparable to the corresponding distances in the similar structures (Wang *et al.*, 2006; Yuan, Qin *et al.*, 2009). The heteropolyanion $[AlW_{11}^{VI}W^V O_4]^{6-}$ is a one-electron-reduced derivative of $[AlW_{12}O_{40}]^{6-}$, similar to other reported representatives (Lan *et al.*, 2008; Meng *et al.*, 2008). We consider that oxalic acid acts as reducing agent reducing W^{VI} to W^V in the reactions.

The most unusual structural feature of the title compound is that each of three surface bridging oxygen atoms (O4) of the polyoxoanion is coordinated to one $[Cu(en)_2(H_2O)]^{2+}$ fragment. The Cu1 center possesses an elongated octahedral geometry defined by the bridging oxygen atom (Cu—O4, 2.718 (9) Å) from the polyoxoanion, a coordination water molecule [Cu—O1W, 2.411 (11) Å] *trans* to O4 atom and four N atoms from two chelating en ligands with equal Cu—N bond lengths 2.002 (9) Å. The bond lengths and angles at Cu1 are consistent with the Jahn–Teller active d^9 electronic configuration of divalent copper. The tri-supported POMs are extended into three-dimensional supramolecular network

via a combination of intermolecular N—H···O and O—H···O hydrogen bonding (Fig. 2).

S2. Experimental

A mixture of $\text{Na}_2\text{WO}_4 \cdot 2\text{H}_2\text{O}$ (0.658 g, 2 mmol), $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (0.25 g, 1 mmol), $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ (0.189 g, 1.5 mmol), NaAlO_2 (0.10 g, 1.25 mmol) and H_2O (15 mL) was mixed and stirred for 30 min, and the pH was adjusted to 7 with en. The resulting suspension was transferred to a Teflon-lined autoclave (25 ml) and kept at 180°C for 3 days. After slow cooling to room temperature for 2 days, blue prism crystals were obtained by filtering, washing with distilled water, and drying in desiccators at ambient temperature. The yields were *ca* 42% based on W. Elemental analysis $\text{C}_{12}\text{H}_{54}\text{Cu}_3\text{N}_{12}\text{O}_{43}\text{AlW}_{12}(3478.47)$: Calcd. (%): C, 4.14; H, 1.56; N, 4.83. Found: C, 4.21; H, 1.57; N, 4.94.

S3. Refinement

H atoms bonded to C and N atoms were positioned geometrically and refined as riding atoms, with C—H = 0.97, N—H = 0.90 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$. H atoms attached to the water molecule were located in a difference Fourier map and refined as riding, with O—H = 0.85 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. In the final difference Fourier map, the highest peak and the deepest hole are 0.37 Å and 0.93 Å from atom W2, respectively.





Figure 1


The molecular structure of the title compound with displacement ellipsoids drawn at the 30% probability level and labels shown for the asymmetric unit; the remaining part of the molecule was generated by the symmetry operations: $-y, x-y, z; -x+y, -x, z; x-y, -y, 0.5-z; -x, -x+y, 0.5-z; y, x, 0.5-z$. Only one position of the $\{\text{AlO}_4\}$ unit and one position of the disordered O5, O6 and O7 atoms is shown. H atoms have been omitted for clarity.

**Figure 2**

The crystal packing of the title compound viewed along the [001] direction with N—H···O and O—H···O hydrogen bonds displayed as dashed lines.

$\mu_3\text{-Dodecatungsto(V,VI)aluminato-}\kappa^3\text{O:O':O''-tris[aquabis(ethylenediamine-}\kappa^2\text{N,N')copper(II)]}$

Crystal data



$M_r = 3478.47$

Trigonal, $R\bar{3}c$

Hall symbol: -R 3 2 "c

$a = 17.9719 (14)$ Å

$c = 29.335 (5)$ Å

$V = 8206 (2)$ Å³

$Z = 6$

$F(000) = 9252$

$D_x = 4.224$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4488 reflections

$\theta = 2.3\text{--}27.9^\circ$

$\mu = 26.38$ mm⁻¹

$T = 296$ K

Prism, blue

$0.11 \times 0.11 \times 0.10$ mm

Data collection

Rigaku R-AXIS RAPID
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 10 pixels mm⁻¹
 ω scans

Absorption correction: multi-scan
(*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.159$, $T_{\max} = 0.178$

22760 measured reflections

2220 independent reflections

1864 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.071$

$\theta_{\max} = 28.0^\circ$, $\theta_{\min} = 2.3^\circ$

$h = -23 \rightarrow 23$

$k = -23 \rightarrow 23$

$l = -38 \rightarrow 38$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.035$$

$$wR(F^2) = 0.071$$

$$S = 1.10$$

2220 reflections

157 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0089P)^2 + 625.9202P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 1.85 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -3.56 \text{ e } \text{\AA}^{-3}$$

Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.000015 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
A11	0.0000	1.0000	0.2500	0.0086 (12)	
W1	0.12019 (2)	0.89540 (2)	0.251326 (13)	0.01398 (11)	
W2	0.11116 (3)	0.99748 (3)	0.151770 (19)	0.03112 (15)	
Cu1	0.38149 (11)	1.0000	0.2500	0.0250 (4)	
O1	0.1553 (4)	0.8230 (4)	0.2512 (3)	0.035 (2)	
O2	0.1623 (6)	0.9928 (5)	0.1054 (3)	0.035 (2)	
O3	0.0000	0.8191 (5)	0.2500	0.032 (3)	
O4	0.2303 (5)	1.0000	0.2500	0.021 (2)	
O5	0.1345 (9)	0.9100 (9)	0.1856 (6)	0.017 (3)	0.50
O5'	0.0992 (10)	0.9148 (9)	0.1888 (5)	0.014 (3)	0.50
O6	0.1377 (9)	0.9148 (8)	0.3158 (5)	0.016 (3)	0.50
O6'	0.0980 (9)	0.9157 (9)	0.3118 (5)	0.013 (3)	0.50
O7	0.0887 (8)	1.0866 (8)	0.1188 (5)	0.016 (3)*	0.50
O7'	-0.0036 (9)	0.9157 (8)	0.1502 (5)	0.018 (3)	0.50
O8	0.0000 (8)	1.0914 (7)	0.2333 (4)	0.010 (2)	0.50
O9	0.0000	1.0000	0.1883 (7)	0.011 (4)	0.50
O1W	0.5157 (6)	1.0000	0.2500	0.039 (3)	
H1W	0.5318	0.9756	0.2693	0.059*	
C1	0.3609 (8)	1.0205 (8)	0.3444 (4)	0.036 (3)	
H1C	0.3626	1.0033	0.3755	0.043*	
H1D	0.3169	1.0365	0.3420	0.043*	
C2	0.4485 (8)	1.0956 (8)	0.3305 (4)	0.037 (3)	
H2C	0.4615	1.1460	0.3484	0.044*	

H2D	0.4930	1.0811	0.3359	0.044*
N1	0.3427 (6)	0.9487 (6)	0.3121 (3)	0.028 (2)
H1A	0.2861	0.9102	0.3118	0.033*
H1B	0.3712	0.9218	0.3208	0.033*
N2	0.4457 (6)	1.1133 (6)	0.2820 (3)	0.031 (2)
H2A	0.4994	1.1439	0.2707	0.037*
H2B	0.4189	1.1438	0.2780	0.037*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Al1	0.0072 (17)	0.0072 (17)	0.011 (3)	0.0036 (8)	0.000	0.000
W1	0.01112 (18)	0.00943 (17)	0.0236 (2)	0.00683 (14)	-0.00119 (14)	-0.00112 (14)
W2	0.0239 (2)	0.0154 (2)	0.0504 (3)	0.00709 (17)	0.0232 (2)	-0.00150 (19)
Cu1	0.0300 (7)	0.0256 (9)	0.0178 (9)	0.0128 (5)	0.0001 (4)	0.0001 (7)
O1	0.012 (3)	0.013 (3)	0.080 (6)	0.007 (3)	-0.003 (4)	-0.001 (4)
O2	0.056 (5)	0.023 (4)	0.022 (4)	0.017 (4)	0.013 (4)	0.003 (3)
O3	0.008 (4)	0.009 (3)	0.078 (9)	0.004 (2)	0.002 (5)	0.001 (3)
O4	0.005 (3)	0.010 (4)	0.049 (7)	0.005 (2)	-0.001 (2)	-0.003 (4)
O5	0.010 (7)	0.007 (6)	0.034 (9)	0.004 (6)	0.006 (7)	-0.001 (6)
O5'	0.015 (8)	0.015 (7)	0.009 (7)	0.005 (6)	-0.006 (6)	-0.004 (5)
O6	0.011 (7)	0.007 (6)	0.029 (8)	0.005 (6)	0.003 (6)	0.001 (5)
O6'	0.013 (7)	0.016 (7)	0.014 (7)	0.009 (6)	0.000 (6)	0.003 (5)
O7'	0.023 (7)	0.009 (6)	0.025 (8)	0.009 (6)	-0.004 (6)	0.002 (6)
O8	0.011 (5)	0.015 (6)	0.009 (5)	0.009 (5)	0.001 (6)	-0.004 (5)
O9	0.011 (6)	0.011 (6)	0.012 (10)	0.005 (3)	0.000	0.000
O1W	0.026 (4)	0.038 (7)	0.058 (8)	0.019 (3)	0.009 (3)	0.017 (6)
C1	0.044 (7)	0.062 (8)	0.022 (6)	0.042 (7)	0.001 (5)	0.001 (6)
C2	0.039 (7)	0.049 (8)	0.028 (6)	0.026 (6)	-0.014 (5)	-0.012 (6)
N1	0.025 (5)	0.040 (5)	0.024 (5)	0.020 (4)	0.002 (4)	0.002 (4)
N2	0.030 (5)	0.032 (5)	0.035 (6)	0.020 (4)	-0.008 (4)	-0.004 (4)

Geometric parameters (\AA , $^\circ$)

Al1—O8 ⁱ	1.714 (11)	W2—O7 ⁱⁱⁱ	2.079 (13)
Al1—O9	1.81 (2)	W2—O9	2.288 (10)
W1—O1	1.707 (7)	W2—O8 ⁱⁱⁱ	2.421 (11)
W1—O3	1.894 (6)	Cu1—N2	2.002 (9)
W1—O6'	1.894 (6)	Cu1—N1	2.002 (9)
W1—O6	1.921 (16)	Cu1—O1W	2.411 (11)
W1—O4	1.931 (4)	Cu1—O4	2.718 (9)
W1—O5'	1.939 (14)	O1W—H1W	0.8499
W1—O5	1.946 (17)	C1—N1	1.498 (15)
W1—O8 ⁱⁱ	2.232 (11)	C1—C2	1.530 (17)
W1—O8 ⁱⁱⁱ	2.248 (12)	C1—H1C	0.9700
W2—O2	1.668 (8)	C1—H1D	0.9700
W2—O5'	1.765 (14)	C2—N2	1.465 (15)
W2—O7 ^{iv}	1.787 (13)	C2—H2C	0.9700

W2—O6 ^v	1.793 (14)	C2—H2D	0.9700
W2—O7'	1.840 (13)	N1—H1A	0.9000
W2—O6 ^v	2.063 (14)	N1—H1B	0.9000
W2—O5	2.072 (15)	N2—H2A	0.9000
W2—O7	2.076 (13)	N2—H2B	0.9000
O8 ⁱⁱ —Al1—O8 ⁱ	112.2 (3)	O6 ^v —W2—O8 ⁱⁱⁱ	56.2 (5)
O8 ⁱⁱ —Al1—O8 ^{iv}	122.8 (8)	O7'—W2—O8 ⁱⁱⁱ	85.6 (5)
O8 ⁱⁱ —Al1—O9 ⁱⁱ	73.4 (4)	O6 ^v —W2—O8 ⁱⁱⁱ	68.1 (5)
O9 ⁱⁱ —Al1—O9	180.000 (2)	O5—W2—O8 ⁱⁱⁱ	68.2 (5)
O1—W1—O3	99.8 (4)	O7—W2—O8 ⁱⁱⁱ	111.0 (5)
O1—W1—O6'	110.1 (5)	O7 ⁱⁱⁱ —W2—O8 ⁱⁱⁱ	112.0 (5)
O3—W1—O6'	83.4 (5)	O9—W2—O8 ⁱⁱⁱ	53.0 (5)
O1—W1—O6	93.0 (5)	N2—Cu1—N2 ^v	172.2 (5)
O3—W1—O6	100.1 (4)	N2—Cu1—N1	86.2 (4)
O6'—W1—O6	22.1 (4)	N2 ^v —Cu1—N1	94.7 (4)
O1—W1—O4	98.8 (3)	N2—Cu1—N1 ^v	94.7 (4)
O3—W1—O4	161.2 (3)	N2 ^v —Cu1—N1 ^v	86.2 (4)
O6'—W1—O4	92.5 (4)	N1—Cu1—N1 ^v	166.4 (5)
O6—W1—O4	81.2 (4)	N2—Cu1—O1W	86.1 (3)
O1—W1—O5'	108.3 (6)	N2 ^v —Cu1—O1W	86.1 (3)
O3—W1—O5'	81.8 (5)	N1—Cu1—O1W	96.8 (3)
O6'—W1—O5'	140.6 (6)	N1 ^v —Cu1—O1W	96.8 (3)
O6—W1—O5'	158.1 (6)	N2—Cu1—O4	93.9 (3)
O4—W1—O5'	90.2 (4)	N2 ^v —Cu1—O4	93.9 (3)
O1—W1—O5	91.4 (5)	N1—Cu1—O4	83.2 (3)
O3—W1—O5	95.9 (4)	N1 ^v —Cu1—O4	83.2 (3)
O6'—W1—O5	158.3 (6)	O1W—Cu1—O4	180.000 (2)
O6—W1—O5	162.3 (6)	W1—O3—W1 ⁱ	162.3 (6)
O4—W1—O5	81.2 (4)	W1—O4—W1 ^v	114.9 (4)
O5'—W1—O5	20.4 (4)	W1—O4—Cu1	122.5 (2)
O1—W1—O8 ⁱⁱ	166.6 (4)	W1 ^v —O4—Cu1	122.5 (2)
O3—W1—O8 ⁱⁱ	87.3 (4)	O5'—O5—W1	79 (2)
O6'—W1—O8 ⁱⁱ	59.2 (5)	O5'—O5—W2	54.6 (19)
O6—W1—O8 ⁱⁱ	74.6 (5)	W1—O5—W2	120.9 (7)
O4—W1—O8 ⁱⁱ	75.0 (4)	O5—O5'—W2	107 (2)
O5'—W1—O8 ⁱⁱ	83.8 (5)	O5—O5'—W1	80 (2)
O5—W1—O8 ⁱⁱ	99.1 (5)	W2—O5'—W1	141.3 (8)
O1—W1—O8 ⁱⁱⁱ	164.6 (4)	O6'—O6—W1	76.9 (19)
O3—W1—O8 ⁱⁱⁱ	86.8 (4)	O6'—O6—W2 ^v	58.5 (16)
O6'—W1—O8 ⁱⁱⁱ	84.3 (5)	W1—O6—W2 ^v	121.0 (7)
O6—W1—O8 ⁱⁱⁱ	99.5 (5)	O6—O6'—W2 ^v	101.1 (19)
O4—W1—O8 ⁱⁱⁱ	74.6 (4)	O6—O6'—W1	81.0 (19)
O5'—W1—O8 ⁱⁱⁱ	58.7 (5)	W2 ^v —O6'—W1	140.3 (8)
O5—W1—O8 ⁱⁱⁱ	74.0 (5)	O7 ^{iv} —O7—W2	59.1 (11)
O8 ⁱⁱ —W1—O8 ⁱⁱⁱ	25.3 (6)	O7 ^{iv} —O7—W2 ^{iv}	62.2 (11)
O2—W2—O5'	107.3 (6)	W2—O7—W2 ^{iv}	114.8 (7)
O2—W2—O7 ^{iv}	114.1 (5)	O7 ⁱⁱⁱ —O7'—W2 ⁱⁱⁱ	94.5 (13)

O5'—W2—O7 ^{iv}	138.6 (7)	O7 ⁱⁱⁱ —O7'—W2	91.4 (13)
O2—W2—O6 ^v	111.0 (5)	W2 ⁱⁱⁱ —O7'—W2	149.6 (7)
O5'—W2—O6 ^v	95.7 (6)	O8 ⁱ —O8—Al1	73.4 (4)
O7 ^{iv} —W2—O6 ^v	70.1 (6)	O8 ⁱ —O8—W1 ⁱⁱ	78.3 (10)
O2—W2—O7'	111.5 (5)	Al1—O8—W1 ⁱⁱ	124.6 (6)
O5'—W2—O7'	74.1 (6)	O8 ⁱ —O8—W1 ^{iv}	76.4 (10)
O7 ^{iv} —W2—O7'	90.2 (7)	Al1—O8—W1 ^{iv}	123.7 (6)
O6 ^v —W2—O7'	137.4 (6)	W1 ⁱⁱ —O8—W1 ^{iv}	93.3 (4)
O2—W2—O6 ^v	93.7 (5)	O8 ⁱ —O8—W2 ^{iv}	171.0 (3)
O5'—W2—O6 ^v	91.3 (6)	Al1—O8—W2 ^{iv}	115.6 (5)
O7 ^{iv} —W2—O6 ^v	86.5 (6)	W1 ⁱⁱ —O8—W2 ^{iv}	96.3 (4)
O6 ^v —W2—O6 ^v	20.4 (5)	W1 ^{iv} —O8—W2 ^{iv}	96.9 (4)
O7'—W2—O6 ^v	153.6 (6)	Al1—O9—W2 ⁱⁱⁱ	118.0 (5)
O2—W2—O5	91.5 (5)	Al1—O9—W2	118.0 (5)
O5'—W2—O5	18.6 (5)	W2 ⁱⁱⁱ —O9—W2	99.8 (6)
O7 ^{iv} —W2—O5	152.8 (6)	Al1—O9—W2 ^{iv}	118.0 (5)
O6 ^v —W2—O5	92.8 (6)	W2 ⁱⁱⁱ —O9—W2 ^{iv}	99.8 (6)
O7'—W2—O5	88.7 (6)	W2—O9—W2 ^{iv}	99.8 (6)
O6 ^v —W2—O5	82.5 (6)	Cu1—O1W—H1W	126.8
O2—W2—O7	89.1 (5)	N1—C1—C2	106.1 (9)
O5'—W2—O7	161.3 (7)	N1—C1—H1C	110.5
O6 ^v —W2—O7	86.3 (6)	C2—C1—H1C	110.5
O7'—W2—O7	91.8 (5)	N1—C1—H1D	110.5
O6 ^v —W2—O7	96.7 (5)	C2—C1—H1D	110.5
O5—W2—O7	179.0 (6)	H1C—C1—H1D	108.7
O2—W2—O7 ⁱⁱⁱ	86.7 (5)	N2—C2—C1	108.6 (9)
O5'—W2—O7 ⁱⁱⁱ	89.9 (6)	N2—C2—H2C	110.0
O7 ^{iv} —W2—O7 ⁱⁱⁱ	92.2 (5)	C1—C2—H2C	110.0
O6 ^v —W2—O7 ⁱⁱⁱ	158.7 (6)	N2—C2—H2D	110.0
O6 ^v —W2—O7 ⁱⁱⁱ	178.7 (5)	C1—C2—H2D	110.0
O5—W2—O7 ⁱⁱⁱ	98.7 (6)	H2C—C2—H2D	108.3
O7—W2—O7 ⁱⁱⁱ	82.1 (7)	C1—N1—Cu1	107.7 (7)
O2—W2—O9	153.2 (6)	C1—N1—H1A	110.2
O5'—W2—O9	89.3 (6)	Cu1—N1—H1A	110.2
O7 ^{iv} —W2—O9	52.5 (5)	C1—N1—H1B	110.2
O6 ^v —W2—O9	87.3 (6)	Cu1—N1—H1B	110.2
O7'—W2—O9	52.1 (5)	H1A—N1—H1B	108.5
O6 ^v —W2—O9	107.2 (6)	C2—N2—Cu1	107.4 (7)
O5—W2—O9	107.5 (5)	C2—N2—H2A	110.2
O7—W2—O9	72.2 (5)	Cu1—N2—H2A	110.2
O7 ⁱⁱⁱ —W2—O9	72.1 (5)	C2—N2—H2B	110.2
O2—W2—O8 ⁱⁱⁱ	153.7 (4)	Cu1—N2—H2B	110.2
O5'—W2—O8 ⁱⁱⁱ	56.7 (6)	H2A—N2—H2B	108.5
O7 ^{iv} —W2—O8 ⁱⁱⁱ	84.6 (6)		

Symmetry codes: (i) $-x, -x+y, -z+1/2$; (ii) $y-1, x+1, -z+1/2$; (iii) $-x+y-1, -x+1, z$; (iv) $-y+1, x-y+2, z$; (v) $x-y+1, -y+2, -z+1/2$.

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
O1W—H1W···O2 ^{vi}	0.85	2.25	2.856 (9)	128
N1—H1B···O5 ^{vi}	0.90	2.26	3.138 (17)	163
N1—H1B···O5 ^{vii}	0.90	2.30	3.185 (17)	170
N2—H2A···O7 ^{vii}	0.90	2.35	3.101 (17)	141
N2—H2B···O1 ^v	0.90	2.11	2.956 (12)	157

Symmetry codes: (v) $x-y+1, -y+2, -z+1/2$; (vi) $x+1/3, x-y+5/3, z+1/6$; (vii) $-x+2/3, -y+7/3, -z+1/3$.