

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Methyl 4-hydroxy-1,1-dioxo-2-(2-phenylethyl)-2H-1λ⁶,2-benzothiazine-3-carboxylate

 Muhammad Nadeem Arshad,^{a*} Islam Ullah Khan,^b
 Muhammad Zia-ur-Rehman,^c Muhammad Danish^a and
 K. Travis Holman^d

^aDepartment of Chemistry, University of Gujrat (H. H. Campus), Gujrat 50700, Pakistan, ^bMaterials Chemistry Laboratory, Department of Chemistry, GC University, Lahore 54000, Pakistan, ^cApplied Chemistry Research Centre PCSIR Laboratories Complex, Lahore 54600, Pakistan, and ^dDepartment of Chemistry, Georgetown University, 37th and O St NW, Washington, DC 20057, USA
 Correspondence e-mail: mnachemist@hotmail.com


Received 17 November 2011; accepted 21 November 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.049; wR factor = 0.116; data-to-parameter ratio = 17.7.

In the title compound, $\text{C}_{18}\text{H}_{17}\text{NO}_5\text{S}$, the thiazine ring adopts a half-chair conformation and the dihedral angle between the aromatic rings is $79.41(6)^\circ$. An intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond generates an $S(6)$ ring. In the crystal, molecules are linked by weak $\text{C}-\text{H}\cdots\text{O}$ interactions resulting in infinite sheets along the b and c axes.

Related literature

For related structures, see: Arshad *et al.* (2011a); Ahmad *et al.* (2010); Khalid *et al.* (2010). For further synthetic details, see: Arshad *et al.* (2011b). For graph-set notation, see: Bernstein, *et al.* (1995). For ring conformations, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{17}\text{NO}_5\text{S}$
 $M_r = 359.39$
 Monoclinic, $P2_1/c$
 $a = 9.018(2)$ Å
 $b = 19.026(4)$ Å

$c = 10.193(2)$ Å
 $\beta = 106.441(3)^\circ$
 $V = 1677.3(7)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.22$ mm⁻¹
 $T = 100$ K

$0.37 \times 0.23 \times 0.08$ mm

Data collection

Bruker SMART 1K diffractometer
 with a Bruker APEXII CCD
 detector
 Absorption correction: multi-scan
 (SADABS; Bruker, 2007)
 $T_{\min} = 0.922$, $T_{\max} = 0.982$

19773 measured reflections
 4066 independent reflections
 2735 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.094$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.116$
 $S = 1.01$
 4066 reflections
 230 parameters

H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\max} = 0.38$ e Å⁻³
 $\Delta\rho_{\min} = -0.44$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C2}-\text{H2}\cdots\text{O3}^i$	0.95	2.55	3.391 (3)	147
$\text{C17}-\text{H17}\cdots\text{O2}^{ii}$	0.95	2.53	3.205 (3)	128
$\text{C18}-\text{H18}\cdots\text{O4}^{iii}$	0.95	2.49	3.308 (3)	145
$\text{O1}-\text{H1}\cdots\text{O4}$	0.92 (3)	1.74 (3)	2.583 (2)	152 (3)

Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $x - 1, y, z - 1$; (iii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2009); software used to prepare material for publication: X-SEED (Barbour, 2001), WinGX (Farrugia, 1999) and PLATON.

MNA acknowledges the HEC for providing a fellowship under the International Research Support Initiative Program (IRSIP).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6520).

References

- Ahmad, M., Siddiqui, H. L., Zia-ur-Rehman, M., Elsegood, M. R. J. & Weaver, G. W. (2010). *Acta Cryst.* **E66**, o333.
 Arshad, M. N., Khan, I. U., Zia-ur-Rehman, M. & Shafiq, M. (2011b). *Asian J. Chem.* **23**, 2801–2805.
 Arshad, M. N., Khan, I. U., Zia-ur-Rehman, M., Shafiq, M. & Asiri, A. M. (2011a). *Acta Cryst.* **E67**, o1588–o1589.
 Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Bruker (2001). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Khalid, Z., Siddiqui, H. L., Ahmad, M., Aslam, S. & Parvez, M. (2010). *Acta Cryst.* **E66**, o885.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2011). E67, o3445 [https://doi.org/10.1107/S160053681104966X]

Methyl 4-hydroxy-1,1-dioxo-2-(2-phenylethyl)-2H-1 λ ⁶,2-benzothiazine-3-carboxylate

Muhammad Nadeem Arshad, Islam Ullah Khan, Muhammad Zia-ur-Rehman, Muhammad Danish and K. Travis Holman

S1. Comment

Benzothiazines molecules are well explored for their crystallographic studies (Arshad *et al.* 2011a), (Ahmad *et al.* 2010), (Khalid *et al.* 2010).

In the title compound, the nitrogen atom of methyl-4-hydroxy-2H-1, 2-benzothiazine-3-carboxylate 1,1-dioxide was alkylated with phenylethyl group. The thiazine ring (with root mean square deviation = 0.210Å) is oriented at dihedral angle of 13.12 (11)° with respect to the aromatic ring (C1—C6) and the S1 & N1 atoms showed deviations from the least square plane by 0.3121 (10)Å and -0.3310 (12)Å, respectively. The thiazine ring adopts a half chair conformation with puckering amplitude $Q = 0.5160$ (16)Å $\theta = 63.6$ (2)°, $\varphi = 26.3$ (2)° (Cremer & Pople, 1975). The phenyl ring (C13—C17) is oriented at dihedral angle of 79.41 (6)° and 66.53 (5)° with respect to the aromatic (C1—C6) and thiazine rings respectively. The intramolecular O—H \cdots O interaction observed and generates an almost planer S(6) ring (Bernstein, *et al.*, 1995) with the r.m.s deviation of 0.0131Å and produces dihedral angles of 16.02 (33)° & 15.87 (32)° with respect to the thiazine and aromatic (C1—C6) rings respectively.

Weak hydrogen bonding interactions of C—H \cdots O type connect the molecules. The interaction C2—H2 \cdots O3 resulted in the formation of dimers which are further linked along *b* and *c* axes (Fig.2. Tab.1).

S2. Experimental

The title compound was synthesised according to literature procedure (Arshad *et al.* 2011b) and crystalized in methanol under slow evaporation to yield colourless blocks.

S3. Refinement

All the C—H H-atoms were positioned with idealized geometry with C—H_{aromatic} = 0.95 Å, C—H_{methylene} = 0.99 Å and C—H_{methyl} = 0.98 Å and were refined using a riding model with $U_{iso}(H) = 1.2 U_{eq}$ for aromatic C atoms. The O—H H-atom was located via fourier map with O—H = 0.92 (3) Å with $U_{iso}(H) = 1.5 U_{eq}$ for O atom.




Figure 1

The labeled ortep diagram of (I) for which thermal ellipsoids were drawn at 50% probability level.




Figure 2

Unit cell packing for (I) showing the hydrogen bondings using dashed lines.

Methyl 4-hydroxy-1,1-dioxo-2-(2-phenylethyl)-2H-1 λ ^6,2-benzothiazine-3-carboxylate

Crystal data

$C_{18}H_{17}NO_5S$

$M_r = 359.39$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 9.018$ (2) Å

$b = 19.026$ (4) Å

$c = 10.193$ (2) Å

$\beta = 106.441$ (3)°

$V = 1677.3$ (7) Å³

$Z = 4$

$F(000) = 752$

$D_x = 1.423$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2258 reflections

$\theta = 2.3$ – 22.9 °

$\mu = 0.22$ mm⁻¹

$T = 100$ K

Block, colorless

$0.37 \times 0.23 \times 0.08$ mm

Data collection

Bruker SMART 1K

diffractometer with a Bruker APEXII CCD

detector

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2007)

$T_{\min} = 0.922$, $T_{\max} = 0.982$

19773 measured reflections

4066 independent reflections

2735 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.094$

$\theta_{\max} = 28.4^\circ$, $\theta_{\min} = 2.1^\circ$
 $h = -11 \rightarrow 11$

$k = -25 \rightarrow 25$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.116$
 $S = 1.01$
 4066 reflections
 230 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0449P)^2 + 0.2677P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.38 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.44 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.29319 (6)	0.07784 (3)	0.51969 (5)	0.01673 (14)
O3	0.30860 (16)	0.00885 (8)	0.46983 (15)	0.0214 (4)
O5	-0.10038 (17)	0.12689 (8)	0.61803 (15)	0.0231 (4)
O1	0.13070 (18)	0.28877 (8)	0.48807 (17)	0.0251 (4)
O4	-0.07394 (18)	0.24350 (9)	0.60086 (17)	0.0276 (4)
N1	0.10892 (18)	0.09548 (9)	0.48635 (17)	0.0154 (4)
O2	0.36535 (16)	0.09408 (8)	0.65995 (14)	0.0227 (4)
C1	0.3551 (2)	0.14133 (12)	0.4212 (2)	0.0170 (5)
C7	0.1578 (2)	0.22064 (12)	0.4703 (2)	0.0182 (5)
C8	0.0779 (2)	0.16750 (11)	0.5085 (2)	0.0177 (5)
C5	0.3377 (2)	0.26007 (12)	0.3379 (2)	0.0220 (5)
H5	0.2933	0.3057	0.3300	0.026*
C9	-0.0383 (2)	0.18329 (12)	0.5789 (2)	0.0207 (5)
C6	0.2849 (2)	0.20757 (11)	0.4087 (2)	0.0180 (5)
C2	0.4731 (2)	0.12640 (12)	0.3629 (2)	0.0214 (5)
H2	0.5193	0.0811	0.3720	0.026*
C17	-0.3703 (2)	0.07609 (13)	-0.0561 (2)	0.0235 (5)
H17	-0.4494	0.1074	-0.1035	0.028*
C18	-0.2463 (2)	0.10137 (12)	0.0471 (2)	0.0209 (5)
H18	-0.2404	0.1500	0.0694	0.025*
C11	-0.0022 (2)	0.05922 (11)	0.3704 (2)	0.0170 (5)
H11A	-0.1082	0.0670	0.3773	0.020*

H11B	0.0185	0.0081	0.3786	0.020*
C3	0.5220 (3)	0.17932 (13)	0.2911 (2)	0.0270 (6)
H3	0.6021	0.1701	0.2496	0.032*
C15	-0.2653 (3)	-0.03981 (13)	-0.0196 (2)	0.0247 (5)
H15	-0.2717	-0.0884	-0.0421	0.030*
C4	0.4549 (3)	0.24549 (13)	0.2793 (2)	0.0257 (5)
H4	0.4901	0.2813	0.2303	0.031*
C12	0.0052 (2)	0.08358 (12)	0.2302 (2)	0.0225 (5)
H12A	0.0047	0.1356	0.2273	0.027*
H12B	0.1029	0.0671	0.2146	0.027*
C14	-0.1420 (3)	-0.01493 (13)	0.0835 (2)	0.0249 (5)
H14	-0.0642	-0.0467	0.1314	0.030*
C13	-0.1300 (2)	0.05577 (12)	0.1181 (2)	0.0196 (5)
C16	-0.3799 (3)	0.00572 (13)	-0.0903 (2)	0.0239 (5)
H16	-0.4643	-0.0113	-0.1619	0.029*
C10	-0.2150 (3)	0.13972 (14)	0.6903 (3)	0.0314 (6)
H10A	-0.2992	0.1682	0.6329	0.047*
H10B	-0.1669	0.1649	0.7756	0.047*
H10C	-0.2564	0.0948	0.7111	0.047*
H1	0.057 (3)	0.2878 (14)	0.534 (3)	0.047*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0166 (2)	0.0155 (3)	0.0161 (3)	0.0009 (2)	0.0015 (2)	0.0007 (2)
O3	0.0204 (7)	0.0153 (8)	0.0269 (8)	0.0031 (6)	0.0042 (7)	-0.0005 (7)
O5	0.0245 (8)	0.0217 (9)	0.0271 (9)	-0.0026 (7)	0.0138 (7)	-0.0042 (7)
O1	0.0285 (9)	0.0147 (9)	0.0347 (10)	0.0009 (7)	0.0130 (8)	-0.0018 (7)
O4	0.0287 (9)	0.0193 (9)	0.0370 (10)	0.0027 (7)	0.0127 (8)	-0.0047 (7)
N1	0.0155 (8)	0.0138 (10)	0.0158 (9)	0.0005 (7)	0.0025 (7)	0.0009 (7)
O2	0.0237 (8)	0.0235 (9)	0.0171 (8)	-0.0012 (7)	-0.0005 (7)	0.0002 (7)
C1	0.0154 (9)	0.0186 (12)	0.0136 (10)	-0.0025 (9)	-0.0016 (8)	-0.0005 (9)
C7	0.0188 (10)	0.0161 (12)	0.0172 (11)	0.0004 (9)	0.0009 (9)	-0.0006 (9)
C8	0.0190 (10)	0.0174 (12)	0.0152 (10)	0.0025 (9)	0.0026 (9)	-0.0016 (9)
C5	0.0237 (11)	0.0182 (12)	0.0211 (12)	-0.0006 (9)	0.0018 (10)	0.0023 (9)
C9	0.0182 (10)	0.0210 (13)	0.0204 (11)	-0.0007 (9)	0.0015 (9)	-0.0017 (10)
C6	0.0175 (10)	0.0179 (12)	0.0158 (11)	-0.0035 (9)	0.0002 (9)	-0.0018 (9)
C2	0.0190 (10)	0.0219 (13)	0.0209 (11)	-0.0003 (9)	0.0020 (9)	-0.0029 (9)
C17	0.0217 (11)	0.0287 (14)	0.0186 (11)	0.0019 (10)	0.0035 (9)	0.0039 (10)
C18	0.0259 (11)	0.0193 (12)	0.0177 (11)	-0.0029 (10)	0.0067 (10)	-0.0004 (9)
C11	0.0162 (10)	0.0177 (12)	0.0160 (10)	-0.0016 (8)	0.0030 (9)	-0.0007 (8)
C3	0.0223 (11)	0.0353 (16)	0.0248 (12)	-0.0029 (10)	0.0089 (10)	-0.0019 (11)
C15	0.0301 (12)	0.0205 (13)	0.0243 (12)	0.0002 (10)	0.0089 (10)	-0.0030 (10)
C4	0.0256 (12)	0.0267 (14)	0.0245 (13)	-0.0067 (10)	0.0066 (10)	0.0046 (10)
C12	0.0227 (10)	0.0248 (13)	0.0182 (11)	-0.0036 (10)	0.0030 (9)	0.0031 (10)
C14	0.0243 (11)	0.0253 (14)	0.0234 (12)	0.0041 (10)	0.0040 (10)	0.0020 (10)
C13	0.0208 (11)	0.0247 (13)	0.0136 (10)	-0.0026 (9)	0.0053 (9)	0.0001 (9)
C16	0.0237 (11)	0.0287 (14)	0.0170 (11)	-0.0045 (10)	0.0017 (9)	-0.0036 (10)

C10	0.0303 (12)	0.0354 (16)	0.0360 (14)	-0.0049 (11)	0.0215 (12)	-0.0120 (12)
-----	-------------	-------------	-------------	--------------	-------------	--------------

Geometric parameters (Å, °)

S1—O2	1.4270 (15)	C17—C18	1.386 (3)
S1—O3	1.4285 (16)	C17—H17	0.9500
S1—N1	1.6339 (17)	C18—C13	1.395 (3)
S1—C1	1.760 (2)	C18—H18	0.9500
O5—C9	1.323 (3)	C11—C12	1.521 (3)
O5—C10	1.449 (3)	C11—H11A	0.9900
O1—C7	1.341 (3)	C11—H11B	0.9900
O1—H1	0.92 (3)	C3—C4	1.387 (3)
O4—C9	1.227 (3)	C3—H3	0.9500
N1—C8	1.429 (3)	C15—C14	1.379 (3)
N1—C11	1.485 (3)	C15—C16	1.384 (3)
C1—C2	1.387 (3)	C15—H15	0.9500
C1—C6	1.400 (3)	C4—H4	0.9500
C7—C8	1.361 (3)	C12—C13	1.511 (3)
C7—C6	1.475 (3)	C12—H12A	0.9900
C8—C9	1.459 (3)	C12—H12B	0.9900
C5—C4	1.382 (3)	C14—C13	1.387 (3)
C5—C6	1.392 (3)	C14—H14	0.9500
C5—H5	0.9500	C16—H16	0.9500
C2—C3	1.389 (3)	C10—H10A	0.9800
C2—H2	0.9500	C10—H10B	0.9800
C17—C16	1.380 (3)	C10—H10C	0.9800
O2—S1—O3	119.52 (9)	C13—C18—H18	119.9
O2—S1—N1	108.21 (9)	N1—C11—C12	114.02 (17)
O3—S1—N1	108.08 (9)	N1—C11—H11A	108.7
O2—S1—C1	107.13 (9)	C12—C11—H11A	108.7
O3—S1—C1	110.45 (10)	N1—C11—H11B	108.7
N1—S1—C1	102.04 (9)	C12—C11—H11B	108.7
C9—O5—C10	116.10 (18)	H11A—C11—H11B	107.6
C7—O1—H1	103.7 (17)	C4—C3—C2	120.5 (2)
C8—N1—C11	116.93 (16)	C4—C3—H3	119.7
C8—N1—S1	113.41 (13)	C2—C3—H3	119.7
C11—N1—S1	119.16 (14)	C14—C15—C16	120.3 (2)
C2—C1—C6	121.8 (2)	C14—C15—H15	119.9
C2—C1—S1	120.63 (17)	C16—C15—H15	119.9
C6—C1—S1	117.50 (16)	C5—C4—C3	120.9 (2)
O1—C7—C8	123.2 (2)	C5—C4—H4	119.6
O1—C7—C6	114.46 (19)	C3—C4—H4	119.6
C8—C7—C6	122.3 (2)	C13—C12—C11	111.30 (18)
C7—C8—N1	121.54 (19)	C13—C12—H12A	109.4
C7—C8—C9	120.0 (2)	C11—C12—H12A	109.4
N1—C8—C9	118.38 (19)	C13—C12—H12B	109.4
C4—C5—C6	119.7 (2)	C11—C12—H12B	109.4

C4—C5—H5	120.1	H12A—C12—H12B	108.0
C6—C5—H5	120.1	C15—C14—C13	121.0 (2)
O4—C9—O5	123.2 (2)	C15—C14—H14	119.5
O4—C9—C8	122.9 (2)	C13—C14—H14	119.5
O5—C9—C8	113.92 (19)	C14—C13—C18	118.5 (2)
C5—C6—C1	118.8 (2)	C14—C13—C12	121.3 (2)
C5—C6—C7	121.3 (2)	C18—C13—C12	120.1 (2)
C1—C6—C7	119.92 (19)	C17—C16—C15	119.4 (2)
C1—C2—C3	118.3 (2)	C17—C16—H16	120.3
C1—C2—H2	120.8	C15—C16—H16	120.3
C3—C2—H2	120.8	O5—C10—H10A	109.5
C16—C17—C18	120.5 (2)	O5—C10—H10B	109.5
C16—C17—H17	119.8	H10A—C10—H10B	109.5
C18—C17—H17	119.8	O5—C10—H10C	109.5
C17—C18—C13	120.3 (2)	H10A—C10—H10C	109.5
C17—C18—H18	119.9	H10B—C10—H10C	109.5
O2—S1—N1—C8	59.99 (16)	C4—C5—C6—C7	-178.7 (2)
O3—S1—N1—C8	-169.24 (14)	C2—C1—C6—C5	-1.3 (3)
C1—S1—N1—C8	-52.79 (16)	S1—C1—C6—C5	176.40 (16)
O2—S1—N1—C11	-156.23 (15)	C2—C1—C6—C7	178.87 (19)
O3—S1—N1—C11	-25.46 (18)	S1—C1—C6—C7	-3.5 (3)
C1—S1—N1—C11	90.99 (17)	O1—C7—C6—C5	-18.7 (3)
O2—S1—C1—C2	99.50 (18)	C8—C7—C6—C5	162.8 (2)
O3—S1—C1—C2	-32.19 (19)	O1—C7—C6—C1	161.14 (18)
N1—S1—C1—C2	-146.92 (17)	C8—C7—C6—C1	-17.4 (3)
O2—S1—C1—C6	-78.19 (17)	C6—C1—C2—C3	0.3 (3)
O3—S1—C1—C6	150.11 (15)	S1—C1—C2—C3	-177.31 (16)
N1—S1—C1—C6	35.39 (18)	C16—C17—C18—C13	-0.4 (3)
O1—C7—C8—N1	179.24 (18)	C8—N1—C11—C12	71.0 (2)
C6—C7—C8—N1	-2.4 (3)	S1—N1—C11—C12	-71.6 (2)
O1—C7—C8—C9	-3.3 (3)	C1—C2—C3—C4	0.5 (3)
C6—C7—C8—C9	175.06 (18)	C6—C5—C4—C3	-0.6 (3)
C11—N1—C8—C7	-103.1 (2)	C2—C3—C4—C5	-0.4 (3)
S1—N1—C8—C7	41.5 (2)	N1—C11—C12—C13	-168.72 (18)
C11—N1—C8—C9	79.4 (2)	C16—C15—C14—C13	-0.1 (3)
S1—N1—C8—C9	-136.00 (16)	C15—C14—C13—C18	0.5 (3)
C10—O5—C9—O4	-0.2 (3)	C15—C14—C13—C12	-179.6 (2)
C10—O5—C9—C8	179.15 (18)	C17—C18—C13—C14	-0.2 (3)
C7—C8—C9—O4	3.4 (3)	C17—C18—C13—C12	179.85 (19)
N1—C8—C9—O4	-179.04 (19)	C11—C12—C13—C14	-70.4 (3)
C7—C8—C9—O5	-175.93 (18)	C11—C12—C13—C18	109.5 (2)
N1—C8—C9—O5	1.6 (3)	C18—C17—C16—C15	0.8 (3)
C4—C5—C6—C1	1.4 (3)	C14—C15—C16—C17	-0.6 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C2—H2 \cdots O3 ⁱ	0.95	2.55	3.391 (3)	147
C17—H17 \cdots O2 ⁱⁱ	0.95	2.53	3.205 (3)	128
C18—H18 \cdots O4 ⁱⁱⁱ	0.95	2.49	3.308 (3)	145
O1—H1 \cdots O4	0.92 (3)	1.74 (3)	2.583 (2)	152 (3)

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $x-1, y, z-1$; (iii) $x, -y+1/2, z-1/2$.