

2-[4-(2-Hydroxyethoxy)phenyl]-4,4,5,5-tetramethyl-2-imidazoline-1-oxyl 3-oxide

Lin-Lin Jing, Hui-Ping Ma, Xiao-Fei Fan, Lei He and Zheng-Ping Jia*

Department of Pharmacy, Lanzhou General Hospital of PLA, Key Laboratory of the Prevention and Cure of Plateau Environmental Damage, PLA 730050, Lanzhou Gansu, People's Republic of China
Correspondence e-mail: zhengping_jia@yahoo.cn


Received 8 November 2011; accepted 13 November 2011

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.047; wR factor = 0.163; data-to-parameter ratio = 14.2.

In the title compound, $\text{C}_{15}\text{H}_{21}\text{N}_2\text{O}_4$, the imidazoline ring displays a twisted conformation. The dihedral angle between the mean plane of the imidazoline ring and the benzene ring is $33.50(12)^\circ$. In the crystal, molecules are connected by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a zigzag chain along the c axis. The chains are linked by $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For the preparation of the title compound, see: Ullman *et al.* (1974). For biological properties of nitronyl nitroxides, see: Soule *et al.* (2007); Blasig *et al.* (2002); Qin *et al.* (2009); Tanaka *et al.* (2007). For coordination properties of nitronyl nitroxides, see: Masuda *et al.* (2009). For related structures, see: Wang *et al.* (2009); Jing *et al.* (2009). For puckering parameters, see: Cremer & Pople (1975). For pseudorotation parameters, see: Rao *et al.* (1981).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{21}\text{N}_2\text{O}_4$

$M_r = 293.34$

Orthorhombic, $Pbca$

$a = 8.869(3)\text{ \AA}$

$b = 16.050(5)\text{ \AA}$

$c = 20.925(6)\text{ \AA}$

$V = 2978.7(16)\text{ \AA}^3$

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 0.10\text{ mm}^{-1}$

$T = 296\text{ K}$

$0.26 \times 0.23 \times 0.22\text{ mm}$

Data collection

Bruker APEXII CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2007)

$T_{\min} = 0.976$, $T_{\max} = 0.979$

20164 measured reflections
2774 independent reflections

1928 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$

$wR(F^2) = 0.163$

$S = 0.95$

2774 reflections

195 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.22\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.25\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg2$ is the centroid of the benzene C4–C9 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O4—H4 \cdots O2 ⁱ	0.82	2.01	2.828 (3)	173
C12—H12C \cdots O1 ⁱⁱ	0.96	2.54	3.418 (3)	152
C15—H15C \cdots Cg2 ⁱⁱⁱ	0.96	2.80	3.570 (3)	138

Symmetry codes: (i) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (ii) $x - \frac{1}{2}, y, -z + \frac{1}{2}$; (iii) $x + \frac{1}{2}, y, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009).

We thank the Natural Science Foundation of China (grant No. 30772773, 30472186) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5006).

References

- Blasig, I. E., Mertsch, K. & Haseloff, R. F. (2002). *Neuropharmacology*, **43**, 1006–1014.
- Bruker (2007). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Jing, L.-L., Wang, H.-B. & Sun, X.-L. (2009). *Acta Cryst. E65*, o2444.
- Masuda, Y., Kurats, M., Suzuki, S., Kozaki, M., Shiomi, D., Sato, K., Takui, T., Hosokoshi, Y., Miyazaki, Y., Inada, A. & Okada, K. (2009). *J. Am. Chem. Soc.* **131**, 4670–4673.
- Qin, X. Y., Ding, G. R. & Sun, X. L. (2009). *J. Chem. Res.* pp. 511–514.
- Rao, S. T., Westhof, E. & Sundaralingam, M. (1981). *Acta Cryst. A37*, 421–425.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Soule, B. P., Hyodo, F., Matsumoto, K., Simone, N. L., Cook, J. A., Krishna, M. C. & Mitchell, J. B. (2007). *Free Radic. Biol. Med.* **42**, 1632–1650.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Tanaka, K., Furuichi, K., Kozaki, M., Suzuki, S., Shiomi, D., Sato, K., Takui, T. & Okada, K. (2007). *Polyhedron*, **26**, 2021–2026.
- Ullman, E. F., Osiecki, J. H., Boocock, D. G. B. & Darcy, R. (1974). *J. Am. Chem. Soc.* **96**, 7049–7053.
- Wang, H.-B., Jing, L.-L., Gao, P. & Sun, X.-L. (2009). *Acta Cryst. E65*, o2090.

supporting information

Acta Cryst. (2011). E67, o3348 [https://doi.org/10.1107/S160053681104815X]

2-[4-(2-Hydroxyethoxy)phenyl]-4,4,5,5-tetramethyl-2-imidazoline-1-oxyl 3-oxide

Lin-Lin Jing, Hui-Ping Ma, Xiao-Fei Fan, Lei He and Zheng-Ping Jia

S1. Comment

Nitronyl nitroxides, firstly synthesized more than 30 years ago, can be used for coordination with many metal cations, such as Mn^{2+} , Cu^{2+} and Ni^{2+} leading to form some molecule-based magnetic materials (Masuda *et al.*, 2009). They can also react with free radicals such as OH, H_2O_2 and O_2 (Blasig *et al.*, 2002) to protect cells from the attack of free radicals. So they have a lot of biological properties as anticancer, antiradiation and antioxidation (Qin *et al.*, 2009; Tanaka *et al.*, 2007; Soule *et al.*, 2007).


The molecular structure of the title compound is shown in Fig. 1. The least-squares plane of the nitronyl nitroxide ring and the benzene ring are twisted with respect to each other making a dihedral angle of $33.50(12)^\circ$. The puckering parameters of the nitronyl nitroxide ring are $Q(2) = 0.177(2)$ Å and $\varphi = 237.1(7)^\circ$ (Cremer & Pople, 1975). The pseudorotation parameters (Rao *et al.*, 1981) for the nitronyl nitroxide ring are $P = 39.7(4)^\circ$ and $\tau(M) = 18.2(1)^\circ$ for the C1—N1 reference bond with the closest puckering descriptor being twisted on C1—C2. The crystal structure is stabilized by O—H \cdots O, C—H \cdots O and C—H \cdots π hydrogen bonds (Table 1).

S2. Experimental

2,3-Dimethyl-2,3-bis(hydroxylamino) butane (1.48 g, 10.0 mmol) and 4-(4-hydroxyethoxy)benzaldehyde (1.66 g, 10 mmol) were dissolved in methanol (30.0 ml). The reaction was filtered after stirring for 24 h at room temperature. The resulting white powder was washed by cool methanol and suspended in the solution of dichloromethane (30.0 ml). Then the reaction mixture was added to an aqueous solution of $NaIO_4$ (30 ml) and stirred for 15 min in an ice bath to give a dark blue solution. The aqueous phase was extracted with CH_2Cl_2 and the organic layer was combined and dried over Na_2SO_4 . Then the solvent was removed to give a dark blue residue which was purified by flash column chromatography with the elution of *n*-hexane/ ethyl acetate (1:2) to yield 1.61 g (55%) of the title compound as a dark blue powder. Single crystals of the title compound suitable for X-ray diffraction was recrystallized from hexane/dichloromethane (1:1).

S3. Refinement


All H atoms were positioned geometrically and refined using a riding model, with C—H = 0.96 Å (methyl), 0.97 Å (methylene) or 0.93 Å (aryl), and O—H = 0.82 Å, and with $U_{iso}(H) = 1.2U_{eq}(C)$ or $1.5U_{eq}(\text{methyl } C)$.

**Figure 1**

Molecular structure of the title compound, showing the atom labeling scheme. Displacement ellipsoids are drawn at the 30% probability level.

2-[4-(2-Hydroxyethoxy)phenyl]-4,4,5,5-tetramethyl-2-imidazoline-1-oxyl 3-oxide

Crystal data


 $M_r = 293.34$

Orthorhombic, $Pbca$

Hall symbol: -P 2ac 2ab

 $a = 8.869 (3) \text{ \AA}$
 $b = 16.050 (5) \text{ \AA}$
 $c = 20.925 (6) \text{ \AA}$
 $V = 2978.7 (16) \text{ \AA}^3$
 $Z = 8$
 $F(000) = 1256$
 $D_x = 1.308 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3005 reflections

 $\theta = 2.5\text{--}21.6^\circ$
 $\mu = 0.10 \text{ mm}^{-1}$
 $T = 296 \text{ K}$

Block, blue

 $0.26 \times 0.23 \times 0.22 \text{ mm}$

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2007)

 $T_{\min} = 0.976$, $T_{\max} = 0.979$

20164 measured reflections

2774 independent reflections

1928 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$
 $\theta_{\max} = 25.5^\circ$, $\theta_{\min} = 2.5^\circ$
 $h = -10 \rightarrow 10$
 $k = -17 \rightarrow 19$
 $l = -25 \rightarrow 25$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.163$
 $S = 0.95$

2774 reflections

195 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 0.8575P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.22 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.8584 (2)	1.12763 (14)	0.33524 (10)	0.0381 (5)
C2	0.8507 (3)	1.05283 (14)	0.38244 (10)	0.0401 (5)
C3	0.8131 (2)	1.00297 (13)	0.27718 (9)	0.0338 (5)
C4	0.7901 (2)	0.94682 (13)	0.22347 (9)	0.0333 (5)
C5	0.6890 (3)	0.88085 (13)	0.22601 (10)	0.0405 (5)
H5	0.6343	0.8719	0.2633	0.049*
C6	0.6678 (3)	0.82848 (13)	0.17472 (10)	0.0405 (5)
H6	0.5986	0.7851	0.1773	0.049*
C7	0.7503 (3)	0.84085 (13)	0.11912 (9)	0.0351 (5)
C8	0.8535 (3)	0.90560 (15)	0.11621 (10)	0.0417 (6)
H8	0.9098	0.9136	0.0792	0.050*
C9	0.8733 (3)	0.95792 (14)	0.16734 (10)	0.0399 (6)
H9	0.9427	1.0012	0.1647	0.048*
C10	0.6427 (3)	0.72306 (15)	0.06578 (10)	0.0463 (6)
H10A	0.5408	0.7395	0.0770	0.056*
H10B	0.6778	0.6829	0.0971	0.056*
C11	0.6449 (3)	0.68517 (19)	0.00040 (12)	0.0586 (7)
H11A	0.5780	0.6374	-0.0007	0.070*
H11B	0.6087	0.7255	-0.0305	0.070*
C12	0.7205 (3)	1.18423 (16)	0.33598 (13)	0.0550 (7)
H12A	0.7275	1.2236	0.3016	0.083*
H12B	0.7161	1.2135	0.3759	0.083*
H12C	0.6311	1.1512	0.3309	0.083*
C13	1.0013 (3)	1.17980 (17)	0.33876 (13)	0.0569 (7)
H13A	1.0871	1.1452	0.3300	0.085*
H13B	1.0106	1.2034	0.3807	0.085*
H13C	0.9962	1.2238	0.3077	0.085*
C14	0.7400 (4)	1.06293 (18)	0.43721 (12)	0.0653 (8)
H14A	0.6413	1.0741	0.4204	0.098*
H14B	0.7711	1.1085	0.4638	0.098*
H14C	0.7375	1.0126	0.4620	0.098*
C15	1.0041 (3)	1.02474 (19)	0.40759 (13)	0.0638 (8)
H15A	0.9929	0.9733	0.4305	0.096*
H15B	1.0440	1.0665	0.4357	0.096*
H15C	1.0720	1.0167	0.3724	0.096*

N1	0.8557 (2)	1.08321 (11)	0.27221 (8)	0.0368 (4)
N2	0.7976 (2)	0.98429 (11)	0.33930 (8)	0.0374 (5)
O1	0.8756 (2)	1.12342 (10)	0.21999 (7)	0.0558 (5)
O2	0.7595 (2)	0.91285 (10)	0.36180 (7)	0.0581 (5)
O3	0.73915 (18)	0.79412 (10)	0.06528 (7)	0.0464 (4)
O4	0.7906 (3)	0.66024 (15)	-0.01595 (10)	0.0783 (7)
H4	0.7890	0.6369	-0.0508	0.117*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0401 (12)	0.0392 (13)	0.0350 (11)	-0.0033 (10)	0.0006 (9)	-0.0095 (9)
C2	0.0493 (13)	0.0421 (13)	0.0288 (10)	-0.0015 (10)	-0.0004 (9)	-0.0067 (9)
C3	0.0397 (11)	0.0330 (12)	0.0289 (10)	0.0003 (10)	0.0010 (8)	0.0005 (8)
C4	0.0399 (12)	0.0319 (11)	0.0280 (10)	0.0023 (9)	-0.0011 (8)	0.0003 (8)
C5	0.0521 (13)	0.0402 (13)	0.0293 (10)	-0.0037 (11)	0.0089 (9)	-0.0004 (9)
C6	0.0537 (14)	0.0347 (12)	0.0331 (11)	-0.0076 (10)	0.0056 (10)	-0.0016 (9)
C7	0.0421 (12)	0.0342 (11)	0.0289 (10)	0.0037 (9)	-0.0009 (9)	-0.0045 (8)
C8	0.0444 (12)	0.0491 (14)	0.0315 (11)	-0.0040 (11)	0.0113 (9)	-0.0042 (9)
C9	0.0422 (13)	0.0423 (13)	0.0352 (11)	-0.0078 (10)	0.0065 (9)	-0.0050 (9)
C10	0.0536 (14)	0.0489 (14)	0.0363 (11)	-0.0112 (11)	-0.0006 (10)	-0.0037 (10)
C11	0.0661 (18)	0.0676 (18)	0.0420 (14)	-0.0164 (14)	-0.0063 (12)	-0.0132 (12)
C12	0.0545 (16)	0.0485 (15)	0.0621 (16)	0.0082 (12)	-0.0003 (12)	-0.0064 (12)
C13	0.0537 (16)	0.0619 (17)	0.0550 (15)	-0.0171 (13)	-0.0008 (12)	-0.0081 (13)
C14	0.092 (2)	0.0601 (17)	0.0438 (14)	-0.0058 (15)	0.0248 (14)	-0.0106 (12)
C15	0.0679 (18)	0.0708 (19)	0.0527 (15)	0.0060 (15)	-0.0233 (14)	0.0018 (13)
N1	0.0449 (11)	0.0363 (10)	0.0293 (9)	-0.0035 (8)	0.0014 (7)	-0.0010 (7)
N2	0.0505 (11)	0.0355 (10)	0.0261 (8)	-0.0023 (8)	0.0013 (7)	0.0005 (7)
O1	0.0892 (14)	0.0420 (10)	0.0363 (9)	-0.0110 (9)	0.0059 (8)	0.0083 (7)
O2	0.0993 (14)	0.0428 (10)	0.0322 (8)	-0.0155 (10)	0.0016 (8)	0.0072 (7)
O3	0.0640 (11)	0.0435 (9)	0.0317 (8)	-0.0112 (8)	0.0074 (7)	-0.0106 (6)
O4	0.0952 (17)	0.0864 (16)	0.0532 (12)	0.0100 (13)	0.0002 (10)	-0.0280 (10)

Geometric parameters (\AA , $^\circ$)

C1—N1	1.499 (3)	C10—C11	1.497 (3)
C1—C13	1.520 (3)	C10—H10A	0.9700
C1—C12	1.524 (3)	C10—H10B	0.9700
C1—C2	1.556 (3)	C11—O4	1.395 (3)
C2—N2	1.499 (3)	C11—H11A	0.9700
C2—C14	1.518 (3)	C11—H11B	0.9700
C2—C15	1.527 (4)	C12—H12A	0.9600
C3—N2	1.341 (3)	C12—H12B	0.9600
C3—N1	1.346 (3)	C12—H12C	0.9600
C3—C4	1.455 (3)	C13—H13A	0.9600
C4—C5	1.389 (3)	C13—H13B	0.9600
C4—C9	1.398 (3)	C13—H13C	0.9600
C5—C6	1.376 (3)	C14—H14A	0.9600

C5—H5	0.9300	C14—H14B	0.9600
C6—C7	1.389 (3)	C14—H14C	0.9600
C6—H6	0.9300	C15—H15A	0.9600
C7—O3	1.357 (2)	C15—H15B	0.9600
C7—C8	1.386 (3)	C15—H15C	0.9600
C8—C9	1.371 (3)	N1—O1	1.281 (2)
C8—H8	0.9300	N2—O2	1.285 (2)
C9—H9	0.9300	O4—H4	0.8200
C10—O3	1.426 (3)		
N1—C1—C13	108.57 (17)	O4—C11—C10	110.7 (2)
N1—C1—C12	106.21 (18)	O4—C11—H11A	109.5
C13—C1—C12	109.9 (2)	C10—C11—H11A	109.5
N1—C1—C2	101.00 (16)	O4—C11—H11B	109.5
C13—C1—C2	115.53 (19)	C10—C11—H11B	109.5
C12—C1—C2	114.71 (19)	H11A—C11—H11B	108.1
N2—C2—C14	109.3 (2)	C1—C12—H12A	109.5
N2—C2—C15	105.71 (19)	C1—C12—H12B	109.5
C14—C2—C15	110.4 (2)	H12A—C12—H12B	109.5
N2—C2—C1	101.41 (16)	C1—C12—H12C	109.5
C14—C2—C1	115.2 (2)	H12A—C12—H12C	109.5
C15—C2—C1	114.0 (2)	H12B—C12—H12C	109.5
N2—C3—N1	108.50 (17)	C1—C13—H13A	109.5
N2—C3—C4	126.58 (19)	C1—C13—H13B	109.5
N1—C3—C4	124.91 (18)	H13A—C13—H13B	109.5
C5—C4—C9	118.04 (18)	C1—C13—H13C	109.5
C5—C4—C3	122.21 (18)	H13A—C13—H13C	109.5
C9—C4—C3	119.75 (19)	H13B—C13—H13C	109.5
C6—C5—C4	121.61 (19)	C2—C14—H14A	109.5
C6—C5—H5	119.2	C2—C14—H14B	109.5
C4—C5—H5	119.2	H14A—C14—H14B	109.5
C5—C6—C7	119.6 (2)	C2—C14—H14C	109.5
C5—C6—H6	120.2	H14A—C14—H14C	109.5
C7—C6—H6	120.2	H14B—C14—H14C	109.5
O3—C7—C8	115.21 (18)	C2—C15—H15A	109.5
O3—C7—C6	125.3 (2)	C2—C15—H15B	109.5
C8—C7—C6	119.47 (18)	H15A—C15—H15B	109.5
C9—C8—C7	120.63 (19)	C2—C15—H15C	109.5
C9—C8—H8	119.7	H15A—C15—H15C	109.5
C7—C8—H8	119.7	H15B—C15—H15C	109.5
C8—C9—C4	120.6 (2)	O1—N1—C3	125.90 (17)
C8—C9—H9	119.7	O1—N1—C1	120.55 (17)
C4—C9—H9	119.7	C3—N1—C1	113.06 (16)
O3—C10—C11	108.08 (19)	O2—N2—C3	125.57 (17)
O3—C10—H10A	110.1	O2—N2—C2	121.12 (16)
C11—C10—H10A	110.1	C3—N2—C2	112.80 (17)
O3—C10—H10B	110.1	C7—O3—C10	118.67 (16)
C11—C10—H10B	110.1	C11—O4—H4	109.5

H10A—C10—H10B	108.4		
N1—C1—C2—N2	−16.7 (2)	N2—C3—N1—O1	−177.2 (2)
C13—C1—C2—N2	−133.60 (19)	C4—C3—N1—O1	3.5 (3)
C12—C1—C2—N2	97.0 (2)	N2—C3—N1—C1	−5.2 (2)
N1—C1—C2—C14	−134.5 (2)	C4—C3—N1—C1	175.45 (19)
C13—C1—C2—C14	108.6 (2)	C13—C1—N1—O1	−51.1 (3)
C12—C1—C2—C14	−20.8 (3)	C12—C1—N1—O1	67.0 (2)
N1—C1—C2—C15	96.4 (2)	C2—C1—N1—O1	−172.97 (19)
C13—C1—C2—C15	−20.5 (3)	C13—C1—N1—C3	136.5 (2)
C12—C1—C2—C15	−149.9 (2)	C12—C1—N1—C3	−105.4 (2)
N2—C3—C4—C5	29.5 (3)	C2—C1—N1—C3	14.6 (2)
N1—C3—C4—C5	−151.3 (2)	N1—C3—N2—O2	−179.3 (2)
N2—C3—C4—C9	−149.5 (2)	C4—C3—N2—O2	0.0 (4)
N1—C3—C4—C9	29.7 (3)	N1—C3—N2—C2	−7.4 (2)
C9—C4—C5—C6	−1.3 (3)	C4—C3—N2—C2	171.86 (19)
C3—C4—C5—C6	179.6 (2)	C14—C2—N2—O2	−49.8 (3)
C4—C5—C6—C7	0.7 (4)	C15—C2—N2—O2	69.0 (3)
C5—C6—C7—O3	−179.5 (2)	C1—C2—N2—O2	−171.79 (19)
C5—C6—C7—C8	0.3 (3)	C14—C2—N2—C3	138.0 (2)
O3—C7—C8—C9	179.1 (2)	C15—C2—N2—C3	−103.3 (2)
C6—C7—C8—C9	−0.8 (3)	C1—C2—N2—C3	15.9 (2)
C7—C8—C9—C4	0.2 (3)	C8—C7—O3—C10	176.0 (2)
C5—C4—C9—C8	0.8 (3)	C6—C7—O3—C10	−4.1 (3)
C3—C4—C9—C8	179.9 (2)	C11—C10—O3—C7	177.4 (2)
O3—C10—C11—O4	60.5 (3)		

Hydrogen-bond geometry (Å, °)

Cg2 is the centroid of the benzene C4—C9 ring.

D—H···A	D—H	H···A	D···A	D—H···A
O4—H4···O2 ⁱ	0.82	2.01	2.828 (3)	173
C12—H12C···O1 ⁱⁱ	0.96	2.54	3.418 (3)	152
C15—H15C···Cg2 ⁱⁱⁱ	0.96	2.80	3.570 (3)	138

Symmetry codes: (i) $x, -y+3/2, z-1/2$; (ii) $x-1/2, y, -z+1/2$; (iii) $x+1/2, y, -z+1/2$.