

**catena-Poly[bis(dimethylazanium)
[[chloridocopper(II)]-di- μ -chlorido-
[chloridocopper(II)]-di- μ -azido- $\kappa^4 N:N$]]**

Wei-Yi Zhang,^a Li Yang^b and Jie Liu^{c*}

^aDepartment of Obstetrics and Gynecology, The First Affiliated Hospital of Henan University, of Traditional Chinese Medicine, Zhengzhou, 450008, People's Republic of China, ^bHenan Medical College for Staff and Workers, Zhengzhou, 451191, People's Republic of China, and ^cDepartment of Urology, Henan Provincial People's Hospital, Zhengzhou, 450003, People's Republic of China
Correspondence e-mail: liu_jie1011@163.com


Received 11 September 2011; accepted 24 November 2011

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(N-C) = 0.005$ Å;
 R factor = 0.027; wR factor = 0.066; data-to-parameter ratio = 21.3.

The crystal structure of the title complex, $\{(C_2H_8N)[CuCl_2(N_3)]\}_n$, exhibits inorganic chains consisting of Cu(II) cations as well azide and chloride anions. The chains, made up from Cu—Cl—Cu—N—Cu linkages, are aligned parallel to the c axis. This architecture is further stabilized by a number of N—H···Cl hydrogen bonds involving the protonated charge-compensating dimethylamine cations and chloride atoms.

Related literature

For background to polynuclear complexes, see Goher *et al.* (2000); Liu *et al.* (2008); Ribas *et al.* (1994); Saha *et al.* (2005); Vicente *et al.* (1993); Wang *et al.* (2008). For di- or polyalkylamines as templates, see: Cheetham *et al.* (1999); Hagrman *et al.* (1999). For related copper(II) complexes, see: Mautner *et al.* (1999).



Experimental

Crystal data

$(C_2H_8N)[CuCl_2(N_3)]$

$M_r = 222.57$

Monoclinic, $C2/c$

$a = 15.348$ (5) Å

$b = 11.089$ (2) Å

$c = 10.729$ (2) Å

$\beta = 119.73$ (2)°

$V = 1585.7$ (7) Å³

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 3.35$ mm⁻¹

$T = 298$ K

$0.14 \times 0.10 \times 0.08$ mm

Data collection

Bruker APEXII CCD
diffractometer

Absorption correction: multi-scan
(SADABS; Sheldrick, 2003)

$T_{\min} = 0.651$, $T_{\max} = 0.775$

3510 measured reflections
1811 independent reflections
1251 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$
 $wR(F^2) = 0.066$
 $S = 0.94$
1811 reflections
85 parameters

13 restraints
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.41$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.38$ e Å⁻³

Table 1
Selected bond lengths (Å).

Cu1—N1 ⁱ	1.987 (2)	Cu1—Cl1	2.2729 (9)
Cu1—N1	2.002 (2)	Cu1—Cl1 ⁱⁱ	2.8860 (13)
Cu1—Cl2	2.2527 (8)	Cu1—Cu1 ⁱ	3.1460 (7)

Symmetry codes: (i) $-x, -y + 1, -z$; (ii) $-x, y, -z + \frac{1}{2}$.

Table 2
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N4—H1···Cl1 ⁱⁱⁱ	0.99	2.41	3.331 (3)	154
N4—H2···Cl2 ⁱⁱ	0.87	2.50	3.257 (3)	146
N4—H2···Cl1	0.87	2.82	3.270 (2)	114
N4—H2···Cl2	0.87	2.92	3.340 (3)	112

Symmetry codes: (ii) $-x, y, -z + \frac{1}{2}$; (iii) $-x - \frac{1}{2}, -y + \frac{1}{2}, -z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: publCIF (Westrip, 2010).

This study was supported by the Doctoral Research Fund of Henan Chinese Medicine (BSJJ2009-42).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZB2020).

References

- Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cheetham, A. K., Ferey, G. & Loiseau, T. (1999). *Angew. Chem. Int. Ed.* **38**, 3268–3292.
- Goher, M. A. S., Cano, J., Journaux, Y., Abu-Youssef, M. A. M., Mautner, F. A., Escuer, A. & Vicente, R. (2000). *Chem. Eur. J.* **6**, 778–784.
- Hagrman, P. J., Hagrman, D. & Zubietta, J. (1999). *Angew. Chem. Int. Ed.* **38**, 2638–2684.
- Liu, T., Yang, Y. F., Wang, Z. M. & Gao, S. (2008). *Chem. Asian J.* **3**, 950–957.
- Mautner, F. A., Hanna, S., Cortes, R., Lezama, L., Barandika, M. G. & Rojo, T. (1999). *Inorg. Chem.* **38**, 4647–4652.
- Ribas, J., Monfort, M., Diaz, C., Bastos, C. & Solans, X. (1994). *Inorg. Chem.* **33**, 484–489.
- Saha, S., Koner, S., Tuchagues, J. P., Boudalis, A. K., Okamoto, K. I., Banerjee, S. & Mal, D. (2005). *Inorg. Chem.* **44**, 6379–6385.
- Sheldrick, G. M. (2003). SADABS. University of Göttingen, Germany.

metal-organic compounds

- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Vicente, R., Escuer, A., Ribas, J., Fallah, M. S., Solans, X. & Font-Bardid, M. (1993). *Inorg. Chem.* **32**, 1920–1924.
- Wang, X. Y., Wang, Z. M. & Gao, S. (2008). *Chem. Commun.* pp. 281–294.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, m1877–m1878 [https://doi.org/10.1107/S1600536811050537]

catena-Poly[bis(dimethylazanium) [[chloridocopper(II)]-di- μ -chlorido-[chloridocopper(II)]-di- μ -azido- $\kappa^4N:N$]]

Wei-Yi Zhang, Li Yang and Jie Liu

S1. Comment

It is well known that the azide ion is a versatile ligand, and its versatility and efficiency lie in its functionality as a terminal monodentate and a bridging bi-, tri-, and tetradentate ligand. Because of this unique capability, azide attracts a lot of attention in the design of mono- or multidimensional metal-assembled azido complexes. (Vicente *et al.*, 1993; Ribas *et al.*, 1994; Goher *et al.*, 2000; Saha *et al.*, 2005; Liu *et al.*, 2008). Having control over the molecular dimensions and geometry of the metal-ligand moiety in the compounds may lead to the control over their magnetic properties.(Wang *et al.*, 2008). Di- or polyalkylamines, if protonated, could be conveniently used as cationic templates, and they have been widely employed in making metal oxalates, metal phosphates, and oxometalates.(Cheetham *et al.*, 1999; Hagrman *et al.*, 1999).In order to study the coordination behavior of the azide ion and templates, we synthesized herein the title complex $[(\text{NH}_2(\text{CH}_3)_2)(\text{CuN}_3\text{Cl}_2)]_n$. As shown in Figure 1, each asymmetric unit contains one Cu(II) atom, two chloride atoms, one azide atom and one dimethylamine cation. This architecture is further stabilized by a number of N—H···Cl hydrogen bonds involving the protonated charge-compensating dimethylamine cations and chloride atoms.(Figure 2). The bond distances for Cu—N are 1.984 (2) and 2.001 (2) Å, respectively. and the angles for N—Cu—Cl are between 92.86 (6) and 167.48 (6)°. The Cu—Cl bond lengths are 2.2526 (8) Å, 2.2725 (10) Å, respectively. and the bond angles for N—Cu—N and Cl—Cu—Cl are 75.74 (10) and 94.61 (2)°, respectively. These bond distances and bond angles are in agreement with those found in the reported copper compounds(Mautner *et al.*, 1999).


S2. Experimental

A mixture of methanol and water (1:1, 2 ml) was gently layered on the top of a solution of $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ (37.1 mg, 0.1 mmol) in water (3 ml). A solution of dimethylamine (18 mg, 0.4 mmol), NaN_3 (13 mg, 0.2 mmol) and hydrochloric acid (40.5 mg, 0.4 mmol, 36%) in methanol (10 ml) was added carefully as the third layer. Green crystals were obtained after 3 weeks, washed with ethanol and ether, and dried in air.

S3. Refinement

During refinement, H atoms were placed in calculated positions and allowed to ride, with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$.

Data collection: *APEX2*(Bruker, 2007); cell refinemetnet: *SAINT*(Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structures: *SHELXS97*(Sheldrick, 2008); program(s) used to refine structures: *SHELXL97*(Sheldrick, 2008); molecular graphics: *SHELXTL*(Sheldrick, 2008); software used to prepare material for publication: *publCIF*.

**Figure 1**

The molecular structure for title compound. Displacement ellipsoids at the 30% probability level. Symmetry codes: (i) -
 $x, -y + 1, -z$

catena-Poly[bis(dimethylazanium) [[chloridocopper(II)]-di- μ -chlorido-[chloridocopper(II)]-di- μ -azido- $\kappa^4N:N$]]

Crystal data



$M_r = 222.57$

Monoclinic, $C2/c$

$a = 15.348 (5)$ Å

$b = 11.089 (2)$ Å

$c = 10.729 (2)$ Å

$\beta = 119.73 (2)^\circ$

$V = 1585.7 (7)$ Å³

$Z = 8$

$F(000) = 888$

$D_x = 1.865 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1037 reflections

$\theta = 2.6\text{--}24.6^\circ$

$\mu = 3.35 \text{ mm}^{-1}$

$T = 298$ K

Block, green

$0.14 \times 0.10 \times 0.08$ mm

Data collection

Bruker APEXII CCD
 diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and ω scans

Absorption correction: multi-scan
 $(SADABS; Sheldrick, 2003)$

$T_{\min} = 0.651$, $T_{\max} = 0.775$

3510 measured reflections

1811 independent reflections

1251 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.7^\circ$

$h = -19 \rightarrow 19$

$k = -14 \rightarrow 14$

$l = -13 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.027$

$wR(F^2) = 0.066$

$S = 0.94$

1811 reflections

85 parameters

13 restraints

Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0361P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.012$
 $\Delta\rho_{\max} = 0.41 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.38 \text{ e \AA}^{-3}$

Extinction correction: *SHELXL*,
 $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.0036 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	-0.03408 (2)	0.39407 (3)	0.06367 (3)	0.03133 (14)
Cl1	-0.14064 (5)	0.39423 (5)	0.15347 (8)	0.03677 (18)
Cl2	-0.03672 (5)	0.19228 (5)	0.03779 (7)	0.0404 (2)
N1	-0.04383 (19)	0.57303 (19)	0.0352 (3)	0.0409 (6)
N2	-0.08322 (18)	0.64469 (19)	0.0760 (3)	0.0388 (6)
N3	-0.1205 (2)	0.7121 (2)	0.1134 (3)	0.0659 (9)
N4	-0.1737 (2)	0.1071 (2)	0.1860 (3)	0.0492 (6)
H1	-0.2164	0.1291	0.0834	0.059*
H2	-0.1180	0.1479	0.2287	0.059*
C1	-0.1482 (3)	-0.0211 (3)	0.1877 (4)	0.0631 (9)
H1A	-0.1133	-0.0496	0.2850	0.095*
H1B	-0.1062	-0.0303	0.1455	0.095*
H1C	-0.2087	-0.0669	0.1337	0.095*
C2	-0.2303 (3)	0.1339 (3)	0.2608 (4)	0.0579 (8)
H2A	-0.2921	0.0894	0.2167	0.087*
H2B	-0.2446	0.2186	0.2544	0.087*
H2C	-0.1911	0.1110	0.3598	0.087*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0392 (2)	0.02505 (19)	0.0368 (2)	0.00100 (14)	0.02419 (17)	0.00218 (13)
Cl1	0.0405 (4)	0.0355 (4)	0.0434 (4)	0.0016 (3)	0.0277 (3)	0.0042 (3)
Cl2	0.0520 (5)	0.0270 (3)	0.0459 (5)	-0.0020 (3)	0.0271 (4)	-0.0012 (3)
N1	0.0609 (17)	0.0280 (11)	0.0535 (16)	0.0035 (10)	0.0433 (14)	0.0048 (10)
N2	0.0495 (15)	0.0285 (12)	0.0501 (16)	0.0050 (10)	0.0336 (13)	0.0062 (10)
N3	0.087 (2)	0.0483 (17)	0.092 (2)	0.0157 (15)	0.067 (2)	0.0034 (14)
N4	0.0569 (14)	0.0425 (11)	0.0524 (14)	-0.0084 (10)	0.0303 (11)	-0.0003 (9)
C1	0.0676 (17)	0.0421 (14)	0.0641 (17)	-0.0036 (14)	0.0208 (15)	0.0025 (13)
C2	0.0592 (17)	0.0641 (15)	0.0575 (17)	-0.0108 (14)	0.0342 (14)	0.0001 (13)

Geometric parameters (\AA , $^{\circ}$)

Cu1—N1 ⁱ	1.987 (2)	N4—C1	1.471 (4)
Cu1—N1	2.002 (2)	N4—C2	1.477 (4)
Cu1—Cl2	2.2527 (8)	N4—H1	0.9931
Cu1—Cl1	2.2729 (9)	N4—H2	0.8693
Cu1—Cl1 ⁱⁱ	2.8860 (13)	C1—H1A	0.9600
Cu1—Cu1 ⁱ	3.1460 (7)	C1—H1B	0.9600
Cl1—Cu1 ⁱⁱ	2.8860 (13)	C1—H1C	0.9600
N1—N2	1.205 (3)	C2—H2A	0.9600
N1—Cu1 ⁱ	1.987 (2)	C2—H2B	0.9600
N2—N3	1.129 (3)	C2—H2C	0.9600
N1 ⁱ —Cu1—N1	75.87 (10)	N3—N2—N1	179.6 (3)
N1 ⁱ —Cu1—Cl2	95.48 (7)	C1—N4—C2	114.3 (2)
N1—Cu1—Cl2	166.02 (7)	C1—N4—H1	106.0
N1 ⁱ —Cu1—Cl1	167.54 (7)	C2—N4—H1	108.0
N1—Cu1—Cl1	92.79 (7)	C1—N4—H2	108.1
Cl2—Cu1—Cl1	94.60 (3)	C2—N4—H2	107.4
N1 ⁱ —Cu1—Cl1 ⁱⁱ	94.01 (8)	H1—N4—H2	113.1
N1—Cu1—Cl1 ⁱⁱ	96.89 (7)	N4—C1—H1A	109.5
Cl2—Cu1—Cl1 ⁱⁱ	94.63 (2)	N4—C1—H1B	109.5
Cl1—Cu1—Cl1 ⁱⁱ	92.46 (3)	H1A—C1—H1B	109.5
N1 ⁱ —Cu1—Cu1 ⁱ	38.11 (6)	N4—C1—H1C	109.5
N1—Cu1—Cu1 ⁱ	37.76 (7)	H1A—C1—H1C	109.5
Cl2—Cu1—Cu1 ⁱ	132.67 (3)	H1B—C1—H1C	109.5
Cl1—Cu1—Cu1 ⁱ	130.36 (2)	N4—C2—H2A	109.5
Cl1 ⁱⁱ —Cu1—Cu1 ⁱ	96.92 (2)	N4—C2—H2B	109.5
Cu1—Cl1—Cu1 ⁱⁱ	87.54 (3)	H2A—C2—H2B	109.5
N2—N1—Cu1 ⁱ	128.05 (19)	N4—C2—H2C	109.5
N2—N1—Cu1	127.70 (19)	H2A—C2—H2C	109.5
Cu1 ⁱ —N1—Cu1	104.13 (10)	H2B—C2—H2C	109.5

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x, y, -z+1/2$.

Hydrogen-bond geometry (\AA , $^{\circ}$)

D—H···A	D—H	H···A	D···A	D—H···A
N4—H1···Cl1 ⁱⁱⁱ	0.99	2.41	3.331 (3)	154
N4—H2···Cl2 ⁱⁱ	0.87	2.50	3.257 (3)	146
N4—H2···Cl1	0.87	2.82	3.270 (2)	114
N4—H2···Cl2	0.87	2.92	3.340 (3)	112

Symmetry codes: (ii) $-x, y, -z+1/2$; (iii) $-x-1/2, -y+1/2, -z$.