

1-Prop-2-ynyl-1*H*-benzimidazol-2-amine

Alka Agarwal,^{a*} Manavendra K. Singh^a and Satish K. Awasthi^{b*}

^aDepartment of Medicinal Chemistry, Institute of Medical Sciences, Banaras Hindu University, Varanasi 225 001, UP, India, and ^bChemical Biology Laboratory, Department of Chemistry, University of Delhi, Delhi 110 007, India
Correspondence e-mail: dralka@bhu.ac.in,


Received 24 September 2011; accepted 15 October 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.054; wR factor = 0.137; data-to-parameter ratio = 25.2.

In the title compound, $\text{C}_{10}\text{H}_9\text{N}_3$, the benzimidazol-2-amine and $\text{CH}_2-\text{C}\equiv\text{CH}$ units are not coplanar, with a dihedral angle of 60.36° between their mean planes. The crystal structure is stabilized by intermolecular $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonding and $\pi-\pi$ interactions [centroid–centroid distances $3.677(1)$ and $3.580(1)\text{ \AA}$], assembling the molecules into a supramolecular structure with a three-dimensional network.

Related literature

For the biological activities of benzimidazoles, see: Nawrocka *et al.* (1999); Cuberens & Contijoch (1997); Mor *et al.* (2004); de Dios *et al.* (2005). For polyfunctionality and antiviral activity of 2-aminobenzimidazoles, see: Garuti & Roberti (2002); Andries *et al.* (2003). For antiproliferative properties, see: Garuti *et al.* (1998); Nawrocka *et al.* (2005). For inhibition activity against various strains of bacteria, fungi and yeasts, see: Nofal *et al.* (2002); Omar *et al.* (1996); Del Poenta *et al.* (1999). For structural analysis of small molecules, see: Singh, Agarwal, Mahawar & Awasthi (2011); Singh, Singh *et al.* (2011); Singh, Agarwal & Awasthi (2011); Agarwal *et al.* (2011). For the synthesis, see: Lilienkampf *et al.* (2009).



Experimental

Crystal data

$\text{C}_{10}\text{H}_9\text{N}_3$
 $M_r = 171.20$
Monoclinic, $C2/c$
 $a = 15.385(2)\text{ \AA}$
 $b = 12.1433(12)\text{ \AA}$

$c = 9.4653(10)\text{ \AA}$
 $\beta = 95.755(11)^\circ$
 $V = 1759.5(3)\text{ \AA}^3$
 $Z = 8$
Mo $K\alpha$ radiation

$\mu = 0.08\text{ mm}^{-1}$
 $T = 293\text{ K}$

$0.39 \times 0.36 \times 0.20\text{ mm}$

Data collection

Oxford Diffraction Xcalibur
Sapphire3 diffractometer
Absorption correction: multi-scan
(*CrysAlis PRO*; Oxford
Diffraction, 2009)
 $T_{\min} = 0.677$, $T_{\max} = 1.000$

9853 measured reflections
3178 independent reflections
2505 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$
 $wR(F^2) = 0.137$
 $S = 1.08$
3178 reflections
126 parameters

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\max} = 0.25\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.18\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N3—H1N3 \cdots N1 ⁱ	0.90 (2)	2.36 (2)	3.1823 (18)	153.5 (18)
N3—H2N3 \cdots N1 ⁱⁱ	0.860 (18)	2.241 (19)	3.0591 (15)	158.6 (15)

Symmetry codes: (i) $-x + 1, -y, -z + 2$; (ii) $x, -y, z - \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2009); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

AA is thankful to the University Grant Commission (UGC) (scheme No. 34–311/2008), New Delhi, and Banaras Hindu University, Varanasi, India, for financial assistance. The authors are very grateful to the Department of Chemistry, Banaras Hindu University, Varanasi, for providing the single-crystal X-ray diffraction facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZJ207).

References

- Agarwal, A., Singh, M. K., Singh, S., Bhattacharya, S. & Awasthi, S. K. (2011). *Acta Cryst. E67*, o2637–o2638.
- Andries, K., Moeremans, M., Gevers, T., Willebroeds, R., Sommen, C., Lacrampe, J., Janssens, F. & Wyde, P. R. (2003). *Antiviral. Res.* **60**, 209–219.
- Cuberens, M. & Contijoch, M. R. (1997). *Chem. Pharm. Bull.* **45**, 1287–1292.
- Del Poenta, M., Bixel, A. S., Barchiesi, F., Tidwell, R. R., Boykin, D., Scalise, G. & Perfect, J. R. (1999). *J. Antimicrob. Chemother.* **44**, 223–228.
- Dios, A. de, *et al.* (2005). *J. Med. Chem.* **48**, 2270–2273.
- Garuti, L. & Roberti, M. (2002). *Bioorg. Med. Chem. Lett.* **12**, 2707–2710.
- Garuti, L., Varoli, L., Cermelli, C., Baggio, G., Lupo, L., Malagoli, M. & Castelli, M. (1998). *Anti-Cancer Drug Des.* **13**, 969–980.
- Lilienkampf, A., Mao, J., Wan, B., Wang, Y., Franzblau, S. G. & Kozikowski, A. P. (2009). *J. Med. Chem.* **52**, 2109–2118.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Mor, M., Bordi, F., Silva, C., Rivara, S., Zuliana, V., Vacondio, F., Rivara, M., Barocelli, E., Bertoni, S., Ballabeni, V., Magnanini, F., Impicciatore, M. & Piazzi, P. V. (2004). *Bioorg. Med. Chem.* **12**, 663–674.
- Nawrocka, W., Sztuba, B., Liszkiewicz, H., Kowalska, M. W., Wietrzyk, J., Nevozhai, D. & Opolski, A. (2005). *Pol. J. Chem.* **79**, 709–716.

organic compounds

- Nawrocka, W., Zimecki, M., Kuznicki, T. & Kowalska, M. W. (1999). *Arch. Pharm.* **332**, 85–90.
- Nofal, Z. M., Fahmy, H. H. & Mohamed, H. S. (2002). *Arch. Pharm. Res.* **25**, 250–257.
- Omar, M. T., Fahmy, H. H. & Mohamed, H. S. (1996). *Egypt. J. Pharm. Sci.* **37**, 606–620.
- Oxford Diffraction (2009). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Singh, M. K., Agarwal, A. & Awasthi, S. K. (2011). *Acta Cryst. E* **67**, o1137.
- Singh, M. K., Agarwal, A., Mahawar, C. & Awasthi, S. K. (2011). *Acta Cryst. E* **67**, o1382.
- Singh, S., Singh, M. K., Agarwal, A., Hussain, F. & Awasthi, S. K. (2011). *Acta Cryst. E* **67**, o1616–o1617.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, o3213–o3214 [https://doi.org/10.1107/S1600536811042772]

1-Prop-2-ynyl-1H-benzimidazol-2-amine

Alka Agarwal, Manavendra K. Singh and Satish K. Awasthi

S1. Comment

The 2-aminobenzimidazole compounds showed different biological activities such as immunotropic, diuretic, antihistamine as well as highly selective p38a MAP inhibition properties (Nawrocka *et al.*, 1999, Cuberens & Contijoch 1997, Mor *et al.*, 2004, De Dios *et al.*, 2005). The polyfunctionality of the 2-aminobenzimidazole molecule is due to the cyclic guanidine moiety which acts as a building block for the synthesis of a large number of benzimidazole derivatives. The various benzimidazole derivatives such as enviradene and enviroxime and other 2-aminobenzimidazole derivatives have been synthesized and screened for antiviral activity (Garuti & Roberti 2002, Andries *et al.*, 2003). Moreover, a number of 2-aminobenzimidazoles have exhibited antiproliferative properties (Garuti, Varoli *et al.*, 1998, Nawrocka *et al.*, 2005). Further, different substituted 2-aminobenzimidazoles have been found to possess inhibition activity against various strains of bacteria, fungi and yeasts in vivo and in vitro (Nofal *et al.*, 2002, Omar *et al.*, 1996, Del Poenta *et al.*, 1999). In continuation with our recent work on structural analysis of small molecule (Singh, Agarwal, Mahawar & Awasthi 2011, Singh, Singh, Agarwal, Hussain & Awasthi 2011, Singh, Agarwal & Awasthi 2011, Agarwal *et al.*, 2011), we wish to report here the crystal structure of 1-prop-2-ynyl-1H-benzimidazol-2-ylamine (Figure1).

In the title compound, the C7-N3 single bond length (1.35 Å) is shorter than normal C-N (1.47 Å) bond suggesting a delocalized double bond in benzimidazole moiety. Again, in the crystal structure, the title compound is stabilized by intermolecular N-H···N hydrogen bonding and π ··· π interaction assembling, thus forming into supramolecule type assembly with a tri-dimensional network (Figure 2, Table 1). π ··· π interactions form a dimer, the ring A and B of an benzimidazole skeleton stacks with the ring B and A of another adjacent benzimidazole skeleton, respectively. The distance of CgA and CgB is 3.641 Å, where CgA and CgB are the center of ring A and B, respectively and the centroid - centroid distance between two adjacent benzimidazole ring is 3.580 Å and the bond distance between atoms C5···C10 is found to be 3.282 Å as well as the bond distance between atoms C1···C5 is found to be 3.369 Å, which helps in overall crystal structure packing as well as stabilization (Figure 2). In the molecule, 2-aminobenzimidazole ring and CH₂-C≡CH are non-planar with dihedral angle is 60.36. The CCDC No. of crystal is 843876.

S2. Experimental

The synthesis of the title compound was carried out according to the published procedure (Lilienkampf *et al.*, 2009). Briefly, to a solution of 2- aminobenzimidazole (0.50 g m, 3.76 mmol) in dry acetone was added anhydrous K₂CO₃ (3.112 g m, 22.55 mmol) and reaction mixture was further refluxed for 15–30 minutes. Subsequently, KI (0.312 g m, 1.88 mmol) and propargyl bromide (0.39 ml, 4.37 mmol) were added into it and further refluxed the reaction mixture for 18 hrs. After this period, the reaction mixture was cooled, filtered and the filtrate was concentrated *invacuo* up to 10 ml and left for few days at room temperature which gave transparent crystals after slow evaporation of solvent. Yield = 30%, MS (Macromass G) m/z = 188 (M^+), R_f0.59 (98:2, CH₂Cl₂: MeOH) Elemental analysis (Perkin –Elmer 240°C elemental analyzer)Calculated for: C₁₀ H₉ N₃(%) C- 70.2, H-5.3, N -24.5 found C-69.9, H-5.5, N -24.6. ¹H NMR (CDCl₃), 7.71–

7.68 (m, 1H), 7.44–7.42 (m, 1H), 7.26–7.21 (m, 1H), 7.07–7.02 (m, 1H), 4.88 (s, 2H, NH₂), 4.71 (s, 2H, CH₂), 2.43 (s, 1H, CH).

S3. Refinement

All H atoms were located from difference Fourier map (range of C—H = 0.93 - 0.97 Å and N—H = 0.86 (18) - 0.90 (2) Å) and allowed to refine freely.






Figure 1

ORTEP diagram of the molecule with thermal ellipsoids drawn at 50% probability level Color code: White: C; blue: N; white: H.

**Figure 2**

Packing diagram of molecule viewed through a plane showing supramolecule arrangement and intermolecular N—H···N (red and purple doted line) hydrogen bonding, C—H···C (Green doted line) interactions and $\pi\cdots\pi$ (magneta and light blue doted line) interaction.

**Figure 3**

The synthesis of the title compound.

1-Prop-2-ynyl-1H-benzimidazol-2-amine

Crystal data

$C_{10}H_9N_3$
 $M_r = 171.20$
Monoclinic, $C2/c$
Hall symbol: -C 2yc
 $a = 15.385 (2)$ Å
 $b = 12.1433 (12)$ Å
 $c = 9.4653 (10)$ Å
 $\beta = 95.755 (11)^\circ$
 $V = 1759.5 (3)$ Å³
 $Z = 8$

$F(000) = 720.0$
 $D_x = 1.293 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 2285 reflections
 $\theta = 3.3\text{--}29.2^\circ$
 $\mu = 0.08 \text{ mm}^{-1}$
 $T = 293 \text{ K}$
Block, clear white
 $0.39 \times 0.36 \times 0.20$ mm

Data collection

Oxford Diffraction Xcalibur Sapphire3 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2009)
 $T_{\min} = 0.677$, $T_{\max} = 1.000$

9853 measured reflections
 3178 independent reflections
 2505 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$
 $\theta_{\max} = 32.5^\circ$, $\theta_{\min} = 3.0^\circ$
 $h = -19 \rightarrow 23$
 $k = -18 \rightarrow 18$
 $l = -13 \rightarrow 14$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.054$
 $wR(F^2) = 0.137$
 $S = 1.08$
 3178 reflections
 126 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0598P)^2 + 0.678P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.006$
 $\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
H2N3	0.4017 (11)	-0.0261 (14)	0.750 (2)	0.056 (5)*
H1N3	0.4441 (14)	-0.0599 (17)	0.892 (2)	0.082 (6)*
N2	0.35287 (7)	0.16350 (8)	0.82062 (9)	0.0344 (2)
N1	0.40855 (7)	0.10067 (8)	1.03481 (10)	0.0370 (2)
C5	0.38413 (7)	0.21017 (9)	1.04715 (11)	0.0327 (2)
C7	0.38975 (8)	0.07726 (9)	0.89807 (11)	0.0347 (2)
N3	0.40152 (10)	-0.02285 (9)	0.84084 (12)	0.0507 (3)
C6	0.34953 (7)	0.25096 (9)	0.91465 (11)	0.0332 (2)
C4	0.39113 (9)	0.27854 (11)	1.16533 (13)	0.0416 (3)
H4	0.4128	0.2525	1.2544	0.050*
C9	0.38623 (9)	0.24521 (11)	0.60031 (12)	0.0446 (3)
C3	0.36468 (10)	0.38702 (11)	1.14576 (15)	0.0495 (3)
H3	0.3700	0.4350	1.2227	0.059*
C8	0.32884 (9)	0.16969 (11)	0.66794 (12)	0.0448 (3)
H8A	0.2689	0.1947	0.6500	0.054*

H8B	0.3326	0.0969	0.6268	0.054*
C1	0.32143 (9)	0.35828 (10)	0.89445 (15)	0.0439 (3)
H1	0.2979	0.3839	0.8062	0.053*
C2	0.33019 (10)	0.42580 (10)	1.01311 (17)	0.0508 (3)
H2	0.3126	0.4989	1.0040	0.061*
C10	0.43272 (11)	0.30823 (14)	0.55223 (17)	0.0580 (4)
H10	0.4697	0.3583	0.5140	0.070*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N2	0.0425 (5)	0.0342 (5)	0.0259 (4)	0.0015 (4)	0.0010 (3)	0.0029 (3)
N1	0.0490 (6)	0.0362 (5)	0.0257 (4)	0.0108 (4)	0.0030 (4)	0.0004 (3)
C5	0.0345 (5)	0.0338 (5)	0.0301 (5)	0.0030 (4)	0.0054 (4)	0.0003 (4)
C7	0.0417 (6)	0.0349 (5)	0.0277 (5)	0.0055 (4)	0.0048 (4)	0.0016 (4)
N3	0.0769 (9)	0.0425 (6)	0.0319 (5)	0.0186 (6)	0.0013 (5)	-0.0051 (4)
C6	0.0347 (5)	0.0322 (5)	0.0331 (5)	-0.0005 (4)	0.0051 (4)	0.0026 (4)
C4	0.0477 (7)	0.0439 (6)	0.0336 (5)	0.0033 (5)	0.0055 (5)	-0.0059 (5)
C9	0.0586 (8)	0.0469 (7)	0.0289 (5)	0.0108 (6)	0.0069 (5)	0.0047 (5)
C3	0.0562 (8)	0.0403 (6)	0.0534 (8)	0.0005 (5)	0.0118 (6)	-0.0139 (6)
C8	0.0554 (8)	0.0482 (7)	0.0289 (5)	-0.0018 (6)	-0.0051 (5)	0.0047 (5)
C1	0.0481 (7)	0.0341 (5)	0.0491 (7)	0.0021 (5)	0.0029 (5)	0.0077 (5)
C2	0.0554 (8)	0.0308 (5)	0.0669 (9)	0.0035 (5)	0.0099 (7)	-0.0020 (6)
C10	0.0669 (10)	0.0587 (8)	0.0513 (8)	0.0076 (7)	0.0205 (7)	0.0105 (7)

Geometric parameters (\AA , ^\circ)

N2—C7	1.3684 (14)	C4—H4	0.9300
N2—C6	1.3899 (14)	C9—C10	1.170 (2)
N2—C8	1.4573 (14)	C9—C8	1.4641 (19)
N1—C7	1.3284 (14)	C3—C2	1.395 (2)
N1—C5	1.3898 (14)	C3—H3	0.9300
C5—C4	1.3885 (16)	C8—H8A	0.9700
C5—C6	1.4028 (15)	C8—H8B	0.9700
C7—N3	1.3505 (15)	C1—C2	1.386 (2)
N3—H2N3	0.860 (18)	C1—H1	0.9300
N3—H1N3	0.90 (2)	C2—H2	0.9300
C6—C1	1.3803 (16)	C10—H10	0.9300
C4—C3	1.3856 (19)		
C7—N2—C6	106.36 (9)	C5—C4—H4	121.2
C7—N2—C8	128.39 (10)	C10—C9—C8	176.81 (15)
C6—N2—C8	125.00 (10)	C4—C3—C2	121.42 (12)
C7—N1—C5	104.62 (9)	C4—C3—H3	119.3
C4—C5—N1	129.79 (11)	C2—C3—H3	119.3
C4—C5—C6	120.06 (10)	N2—C8—C9	111.21 (11)
N1—C5—C6	110.13 (9)	N2—C8—H8A	109.4
N1—C7—N3	123.96 (11)	C9—C8—H8A	109.4

N1—C7—N2	113.34 (10)	N2—C8—H8B	109.4
N3—C7—N2	122.63 (10)	C9—C8—H8B	109.4
C7—N3—H2N3	117.1 (11)	H8A—C8—H8B	108.0
C7—N3—H1N3	110.8 (13)	C6—C1—C2	116.27 (12)
H2N3—N3—H1N3	116.2 (17)	C6—C1—H1	121.9
C1—C6—N2	131.60 (11)	C2—C1—H1	121.9
C1—C6—C5	122.87 (11)	C1—C2—C3	121.79 (12)
N2—C6—C5	105.52 (9)	C1—C2—H2	119.1
C3—C4—C5	117.57 (12)	C3—C2—H2	119.1
C3—C4—H4	121.2	C9—C10—H10	180.0
C7—N1—C5—C4	-178.09 (12)	N1—C5—C6—C1	-178.69 (11)
C7—N1—C5—C6	0.59 (13)	C4—C5—C6—N2	179.14 (10)
C5—N1—C7—N3	-178.29 (13)	N1—C5—C6—N2	0.31 (12)
C5—N1—C7—N2	-1.34 (14)	N1—C5—C4—C3	177.22 (12)
C6—N2—C7—N1	1.56 (14)	C6—C5—C4—C3	-1.35 (18)
C8—N2—C7—N1	176.05 (11)	C5—C4—C3—C2	1.6 (2)
C6—N2—C7—N3	178.56 (12)	C7—N2—C8—C9	-110.26 (14)
C8—N2—C7—N3	-7.0 (2)	C6—N2—C8—C9	63.27 (15)
C7—N2—C6—C1	177.81 (13)	C10—C9—C8—N2	-27 (3)
C8—N2—C6—C1	3.1 (2)	N2—C6—C1—C2	-177.87 (12)
C7—N2—C6—C5	-1.07 (12)	C5—C6—C1—C2	0.85 (18)
C8—N2—C6—C5	-175.79 (11)	C6—C1—C2—C3	-0.6 (2)
C4—C5—C6—C1	0.14 (18)	C4—C3—C2—C1	-0.6 (2)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N3—H1N3···N1 ⁱ	0.90 (2)	2.36 (2)	3.1823 (18)	153.5 (18)
N3—H2N3···N1 ⁱⁱ	0.860 (18)	2.241 (19)	3.0591 (15)	158.6 (15) (2)

Symmetry codes: (i) $-x+1, -y, -z+2$; (ii) $x, -y, z-1/2$.