

N-[2-(4-Bromobenzoyl)ethyl]isopropylaminium chloride

Abdullah Aydin,^{a*} Mehmet Akkurt,^b Halise Inci Gul,^c Ebru Mete^d and Ertan Sahin^d

^aDepartment of Science Education, Faculty of Education, Kastamonu University, 37200 Kastamonu, Turkey, ^bDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^cDepartment of Pharmaceutical Chemistry, Faculty of Pharmacy, Ataturk University, 25240 Erzurum, Turkey, and ^dDepartment of Chemistry, Faculty of Sciences, Ataturk University, 25240 Erzurum, Turkey
Correspondence e-mail: aaydin@kastamonu.edu.tr


Received 17 November 2011; accepted 6 December 2011

Key indicators: single-crystal X-ray study; $T = 294\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.009\text{ \AA}$; R factor = 0.076; wR factor = 0.181; data-to-parameter ratio = 18.5.

The crystal structure of the title compound, $\text{C}_{12}\text{H}_{17}\text{BrNO}^+ \cdot \text{Cl}^-$, is stabilized by $\text{N}-\text{H} \cdots \text{Cl}$ and $\text{C}-\text{H} \cdots \text{O}$ hydrogen bonds, forming a three-dimensional network. The interactions framework is completed by $\text{C}-\text{H} \cdots \pi$ contacts between a methylene group and the benzene ring of a symmetry-related molecule.

Related literature

For details of the pharmacological effects of Mannich bases and for the synthesis, see: Dimmock & Kumar (1997); Gul, Gul, *et al.* (2005); Gul, Sahin *et al.* (2005); Gul *et al.* (2007); Mete *et al.* (2011); Kucukoglu *et al.* (2011); Canturk *et al.* (2008); Chen *et al.* (1991); Suleyman *et al.* (2007). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{12}\text{H}_{17}\text{BrNO}^+ \cdot \text{Cl}^-$
 $M_r = 306.62$
Tetragonal, $P4_3/n$
 $a = 19.7122(4)\text{ \AA}$
 $c = 7.1738(2)\text{ \AA}$
 $V = 2787.53(11)\text{ \AA}^3$

Data collection

Rigaku R-Axis RAPID-S diffractometer
Absorption correction: multi-scan (Blessing, 1995)
 $T_{\min} = 0.632$, $T_{\max} = 0.709$

50060 measured reflections
2836 independent reflections
1617 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.151$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.076$
 $wR(F^2) = 0.181$
 $S = 1.07$
2836 reflections
153 parameters
2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.67\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.81\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$ is the centroid of the benzene ring.

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
N1—HN2···Cl1	0.86 (4)	2.26 (4)	3.102 (4)	166 (5)
N1—HN1···Cl1 ⁱ	0.86 (6)	2.27 (6)	3.133 (5)	177 (9)
Cl2—H12B···O1 ⁱⁱ	0.96	2.60	3.378 (7)	139
C9—H9B···Cg1 ⁱⁱⁱ	0.97	3.00	3.943 (6)	164

Symmetry codes: (i) $-y + \frac{1}{2}, x - 1, -z + \frac{3}{2}$; (ii) $-x + 1, -y, -z + 2$; (iii) $-x + 1, -y, -z + 1$.

Data collection: *CrystalClear* (Rigaku/MSC, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors are indebted to the Department of Chemistry, Ataturk University, Erzurum, Turkey, for use of the X-ray diffractometer purchased under grant No. 2003/219 of the University Research Fund.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2400).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Molterini, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Canturk, P., Kucukoglu, K., Topcu, Z., Gul, M. & Gul, H. I. (2008). *Arzneim. Forsch.* **58**, 686–691.
- Chen, H. T., Jing, Y. K., Ji, Z. Z. & Zhang, B. F. (1991). *Yaoxue Xuebao*, **26**, 183–192.
- Dimmock, J. R. & Kumar, P. (1997). *Curr. Med. Chem.* **4**, 1–22.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gul, M., Gul, H. I., Das, U. & Hanninen, O. (2005). *Arzneim. Forsch.* **55**, 332–337.
- Gul, H. I., Sahin, F., Gul, M., Ozturk, S. & Yerdelen, K. O. (2005). *Arch. Pharm.* **338**, 335–338.
- Gul, H. I., Yerdelen, K. O., Gul, M., Das, U., Pandit, B., Li, P.-K., Secen, H. & Sahin, F. (2007). *Arch. Pharm.* **340**, 195–201.
- Kucukoglu, K., Gul, M., Atalay, M., Mete, E., Kazaz, C., Hanninen, O. & Gul, H. I. (2011). *Arzneim. Forsch.* **61**, 366–371.
- Mete, E., Gul, H. I., Cetin-Atalay, R., Das, U., Sahin, E., Gul, M., Kazaz, C. & Dimmock, J. R. (2011). *Arch. Pharm. Chem. Life Sci.* **11**, 333–339.
- Rigaku/MSC (2005). *CrystalClear*. Rigaku/MSC, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Suleyman, H., Gul, H. I., Gul, M., Alkan, M. & Gocer, F. (2007). *Biol. Pharm. Bull.* **30**, 63–67.

supporting information

Acta Cryst. (2012). E68, o71 [doi:10.1107/S1600536811052640]

N-[2-(4-Bromobenzoyl)ethyl]isopropylaminium chloride

Abdullah Aydin, Mehmet Akkurt, Halise Inci Gul, Ebru Mete and Ertan Sahin

S1. Comment

Mannich bases are generally formed by the reaction between formaldehyde, a secondary amine and a compound containing reactive hydrogen atoms. On occasion, aldehydes other than formaldehyde may be employed and the secondary amine may be replaced by ammonia and primary amines. This process is known as the Mannich reaction (Dimmock & Kumar, 1997).

Mannich bases display varied biological activities such as antimicrobial (Gul, Sahin, *et al.*, 2005), cytotoxic (Gul, Gul *et al.*, 2005; Gul *et al.*, 2007; Mete *et al.*, 2011; Kucukoglu *et al.*, 2011), anticancer (Dimmock & Kumar, 1997; Chen *et al.*, 1991), anti-inflammatory (Suleyman *et al.*, 2007), and DNA topoisomerase I inhibiting properties (Canturk *et al.*, 2008).

A Mannich base having at least one hydrogen atom at the β position of amine group can undergo a deamination process to generate an α,β -unsaturated ketone moiety.

In the molecule of the title compound (Fig. 1), the bond lengths are within normal ranges (Allen *et al.*, 1987), as well as bond angles.

In the crystal structure, molecules are linked *via* intermolecular N—H \cdots Cl and C—H \cdots O hydrogen bonds (Table 1, Fig. 2), forming a three dimensional network. Furthermore, a C—H \cdots π interaction (Table 1) contributes to the stabilization of the crystal packing.

S2. Experimental

A mixture of the appropriate ketone (50 mmol), *para*-formaldehyde (50 mmol), and *iso*-propylamine hydrochloride (27 mmol) was heated in an oil bath at 403 K. The reaction vessel was then removed from the oil bath and when the temperature of the mixture dropped to 338 K, ethyl acetate (40–80 ml) was added. The mixture was stirred at room temperature for 24 h. and the resulting precipitates were then collected and the Mannich base (I) was passed through a column of silica gel 60 (70–230 mesh) using methanol as eluent. After evaporation of the solvent, the product was recrystallized from methanol. *M.p.*: 447–449 K. Yield: 38%. $^1\text{H-NMR}$ (CDCl_3) δ 1.49 (d, J = 6.8 Hz, 6H, $\text{CH}(\text{CH}_3)_2$), 3.34–3.38 (m, 3H, $\text{CH}(\text{CH}_3)_2$ and 2 x H-2), 3.73 (t, J = 7.3 Hz, 2H, 2 x H-3), 7.50 (d, J = 8.4 Hz, 2H, H-3'/5'), 7.76 (d, 2H, J = 8.4 Hz, H-2'/6'), 9.55 (brs, 2H, NH_2^+). $^{13}\text{C-NMR}$ (CDCl_3) δ 19.4 ($\text{CH}(\text{CH}_3)_2$), 35.3, 40.2, 51.3, 129.4, 129.8, 132.3, 134.7, 195.8 (CO); MS (EI) m/z : 254 ($M - \text{CH}_3$) $^+$, 256 ($M - \text{CH}_3 + 2$) $^+$, 270.2 ($M + \text{H}$) $^+$, 272.2 ($M + \text{H} + 2$) $^+$. IR (KBr, cm^{-1}): 2462 (NH_2^+), 1684 (CO). Calcd. for $\text{C}_{12}\text{H}_{17}\text{BrClNO}$ (306.63): C 47.00, H 5.59, N 4.57. Found: C 46.74, H 5.52, N 4.59 (Mete *et al.*, 2011).

S3. Refinement

The H atoms of the NH_2 group, HN1 and HN2, were located in a difference map and refined with a distance restraint of N—H = 0.86 (1) Å. Their displacement parameters were calculated as $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{N}1)$. The other H atoms were positioned geometrically with C—H = 0.93 (aromatic), 0.96 (methyl), 0.97 (methylene) and 0.98 Å (methine), and

constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{methyl C})$.





Figure 1

The title compound with displacement ellipsoids for non-H atoms shown at the 30% probability level.

**Figure 2**

The packing and hydrogen bonding of the title salt viewed down the *c* axis. H atoms not involved in hydrogen bonds are omitted for the sake of clarity.

N-[2-(4-Bromobenzoyl)ethyl]isopropylaminium chloride

Crystal data



$M_r = 306.62$

Tetragonal, $P4_2/n$

Hall symbol: -P 4bc

$a = 19.7122 (4)$ Å

$c = 7.1738 (2)$ Å

$V = 2787.53 (11)$ Å³

$Z = 8$

$F(000) = 1248$

$D_x = 1.461$ Mg m⁻³

Melting point: 447 K

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3888 reflections

$\theta = 2.9\text{--}26.4^\circ$

$\mu = 3.12$ mm⁻¹

$T = 294$ K

Block, white

$0.15 \times 0.13 \times 0.11$ mm

Data collection

Rigaku R-AXIS RAPID-S
diffractometer

Radiation source: Sealed Tube

Graphite Monochromator monochromator

Detector resolution: 10.0000 pixels mm⁻¹

dtprofit.ref scans

Absorption correction: multi-scan
(Blessing, 1995)

$T_{\min} = 0.632$, $T_{\max} = 0.709$

50060 measured reflections

2836 independent reflections
 1617 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.151$
 $\theta_{\text{max}} = 26.4^\circ, \theta_{\text{min}} = 2.9^\circ$

$h = -24 \rightarrow 24$
 $k = -24 \rightarrow 24$
 $l = -8 \rightarrow 8$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.076$
 $wR(F^2) = 0.181$
 $S = 1.07$
 2836 reflections
 153 parameters
 2 restraints
 0 constraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0581P)^2 + 4.3122P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.67 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.81 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.66757 (4)	0.19433 (5)	-0.04514 (15)	0.1156 (4)
O1	0.5235 (2)	0.0463 (2)	0.7192 (7)	0.0850 (17)
N1	0.6128 (2)	-0.1377 (2)	0.8767 (7)	0.0540 (17)
C1	0.6340 (3)	0.0359 (3)	0.3195 (9)	0.065 (2)
C2	0.6561 (3)	0.0740 (3)	0.1698 (10)	0.073 (3)
C3	0.6384 (3)	0.1411 (3)	0.1604 (10)	0.069 (2)
C4	0.5995 (3)	0.1707 (3)	0.2962 (10)	0.074 (3)
C5	0.5762 (3)	0.1323 (3)	0.4438 (9)	0.067 (2)
C6	0.5940 (3)	0.0644 (3)	0.4585 (8)	0.057 (2)
C7	0.5686 (3)	0.0245 (3)	0.6205 (9)	0.061 (2)
C8	0.5994 (3)	-0.0435 (3)	0.6592 (8)	0.063 (2)
C9	0.5742 (3)	-0.0744 (3)	0.8392 (9)	0.061 (2)
C10	0.5984 (3)	-0.1725 (3)	1.0610 (8)	0.0583 (19)
C11	0.6462 (3)	-0.2324 (3)	1.0768 (9)	0.070 (2)
C12	0.5251 (3)	-0.1928 (3)	1.0763 (8)	0.069 (2)
C11	0.75617 (8)	-0.07602 (8)	0.9322 (2)	0.0682 (6)
H1	0.64610	-0.00960	0.32750	0.0780*
HN1	0.604 (4)	-0.168 (3)	0.794 (9)	0.1390*
HN2	0.6548 (13)	-0.126 (4)	0.877 (12)	0.1390*
H2	0.68250	0.05440	0.07690	0.0870*
H4	0.58890	0.21660	0.28900	0.0890*
H5	0.54840	0.15200	0.53360	0.0800*
H8A	0.64830	-0.03880	0.66520	0.0760*
H8B	0.58880	-0.07390	0.55690	0.0760*
H9A	0.58050	-0.04270	0.94100	0.0730*
H9B	0.52610	-0.08460	0.82920	0.0730*
H10	0.60870	-0.14060	1.16200	0.0700*
H11A	0.63550	-0.26500	0.98190	0.1050*
H11B	0.64120	-0.25290	1.19730	0.1050*
H11C	0.69210	-0.21710	1.06120	0.1050*

H12A	0.51320	-0.22120	0.97260	0.1030*
H12B	0.49710	-0.15300	1.07580	0.1030*
H12C	0.51820	-0.21730	1.19040	0.1030*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0916 (6)	0.1139 (7)	0.1414 (9)	0.0018 (4)	0.0247 (5)	0.0597 (6)
O1	0.091 (3)	0.074 (3)	0.090 (3)	0.020 (2)	0.021 (3)	0.008 (3)
N1	0.055 (3)	0.049 (3)	0.058 (3)	0.002 (2)	-0.001 (2)	0.007 (2)
C1	0.061 (4)	0.055 (4)	0.080 (4)	0.009 (3)	-0.003 (3)	0.007 (3)
C2	0.067 (4)	0.067 (4)	0.084 (5)	0.003 (3)	0.006 (3)	0.011 (4)
C3	0.049 (3)	0.070 (4)	0.087 (5)	-0.006 (3)	-0.004 (3)	0.023 (4)
C4	0.075 (4)	0.056 (4)	0.090 (5)	0.002 (3)	-0.008 (4)	0.013 (4)
C5	0.072 (4)	0.051 (3)	0.077 (4)	0.005 (3)	-0.003 (3)	0.000 (3)
C6	0.062 (4)	0.050 (3)	0.060 (4)	-0.002 (3)	-0.010 (3)	0.004 (3)
C7	0.056 (4)	0.060 (4)	0.068 (4)	0.000 (3)	-0.004 (3)	-0.001 (3)
C8	0.064 (4)	0.061 (4)	0.065 (4)	0.006 (3)	0.001 (3)	0.009 (3)
C9	0.060 (4)	0.053 (3)	0.070 (4)	0.003 (3)	-0.003 (3)	0.005 (3)
C10	0.068 (4)	0.056 (3)	0.051 (3)	-0.003 (3)	-0.004 (3)	0.005 (3)
C11	0.070 (4)	0.072 (4)	0.067 (4)	0.009 (3)	-0.008 (3)	0.013 (3)
C12	0.067 (4)	0.075 (4)	0.064 (4)	-0.004 (3)	0.008 (3)	0.006 (3)
Cl1	0.0608 (9)	0.0857 (11)	0.0580 (9)	-0.0135 (7)	0.0045 (7)	-0.0045 (8)

Geometric parameters (\AA , $^\circ$)

Br1—C3	1.899 (7)	C10—C11	1.515 (8)
O1—C7	1.215 (7)	C1—H1	0.9300
N1—C9	1.486 (7)	C2—H2	0.9300
N1—C10	1.516 (8)	C4—H4	0.9300
N1—HN2	0.86 (4)	C5—H5	0.9300
N1—HN1	0.86 (6)	C8—H8A	0.9700
C1—C2	1.381 (9)	C8—H8B	0.9700
C1—C6	1.390 (9)	C9—H9A	0.9700
C2—C3	1.370 (8)	C9—H9B	0.9700
C3—C4	1.370 (9)	C10—H10	0.9800
C4—C5	1.380 (9)	C11—H11A	0.9600
C5—C6	1.388 (8)	C11—H11B	0.9600
C6—C7	1.490 (9)	C11—H11C	0.9600
C7—C8	1.498 (8)	C12—H12A	0.9600
C8—C9	1.512 (9)	C12—H12B	0.9600
C10—C12	1.503 (8)	C12—H12C	0.9600
C9—N1—C10	116.2 (4)	C5—C4—H4	120.00
HN2—N1—HN1	113 (8)	C4—C5—H5	120.00
C9—N1—HN1	111 (5)	C6—C5—H5	120.00
C10—N1—HN2	107 (6)	C7—C8—H8A	109.00
C9—N1—HN2	106 (5)	C7—C8—H8B	109.00

C10—N1—HN1	105 (4)	C9—C8—H8A	109.00
C2—C1—C6	121.1 (6)	C9—C8—H8B	109.00
C1—C2—C3	118.9 (6)	H8A—C8—H8B	108.00
Br1—C3—C2	119.7 (5)	N1—C9—H9A	110.00
Br1—C3—C4	119.1 (5)	N1—C9—H9B	110.00
C2—C3—C4	121.3 (6)	C8—C9—H9A	110.00
C3—C4—C5	119.9 (6)	C8—C9—H9B	110.00
C4—C5—C6	120.2 (6)	H9A—C9—H9B	108.00
C5—C6—C7	118.9 (5)	N1—C10—H10	108.00
C1—C6—C7	122.5 (5)	C11—C10—H10	108.00
C1—C6—C5	118.6 (5)	C12—C10—H10	108.00
O1—C7—C6	120.9 (5)	C10—C11—H11A	109.00
C6—C7—C8	118.7 (5)	C10—C11—H11B	109.00
O1—C7—C8	120.3 (6)	C10—C11—H11C	109.00
C7—C8—C9	112.7 (5)	H11A—C11—H11B	110.00
N1—C9—C8	109.0 (5)	H11A—C11—H11C	110.00
N1—C10—C12	111.4 (5)	H11B—C11—H11C	110.00
C11—C10—C12	112.6 (5)	C10—C12—H12A	109.00
N1—C10—C11	107.5 (5)	C10—C12—H12B	110.00
C2—C1—H1	119.00	C10—C12—H12C	109.00
C6—C1—H1	119.00	H12A—C12—H12B	109.00
C1—C2—H2	121.00	H12A—C12—H12C	109.00
C3—C2—H2	121.00	H12B—C12—H12C	109.00
C3—C4—H4	120.00		
C9—N1—C10—C11	-176.0 (5)	C3—C4—C5—C6	2.3 (9)
C9—N1—C10—C12	60.1 (6)	C4—C5—C6—C1	-1.7 (9)
C10—N1—C9—C8	174.9 (4)	C4—C5—C6—C7	179.4 (6)
C6—C1—C2—C3	0.5 (9)	C5—C6—C7—C8	-166.9 (5)
C2—C1—C6—C5	0.3 (9)	C1—C6—C7—O1	-165.4 (6)
C2—C1—C6—C7	179.2 (6)	C1—C6—C7—C8	14.2 (9)
C1—C2—C3—C4	0.1 (10)	C5—C6—C7—O1	13.5 (9)
C1—C2—C3—Br1	-179.5 (5)	O1—C7—C8—C9	-6.8 (8)
C2—C3—C4—C5	-1.4 (10)	C6—C7—C8—C9	173.6 (5)
Br1—C3—C4—C5	178.1 (5)	C7—C8—C9—N1	-173.8 (5)

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the benzene ring.

D—H···A	D—H	H···A	D···A	D—H···A
N1—HN2···C11	0.86 (4)	2.26 (4)	3.102 (4)	166 (5)
N1—HN1···C11 ⁱ	0.86 (6)	2.27 (6)	3.133 (5)	177 (9)
C12—H12B···O1 ⁱⁱ	0.96	2.60	3.378 (7)	139
C9—H9B···Cg1 ⁱⁱⁱ	0.97	3.00	3.943 (6)	164

Symmetry codes: (i) $-y+1/2, x-1, -z+3/2$; (ii) $-x+1, -y, -z+2$; (iii) $-x+1, -y, -z+1$.