

Ammonium diamminesilver(I) bis(5-chloro-2-hydroxybenzenesulfonate) trihydrate

 Zhao-Peng Deng,^a Shan Gao^a and Seik Weng Ng^{b,c*}

^aKey Laboratory of Functional Inorganic Material Chemistry, Ministry of Education, Heilongjiang University, Harbin 150080, People's Republic of China, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^cChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: seikweng@um.edu.my


Received 25 January 2012; accepted 26 January 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.032; wR factor = 0.084; data-to-parameter ratio = 16.9.

The reaction of silver nitrate with 5-chloro-2-hydroxybenzenesulfonic acid in the presence of ammonia yielded the title salt, $(\text{NH}_4)[\text{Ag}(\text{NH}_3)_2](\text{C}_6\text{H}_4\text{ClO}_4\text{S})_2 \cdot 3\text{H}_2\text{O}$. The Ag^{I} ion shows linear coordination [$\text{N}-\text{Ag}-\text{N} = 175.2$ (1)°]. The ammonium and diamminesilver cations, the benzenesulfonate anion and the lattice water molecules interact through an intricate network of $\text{N}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds to form a three-dimensional network.

Related literature

For a review of metal arenesulfonates, see: Cai (2004).



Experimental

Crystal data

$(\text{NH}_4)[\text{Ag}(\text{NH}_3)_2] \cdot (\text{C}_6\text{H}_4\text{ClO}_4\text{S})_2 \cdot 3\text{H}_2\text{O}$
 $M_r = 629.23$
 Orthorhombic, $P2_12_12_1$
 $a = 8.8814$ (8) Å
 $b = 9.8586$ (10) Å
 $c = 26.434$ (3) Å

$V = 2314.5$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.34$ mm⁻¹
 $T = 293$ K
 $0.19 \times 0.16 \times 0.13$ mm

Data collection

Rigaku R-Axis RAPID IP diffractometer

Absorption correction: multi-scan (ABSCOR; Higashi, 1995)
 $T_{\text{min}} = 0.785$, $T_{\text{max}} = 0.845$
 22666 measured reflections

5278 independent reflections
 4959 reflections with $I > 2\sigma(I)$

 $R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.084$
 $S = 1.04$
 5278 reflections
 312 parameters
 19 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.78$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.46$ e Å⁻³

Absolute structure: Flack (1983), with 2953 Friedel pairs
 Flack parameter: 0.02 (2)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O4}-\text{H4} \cdots \text{O3}$	0.84	2.38	2.970 (3)	127
$\text{O8}-\text{H8} \cdots \text{O4}$	0.84	2.04	2.630 (2)	127
$\text{O1w}-\text{H11} \cdots \text{O5}^{\text{i}}$	0.84 (1)	2.00 (1)	2.838 (3)	174 (4)
$\text{O1w}-\text{H12} \cdots \text{O3w}$	0.85 (1)	1.95 (1)	2.794 (3)	173 (4)
$\text{O2w}-\text{H21} \cdots \text{O3}$	0.83 (1)	2.45 (4)	2.974 (3)	122 (4)
$\text{O2w}-\text{H21} \cdots \text{O4}$	0.83 (1)	2.45 (3)	3.127 (3)	140 (5)
$\text{O2w}-\text{H22} \cdots \text{O5}^{\text{ii}}$	0.84 (1)	2.05 (2)	2.851 (3)	159 (5)
$\text{O3w}-\text{H31} \cdots \text{O2}^{\text{iii}}$	0.85 (1)	2.08 (1)	2.927 (3)	177 (5)
$\text{O3w}-\text{H32} \cdots \text{O7}^{\text{iv}}$	0.85 (1)	2.02 (2)	2.843 (3)	163 (5)
$\text{N1}-\text{H1a} \cdots \text{O5}$	0.88	2.32	3.11 (1)	148
$\text{N1}-\text{H1c} \cdots \text{O1}^{\text{v}}$	0.88	2.11	2.95 (1)	158
$\text{N2}-\text{H2a} \cdots \text{O1w}^{\text{vi}}$	0.88	2.30	3.15 (1)	163
$\text{N2}-\text{H2b} \cdots \text{O7}^{\text{vii}}$	0.88	2.25	3.07 (1)	153
$\text{N2}-\text{H2c} \cdots \text{O3}$	0.88	2.14	3.02 (1)	171
$\text{N3}-\text{H3a} \cdots \text{O3}^{\text{vi}}$	0.88 (1)	2.19 (1)	3.018 (3)	158 (3)
$\text{N3}-\text{H3b} \cdots \text{O6}^{\text{iv}}$	0.88 (1)	2.02 (1)	2.893 (3)	173 (3)
$\text{N3}-\text{H3c} \cdots \text{O8}$	0.88 (1)	1.99 (1)	2.820 (3)	159 (3)
$\text{N3}-\text{H3d} \cdots \text{O2w}$	0.88 (1)	1.94 (1)	2.826 (4)	177 (3)

Symmetry codes: (i) $x, y-1, z$; (ii) $-x+1, y-\frac{1}{2}, -z+\frac{3}{2}$; (iii) $x+1, y, z$; (iv) $-x+2, y-\frac{1}{2}, -z+\frac{3}{2}$; (v) $x, y+1, z$; (vi) $-x+1, y+\frac{1}{2}, -z+\frac{3}{2}$; (vii) $x-1, y, z$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MS, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

This work was supported by the Key Project of the Natural Science Foundation of Heilongjiang Province (grant No. ZD200903), the Key Project of the Education Bureau of Heilongjiang Province (grant Nos. 12511z023, 2011CJHB006), the Innovation Team of the Education Bureau of Heilongjiang Province (grant No. 2010 t d03), Heilongjiang University (grant No. Hdtd2010-04) and the Ministry of Higher Education of Malaysia (grant No. UM.C/HIR/MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5802).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Cai, J. (2004). *Coord. Chem. Rev.* **248**, 1061–1083.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
 Rigaku/MS (2002). *CrystalClear*. Rigaku/MS Inc., The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2012). E68, m225 [doi:10.1107/S160053681200339X]

Ammonium diamminesilver(I) bis(5-chloro-2-hydroxybenzenesulfonate) trihydrate

Zhao-Peng Deng, Shan Gao and Seik Weng Ng

S1. Comment

Metal arenesulfonates are commonly crystalline materials; their coordination chemistry has been reviewed (Cai, 2004). In this study, the attempt to synthesize the silver(I) derivative of 5-chloro-2-hydroxybenzenesulfonic acid in the presence of ammonia gave instead the ammine-coordinated salt (Scheme I) in which the diamminesilver cation interacts indirectly with the 2-hydrox-5-chlorobenzenesulfonate anion through the coordinated ammine ligands in an outer-sphere type of coordination. In the salt, $[\text{Ag}(\text{NH}_3)_2][\text{NH}_4](\text{C}_6\text{H}_4\text{ClO}_4\text{S})_2 \cdot 3\text{H}_2\text{O}$ (Fig. 1), the Ag^{II} atom shows linear coordination $[\text{N}-\text{Ag}-\text{N}$ 175.2 (1) °]. The ammonium and diamminesilver cations, the benzenesulfonate anion and the lattice water molecules interact through $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds to form a three-dimensional network (Table 1).

S2. Experimental


Silver nitrate (1 mmol) and 5-chloro-2-hydroxy-benzenesulfonic acid (1 mmol) were mixed in water (15 ml). The pH of the solution was adjusted to *ca* 6 by the addition of drops of ammonium hydroxide. The solution was filtered; colorless crystals were isolated after several days. The solution was shielded from light during the crystallization.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.93 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to $1.2U(\text{C})$. The hydroxy H atoms were assumed to be co-planar with the aromatic ring, and these were similarly constrained (O–H 0.84 Å) and their displacement factors were set to $1.5U_{\text{eq}}(\text{O})$. The amino H atoms were similarly constrained (N–H 0.88 Å) and their displacement factors were set to $1.5U_{\text{eq}}(\text{N})$.

The water H-atoms were located in a difference Fourier map, and were refined with distance restraints O–H 0.84 ± 0.01 Å and H \cdots H 1.37 ± 0.01 Å; their temperature factors were tied by a factor of 1.5 times.

The four ammonium H-atoms were located in a difference Fourier map, and were refined with distance restraints N–H 0.88 ± 0.01 Å and H \cdots H 1.43 ± 0.01 Å; their temperature factors were displacement factors were set to $1.5U_{\text{eq}}(\text{O})$.

**Figure 1**

Anisotropic displacement ellipsoid plot (Barbour, 2001) of $[\text{Ag}(\text{NH}_3)_2][\text{NH}_4](\text{C}_6\text{H}_4\text{ClO}_4\text{S})_2 \cdot 3\text{H}_2\text{O}$ at the 30% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

Ammonium diamminesilver(I) bis(5-chloro-2-hydroxybenzenesulfonate) trihydrate

Crystal data

$(\text{NH}_4)[\text{Ag}(\text{NH}_3)_2](\text{C}_6\text{H}_4\text{ClO}_4\text{S})_2 \cdot 3\text{H}_2\text{O}$

$M_r = 629.23$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 8.8814$ (8) Å

$b = 9.8586$ (10) Å

$c = 26.434$ (3) Å

$V = 2314.5$ (4) Å³

$Z = 4$

$F(000) = 1272$

$D_x = 1.806$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 16668 reflections

$\theta = 3.1\text{--}27.4^\circ$

$\mu = 1.34$ mm⁻¹

$T = 293$ K

Prism, colorless

$0.19 \times 0.16 \times 0.13$ mm

Data collection

Rigaku R-AXIS RAPID IP
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scan

Absorption correction: multi-scan

(*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.785$, $T_{\max} = 0.845$

22666 measured reflections

5278 independent reflections

4959 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.038$

$\theta_{\max} = 27.4^\circ$, $\theta_{\min} = 3.1^\circ$

$h = -11 \rightarrow 11$

$k = -12 \rightarrow 12$

$l = -32 \rightarrow 34$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.032$

$wR(F^2) = 0.084$

$S = 1.04$

5278 reflections

312 parameters

19 restraints

Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0556P)^2 + 0.1112P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.78 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.46 \text{ e } \text{\AA}^{-3}$
Absolute structure: Flack (1983), with 2953
Friedel pairs
Absolute structure parameter: 0.02 (2)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ag1	0.37725 (3)	0.63955 (3)	0.851526 (11)	0.06155 (10)
Cl1	0.26926 (8)	0.39184 (8)	1.06066 (2)	0.04433 (17)
Cl2	1.22944 (8)	0.75917 (9)	0.98804 (3)	0.04893 (18)
S1	0.30723 (6)	0.15274 (7)	0.87729 (2)	0.03079 (13)
S2	0.91696 (6)	0.76351 (6)	0.81204 (2)	0.02806 (12)
O1	0.4208 (2)	0.0468 (2)	0.87427 (9)	0.0498 (5)
O2	0.1652 (2)	0.1060 (2)	0.89819 (7)	0.0425 (5)
O3	0.2877 (2)	0.2202 (2)	0.82846 (7)	0.0422 (5)
O4	0.5744 (2)	0.3420 (2)	0.86422 (7)	0.0358 (4)
H4	0.5422	0.2865	0.8426	0.054*
O5	0.75295 (19)	0.7749 (2)	0.81090 (7)	0.0402 (4)
O6	0.9903 (2)	0.8954 (2)	0.81310 (8)	0.0414 (4)
O7	0.9735 (2)	0.6777 (2)	0.77132 (6)	0.0392 (5)
O8	0.8214 (2)	0.48134 (19)	0.84662 (7)	0.0335 (4)
H8	0.7913	0.4028	0.8537	0.050*
O1w	0.6740 (2)	0.0537 (2)	0.81430 (10)	0.0506 (5)
H11	0.692 (4)	-0.0301 (11)	0.8143 (19)	0.076*
H12	0.756 (3)	0.095 (3)	0.8202 (18)	0.076*
O2w	0.5127 (3)	0.2905 (3)	0.74936 (9)	0.0592 (6)
H21	0.516 (5)	0.262 (6)	0.7788 (8)	0.089*
H22	0.433 (3)	0.266 (6)	0.7351 (14)	0.089*
O3w	0.9314 (3)	0.2107 (3)	0.83062 (9)	0.0526 (6)
H31	0.998 (3)	0.177 (4)	0.8501 (11)	0.079*
H32	0.970 (4)	0.216 (5)	0.8012 (7)	0.079*
N1	0.5129 (3)	0.7659 (3)	0.89623 (12)	0.0617 (8)
H1A	0.6036	0.7708	0.8829	0.093*
H1B	0.5191	0.7323	0.9270	0.093*
H1C	0.4730	0.8475	0.8975	0.093*
N2	0.2539 (4)	0.5159 (4)	0.80167 (12)	0.0650 (8)
H2A	0.2936	0.5207	0.7712	0.098*
H2B	0.1599	0.5438	0.8006	0.098*
H2C	0.2567	0.4314	0.8123	0.098*
N3	0.7473 (3)	0.4833 (3)	0.74290 (9)	0.0386 (5)
H3A	0.714 (3)	0.5551 (19)	0.7267 (10)	0.058*
H3B	0.825 (2)	0.449 (3)	0.7270 (10)	0.058*
H3C	0.773 (3)	0.505 (3)	0.7738 (5)	0.058*
H3D	0.674 (2)	0.423 (2)	0.7437 (11)	0.058*

C1	0.3775 (2)	0.2748 (2)	0.92047 (8)	0.0273 (4)
C2	0.5039 (2)	0.3542 (3)	0.90939 (8)	0.0276 (4)
C3	0.5558 (3)	0.4449 (3)	0.94605 (10)	0.0328 (5)
H3	0.6390	0.4990	0.9390	0.039*
C4	0.4859 (3)	0.4562 (3)	0.99284 (10)	0.0342 (5)
H4A	0.5220	0.5163	1.0171	0.041*
C5	0.3610 (3)	0.3760 (3)	1.00269 (8)	0.0316 (5)
C6	0.3062 (3)	0.2859 (3)	0.96722 (9)	0.0310 (5)
H6	0.2223	0.2330	0.9745	0.037*
C7	0.9634 (2)	0.6794 (3)	0.86901 (8)	0.0261 (5)
C8	0.9076 (3)	0.5478 (3)	0.87867 (9)	0.0289 (5)
C9	0.9509 (3)	0.4891 (3)	0.92511 (10)	0.0381 (6)
H9	0.9140	0.4037	0.9335	0.046*
C10	1.0459 (3)	0.5539 (3)	0.95862 (10)	0.0393 (6)
H10	1.0720	0.5125	0.9890	0.047*
C11	1.1019 (3)	0.6808 (3)	0.94671 (9)	0.0347 (5)
C12	1.0603 (3)	0.7457 (3)	0.90287 (9)	0.0318 (5)
H12A	1.0960	0.8323	0.8957	0.038*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ag1	0.06878 (17)	0.05382 (16)	0.06205 (16)	0.01435 (13)	0.00984 (12)	0.00913 (13)
Cl1	0.0497 (3)	0.0586 (5)	0.0247 (2)	0.0016 (3)	0.0049 (3)	-0.0006 (3)
Cl2	0.0467 (3)	0.0593 (5)	0.0407 (3)	0.0043 (3)	-0.0153 (3)	-0.0148 (3)
S1	0.0335 (3)	0.0301 (3)	0.0288 (3)	-0.0049 (2)	0.0004 (2)	-0.0032 (2)
S2	0.0295 (3)	0.0293 (3)	0.0253 (2)	-0.0007 (2)	0.00047 (19)	0.0021 (2)
O1	0.0501 (11)	0.0354 (11)	0.0638 (13)	0.0047 (9)	-0.0003 (10)	-0.0095 (10)
O2	0.0385 (9)	0.0505 (13)	0.0386 (9)	-0.0150 (9)	0.0000 (8)	-0.0016 (9)
O3	0.0535 (10)	0.0453 (12)	0.0278 (8)	-0.0144 (9)	-0.0047 (8)	-0.0002 (8)
O4	0.0370 (8)	0.0407 (11)	0.0297 (8)	-0.0101 (8)	0.0057 (7)	-0.0058 (7)
O5	0.0310 (8)	0.0476 (12)	0.0420 (9)	0.0022 (8)	-0.0026 (7)	0.0069 (9)
O6	0.0496 (10)	0.0301 (10)	0.0446 (10)	-0.0075 (8)	-0.0031 (9)	0.0086 (8)
O7	0.0467 (10)	0.0459 (12)	0.0249 (8)	0.0035 (9)	0.0055 (7)	0.0006 (8)
O8	0.0386 (8)	0.0318 (9)	0.0303 (8)	-0.0108 (7)	0.0002 (7)	-0.0008 (7)
O1w	0.0506 (11)	0.0410 (12)	0.0604 (13)	0.0008 (10)	0.0040 (10)	-0.0035 (11)
O2w	0.0520 (12)	0.0760 (18)	0.0494 (11)	-0.0175 (13)	-0.0092 (10)	0.0148 (13)
O3w	0.0536 (12)	0.0635 (16)	0.0408 (10)	0.0040 (11)	0.0052 (9)	-0.0005 (11)
N1	0.0641 (16)	0.0554 (18)	0.0655 (17)	0.0117 (15)	0.0119 (14)	0.0201 (15)
N2	0.0724 (19)	0.065 (2)	0.0576 (17)	0.0251 (17)	0.0065 (15)	0.0041 (15)
N3	0.0414 (11)	0.0403 (13)	0.0340 (11)	-0.0020 (10)	-0.0015 (9)	0.0003 (9)
C1	0.0284 (9)	0.0276 (11)	0.0259 (10)	0.0015 (9)	-0.0031 (8)	-0.0021 (9)
C2	0.0291 (10)	0.0270 (11)	0.0267 (9)	0.0011 (9)	-0.0009 (8)	0.0007 (9)
C3	0.0321 (11)	0.0322 (13)	0.0341 (12)	-0.0035 (10)	-0.0004 (9)	-0.0022 (10)
C4	0.0390 (12)	0.0341 (14)	0.0294 (11)	-0.0021 (11)	-0.0030 (10)	-0.0033 (10)
C5	0.0348 (11)	0.0371 (14)	0.0227 (10)	0.0044 (11)	0.0015 (9)	0.0014 (9)
C6	0.0309 (10)	0.0345 (14)	0.0277 (11)	-0.0028 (10)	0.0001 (9)	0.0044 (10)
C7	0.0246 (9)	0.0320 (13)	0.0217 (9)	0.0024 (9)	0.0015 (8)	-0.0002 (8)

C8	0.0293 (10)	0.0313 (12)	0.0261 (10)	-0.0023 (9)	0.0042 (9)	-0.0004 (10)
C9	0.0445 (14)	0.0362 (14)	0.0336 (12)	-0.0007 (11)	-0.0002 (11)	0.0056 (11)
C10	0.0434 (13)	0.0457 (16)	0.0289 (11)	0.0098 (12)	-0.0050 (10)	0.0036 (11)
C11	0.0321 (11)	0.0445 (15)	0.0276 (11)	0.0024 (10)	-0.0047 (9)	-0.0094 (10)
C12	0.0297 (10)	0.0338 (13)	0.0318 (11)	0.0006 (10)	0.0003 (8)	-0.0046 (10)

Geometric parameters (Å, °)

Ag1—N1	2.097 (4)	N2—H2A	0.8800
Ag1—N2	2.103 (4)	N2—H2B	0.8800
Cl1—C5	1.743 (2)	N2—H2C	0.8800
Cl2—C11	1.753 (2)	N3—H3A	0.878 (9)
S1—O1	1.454 (2)	N3—H3B	0.877 (9)
S1—O2	1.4522 (19)	N3—H3C	0.875 (9)
S1—O3	1.462 (2)	N3—H3D	0.883 (9)
S1—C1	1.772 (2)	C1—C2	1.400 (3)
S2—O6	1.454 (2)	C1—C6	1.393 (3)
S2—O7	1.4580 (19)	C2—C3	1.397 (4)
S2—O5	1.4613 (18)	C3—C4	1.388 (4)
S2—C7	1.768 (2)	C3—H3	0.9300
O4—C2	1.353 (3)	C4—C5	1.387 (4)
O4—H4	0.8400	C4—H4A	0.9300
O8—C8	1.316 (3)	C5—C6	1.380 (4)
O8—H8	0.8400	C6—H6	0.9300
O1w—H11	0.841 (10)	C7—C12	1.403 (3)
O1w—H12	0.849 (10)	C7—C8	1.413 (4)
O2w—H21	0.828 (10)	C8—C9	1.410 (4)
O2w—H22	0.839 (10)	C9—C10	1.380 (4)
O3w—H31	0.849 (10)	C9—H9	0.9300
O3w—H32	0.852 (10)	C10—C11	1.382 (4)
N1—H1A	0.8800	C10—H10	0.9300
N1—H1B	0.8800	C11—C12	1.374 (4)
N1—H1C	0.8800	C12—H12A	0.9300
N1—Ag1—N2	175.18 (13)	C2—C1—C6	120.4 (2)
O1—S1—O2	113.33 (13)	C2—C1—S1	121.83 (17)
O1—S1—O3	111.12 (14)	C6—C1—S1	117.72 (18)
O2—S1—O3	112.13 (12)	O4—C2—C1	120.4 (2)
O1—S1—C1	106.19 (12)	O4—C2—C3	121.1 (2)
O2—S1—C1	106.03 (11)	C1—C2—C3	118.5 (2)
O3—S1—C1	107.56 (12)	C4—C3—C2	121.4 (2)
O6—S2—O7	112.24 (12)	C4—C3—H3	119.3
O6—S2—O5	112.22 (13)	C2—C3—H3	119.3
O7—S2—O5	111.88 (12)	C3—C4—C5	118.7 (2)
O6—S2—C7	107.34 (12)	C3—C4—H4A	120.7
O7—S2—C7	106.06 (11)	C5—C4—H4A	120.7
O5—S2—C7	106.62 (11)	C4—C5—C6	121.4 (2)
C2—O4—H4	120.0	C4—C5—C11	119.20 (19)

C8—O8—H8	120.0	C6—C5—C11	119.37 (19)
H11—O1w—H12	108 (2)	C5—C6—C1	119.6 (2)
H21—O2w—H22	111 (2)	C5—C6—H6	120.2
H31—O3w—H32	107 (2)	C1—C6—H6	120.2
Ag1—N1—H1A	109.5	C12—C7—C8	121.8 (2)
Ag1—N1—H1B	109.5	C12—C7—S2	117.93 (19)
H1A—N1—H1B	109.5	C8—C7—S2	120.21 (17)
Ag1—N1—H1C	109.5	O8—C8—C9	121.0 (2)
H1A—N1—H1C	109.5	O8—C8—C7	123.0 (2)
H1B—N1—H1C	109.5	C9—C8—C7	116.0 (2)
Ag1—N2—H2A	109.5	C10—C9—C8	122.4 (3)
Ag1—N2—H2B	109.5	C10—C9—H9	118.8
H2A—N2—H2B	109.5	C8—C9—H9	118.8
Ag1—N2—H2C	109.5	C9—C10—C11	119.5 (2)
H2A—N2—H2C	109.5	C9—C10—H10	120.2
H2B—N2—H2C	109.5	C11—C10—H10	120.2
H3A—N3—H3B	110 (1)	C12—C11—C10	121.1 (2)
H3A—N3—H3C	110 (1)	C12—C11—C12	119.6 (2)
H3B—N3—H3C	110 (1)	C10—C11—C12	119.3 (2)
H3A—N3—H3D	108 (1)	C11—C12—C7	119.1 (2)
H3B—N3—H3D	110 (1)	C11—C12—H12A	120.5
H3C—N3—H3D	110 (1)	C7—C12—H12A	120.5
O1—S1—C1—C2	67.4 (2)	O6—S2—C7—C12	-2.1 (2)
O2—S1—C1—C2	-171.8 (2)	O7—S2—C7—C12	118.10 (19)
O3—S1—C1—C2	-51.6 (2)	O5—S2—C7—C12	-122.51 (19)
O1—S1—C1—C6	-110.0 (2)	O6—S2—C7—C8	-179.93 (18)
O2—S1—C1—C6	10.8 (2)	O7—S2—C7—C8	-59.8 (2)
O3—S1—C1—C6	130.9 (2)	O5—S2—C7—C8	59.6 (2)
C6—C1—C2—O4	179.1 (2)	C12—C7—C8—O8	-177.2 (2)
S1—C1—C2—O4	1.7 (3)	S2—C7—C8—O8	0.6 (3)
C6—C1—C2—C3	-0.4 (4)	C12—C7—C8—C9	2.2 (3)
S1—C1—C2—C3	-177.78 (19)	S2—C7—C8—C9	179.98 (18)
O4—C2—C3—C4	-178.8 (2)	O8—C8—C9—C10	177.4 (2)
C1—C2—C3—C4	0.8 (4)	C7—C8—C9—C10	-2.0 (4)
C2—C3—C4—C5	-0.7 (4)	C8—C9—C10—C11	-0.2 (4)
C3—C4—C5—C6	0.2 (4)	C9—C10—C11—C12	2.3 (4)
C3—C4—C5—C11	-178.5 (2)	C9—C10—C11—C12	-177.0 (2)
C4—C5—C6—C1	0.1 (4)	C10—C11—C12—C7	-2.2 (4)
C11—C5—C6—C1	178.78 (19)	C12—C11—C12—C7	177.22 (18)
C2—C1—C6—C5	0.0 (4)	C8—C7—C12—C11	-0.2 (3)
S1—C1—C6—C5	177.48 (19)	S2—C7—C12—C11	-178.01 (18)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O4—H4 \cdots O3	0.84	2.38	2.970 (3)	127
O8—H8 \cdots O4	0.84	2.04	2.630 (2)	127

O1 _w —H11···O5 ⁱ	0.84 (1)	2.00 (1)	2.838 (3)	174 (4)
O1 _w —H12···O3 _w	0.85 (1)	1.95 (1)	2.794 (3)	173 (4)
O2 _w —H21···O3	0.83 (1)	2.45 (4)	2.974 (3)	122 (4)
O2 _w —H21···O4	0.83 (1)	2.45 (3)	3.127 (3)	140 (5)
O2 _w —H22···O5 ⁱⁱ	0.84 (1)	2.05 (2)	2.851 (3)	159 (5)
O3 _w —H31···O2 ⁱⁱⁱ	0.85 (1)	2.08 (1)	2.927 (3)	177 (5)
O3 _w —H32···O7 ^{iv}	0.85 (1)	2.02 (2)	2.843 (3)	163 (5)
N1—H1a···O5	0.88	2.32	3.11 (1)	148
N1—H1c···O1 ^v	0.88	2.11	2.95 (1)	158
N2—H2a···O1 _w ^{vi}	0.88	2.30	3.15 (1)	163
N2—H2b···O7 ^{vii}	0.88	2.25	3.07 (1)	153
N2—H2c···O3	0.88	2.14	3.02 (1)	171
N3—H3a···O3 ^{vi}	0.88 (1)	2.19 (1)	3.018 (3)	158 (3)
N3—H3b···O6 ^{iv}	0.88 (1)	2.02 (1)	2.893 (3)	173 (3)
N3—H3c···O8	0.88 (1)	1.99 (1)	2.820 (3)	159 (3)
N3—H3d···O2 _w	0.88 (1)	1.94 (1)	2.826 (4)	177 (3)

Symmetry codes: (i) $x, y-1, z$; (ii) $-x+1, y-1/2, -z+3/2$; (iii) $x+1, y, z$; (iv) $-x+2, y-1/2, -z+3/2$; (v) $x, y+1, z$; (vi) $-x+1, y+1/2, -z+3/2$; (vii) $x-1, y, z$.