

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Hydroxy-*N'*-(4-hydroxybenzylidene)-3-methylbenzohydrazide

Xi-Hai Shen, Li-Xue Zhu,* Li-Juen Shao and Zhao-Fu Zhu

 Department of Chemistry, Hebei Normal University of Science and Technology, Qinhuangdao 066600, People's Republic of China
 Correspondence e-mail: zhaofu_zhu@163.com


Received 26 December 2011; accepted 27 December 2011

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.044; wR factor = 0.103; data-to-parameter ratio = 14.5.

The title compound, $\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_3$, was prepared by condensing 4-hydroxybenzaldehyde and 2-hydroxy-3-methylbenzohydrazide in methanol. The two benzene rings make a dihedral angle of 19.03 (11)°. An intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond is observed. The crystal structure is stabilized by intermolecular $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds and $\text{C}-\text{H}\cdots\text{O}$ interactions, which lead to the formation of a three-dimensional network.

Related literature

For the crystal structures of similar hydrazone compounds, see: Fun *et al.* (2011); Horkaew *et al.* (2011); Zhi *et al.* (2011); Huang & Wu (2010).



Experimental

Crystal data

 $\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_3$
 $M_r = 270.28$

 Orthorhombic, $P2_12_12_1$
 $a = 7.3872$ (17) Å

 $b = 13.012$ (2) Å
 $c = 13.592$ (2) Å
 $V = 1306.5$ (4) Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 298$ K
 $0.20 \times 0.20 \times 0.17$ mm

Data collection

 Bruker SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\min} = 0.981$, $T_{\max} = 0.984$

 6296 measured reflections
 2663 independent reflections
 1953 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.103$
 $S = 0.99$
 2663 reflections

 184 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.13$ e Å⁻³
 $\Delta\rho_{\min} = -0.19$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O3}-\text{H3}\cdots\text{O2}^i$	0.82	1.86	2.575 (2)	146
$\text{O1}-\text{H1}\cdots\text{O2}^i$	0.82	2.00	2.809 (2)	167
$\text{N2}-\text{H2A}\cdots\text{O1}^{ii}$	0.86	2.36	3.067 (2)	139
$\text{C3}-\text{H3A}\cdots\text{O2}^i$	0.93	2.51	3.208 (2)	132

 Symmetry codes: (i) $-x, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + \frac{1}{2}, -y + 2, z - \frac{1}{2}$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2356).

References

- Bruker (2001). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Fun, H.-K., Horkaew, J. & Chantrapromma, S. (2011). *Acta Cryst.* **E67**, o2644–o2645.
 Horkaew, J., Chantrapromma, S. & Fun, H.-K. (2011). *Acta Cryst.* **E67**, o2985.
 Huang, H.-T. & Wu, H.-Y. (2010). *Acta Cryst.* **E66**, o2729–o2730.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Zhi, F., Wang, R., Zhang, Y., Wang, Q. & Yang, Y.-L. (2011). *Acta Cryst.* **E67**, o2825.

supporting information

Acta Cryst. (2012). E68, o297 [doi:10.1107/S1600536811055930]

2-Hydroxy-*N'*-(4-hydroxybenzylidene)-3-methylbenzohydrazide

Xi-Hai Shen, Li-Xue Zhu, Li-Juen Shao and Zhao-Fu Zhu

S1. Comment

In the last few years, the crystal structures of a number of hydrazone compounds have been reported (Fun *et al.*, 2011; Horkaew *et al.*, 2011; Zhi *et al.*, 2011; Huang & Wu, 2010). As an extension of work on such compounds, we report herein on the synthesis and crystal structure of the title compound.

In the title molecule, Fig. 1, there is an intramolecular O3—H3···O2 hydrogen bond (Table 1). The benzene rings, (C1—C6) and (C9—C14), make a dihedral angle of 19.03 (11)°. All the geometrical parameters are within normal ranges and are comparable with those in similar compounds, mentioned above.

In the crystal, molecules are linked via O—H···O and N—H···O hydrogen bonds and C—H···O interactions, leading to the formation of a three-dimensional network (Table 1 and Fig. 2).

S2. Experimental

4-Hydroxybenzaldehyde (122.1 mg, 1.0 mmol) and 2-hydroxy-3-methylbenzohydrazide (166.2 mg, 1.0 mmol) were mixed in methanol (60 ml). The mixture was refluxed for 30 min, then cooled to room temperature, yielding a colourless solution. Colourless crystals were formed when the solution was left to evaporate in air for several days.

S3. Refinement

All the H atoms were placed in calculated positions and refined as riding atoms: O—H = 0.82 Å, N—H = 0.86 Å, C—H = 0.93 and 0.96 Å for CH and CH₃ H atoms, respectively, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{O,N,C})$, where $k = 1.5$ for OH and CH₃ H-atoms and $k = 1.2$ for all other H-atoms. In the absence of significant anomalous scattering effects the Flack parameter of 2.5 (15) for 1092 Friedel pairs, has no meaning.





Figure 1

The molecular structure of the title molecule, with atom numbering and displacement ellipsoids drawn at the 30% probability level. The intramolecular O—H···O hydrogen bond is drawn as a dashed line.

**Figure 2**

The crystal packing of the title compound, viewed along the *a* axis. The O-H...O and N-H...O hydrogen bonds are drawn as dashed lines.

2-Hydroxy-*N'*-(4-hydroxybenzylidene)-3-methylbenzohydrazide

Crystal data

$C_{15}H_{14}N_2O_3$

$M_r = 270.28$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 7.3872$ (17) Å

$b = 13.012$ (2) Å

$c = 13.592$ (2) Å

$V = 1306.5$ (4) Å³

$Z = 4$

$F(000) = 568$

$D_x = 1.374$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1375 reflections

$\theta = 2.6$ – 24.5°

$\mu = 0.10$ mm⁻¹

$T = 298$ K

Block, colourless

$0.20 \times 0.20 \times 0.17$ mm

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2001)

$T_{\min} = 0.981$, $T_{\max} = 0.984$

6296 measured reflections

2663 independent reflections

1953 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.032$

$\theta_{\max} = 26.5^\circ$, $\theta_{\min} = 3.0^\circ$

$h = -9 \rightarrow 8$

$k = -13 \rightarrow 16$

$l = -17 \rightarrow 15$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.103$
 $S = 0.99$
 2663 reflections
 184 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0494P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.13 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.0995 (3)	0.79013 (12)	0.51891 (12)	0.0453 (5)
N2	0.1159 (3)	0.72214 (12)	0.44083 (12)	0.0455 (5)
H2A	0.1404	0.7437	0.3825	0.055*
O1	0.1126 (2)	1.18743 (10)	0.79368 (10)	0.0516 (4)
H1	0.0544	1.1672	0.8412	0.077*
O2	0.0499 (2)	0.59172 (10)	0.54350 (10)	0.0523 (5)
O3	0.1334 (3)	0.40715 (11)	0.49251 (11)	0.0600 (5)
H3	0.1016	0.4528	0.5303	0.090*
C1	0.1268 (3)	0.96163 (14)	0.57681 (14)	0.0376 (5)
C2	0.0866 (3)	0.93585 (16)	0.67358 (15)	0.0430 (6)
H2	0.0608	0.8679	0.6894	0.052*
C3	0.0846 (3)	1.00949 (16)	0.74632 (14)	0.0433 (6)
H3A	0.0602	0.9908	0.8110	0.052*
C4	0.1188 (3)	1.11133 (14)	0.72333 (15)	0.0395 (5)
C5	0.1604 (3)	1.13823 (17)	0.62779 (15)	0.0479 (6)
H5	0.1856	1.2063	0.6121	0.057*
C6	0.1646 (3)	1.06390 (16)	0.55602 (16)	0.0457 (6)
H6	0.1935	1.0826	0.4919	0.055*
C7	0.1324 (3)	0.88408 (15)	0.49972 (15)	0.0424 (5)
H7	0.1604	0.9034	0.4356	0.051*
C8	0.0922 (3)	0.62132 (15)	0.45912 (15)	0.0410 (5)
C9	0.1136 (3)	0.54807 (15)	0.37750 (14)	0.0393 (5)
C10	0.1344 (3)	0.44378 (16)	0.39901 (15)	0.0437 (5)
C11	0.1573 (3)	0.37050 (17)	0.32441 (17)	0.0487 (6)
C12	0.1510 (3)	0.40437 (19)	0.22839 (17)	0.0541 (6)

H12	0.1634	0.3569	0.1777	0.065*
C13	0.1266 (3)	0.50748 (19)	0.20530 (17)	0.0537 (6)
H13	0.1215	0.5281	0.1399	0.064*
C14	0.1101 (3)	0.57847 (17)	0.27877 (15)	0.0465 (5)
H14	0.0964	0.6476	0.2630	0.056*
C15	0.1874 (4)	0.26079 (17)	0.35184 (19)	0.0692 (8)
H15A	0.2038	0.2206	0.2933	0.104*
H15B	0.0842	0.2357	0.3875	0.104*
H15C	0.2934	0.2555	0.3924	0.104*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0612 (13)	0.0413 (10)	0.0334 (10)	0.0007 (9)	0.0050 (9)	-0.0060 (7)
N2	0.0655 (13)	0.0413 (10)	0.0297 (9)	0.0004 (9)	0.0056 (10)	-0.0025 (8)
O1	0.0721 (12)	0.0415 (8)	0.0411 (8)	-0.0051 (8)	0.0055 (8)	-0.0065 (7)
O2	0.0801 (12)	0.0455 (9)	0.0312 (8)	-0.0016 (8)	0.0081 (8)	0.0011 (7)
O3	0.0948 (13)	0.0435 (9)	0.0416 (9)	0.0010 (10)	0.0016 (10)	0.0033 (7)
C1	0.0427 (12)	0.0367 (11)	0.0335 (11)	0.0039 (10)	-0.0002 (10)	0.0008 (9)
C2	0.0518 (14)	0.0346 (11)	0.0427 (12)	-0.0004 (10)	0.0029 (11)	0.0039 (10)
C3	0.0576 (16)	0.0410 (12)	0.0313 (11)	0.0001 (10)	0.0038 (11)	0.0010 (9)
C4	0.0436 (13)	0.0366 (11)	0.0383 (12)	0.0026 (10)	0.0018 (11)	-0.0053 (9)
C5	0.0598 (15)	0.0362 (11)	0.0475 (14)	-0.0068 (10)	0.0091 (12)	0.0031 (10)
C6	0.0579 (15)	0.0453 (13)	0.0339 (11)	0.0019 (11)	0.0046 (12)	0.0053 (10)
C7	0.0469 (13)	0.0434 (13)	0.0368 (12)	0.0037 (11)	0.0051 (11)	-0.0005 (10)
C8	0.0458 (13)	0.0425 (12)	0.0347 (12)	-0.0010 (10)	-0.0004 (11)	0.0019 (9)
C9	0.0429 (12)	0.0411 (11)	0.0340 (11)	-0.0024 (10)	-0.0009 (11)	-0.0042 (9)
C10	0.0491 (14)	0.0453 (12)	0.0365 (12)	-0.0060 (11)	-0.0005 (11)	-0.0023 (10)
C11	0.0458 (14)	0.0508 (14)	0.0495 (14)	-0.0061 (11)	0.0031 (11)	-0.0102 (11)
C12	0.0519 (15)	0.0624 (16)	0.0481 (14)	-0.0065 (12)	0.0039 (12)	-0.0237 (12)
C13	0.0602 (16)	0.0687 (16)	0.0322 (12)	-0.0023 (14)	0.0010 (12)	-0.0033 (11)
C14	0.0520 (14)	0.0515 (13)	0.0360 (12)	0.0003 (11)	-0.0003 (11)	0.0011 (10)
C15	0.083 (2)	0.0445 (14)	0.0800 (19)	-0.0051 (13)	0.0022 (16)	-0.0157 (13)

Geometric parameters (Å, °)

N1—C7	1.273 (2)	C5—H5	0.9300
N1—N2	1.387 (2)	C6—H6	0.9300
N2—C8	1.347 (2)	C7—H7	0.9300
N2—H2A	0.8600	C8—C9	1.471 (3)
O1—C4	1.377 (2)	C9—C10	1.397 (3)
O1—H1	0.8200	C9—C14	1.399 (3)
O2—C8	1.249 (2)	C10—C11	1.402 (3)
O3—C10	1.357 (2)	C11—C12	1.378 (3)
O3—H3	0.8200	C11—C15	1.492 (3)
C1—C6	1.389 (3)	C12—C13	1.390 (3)
C1—C2	1.389 (3)	C12—H12	0.9300
C1—C7	1.455 (3)	C13—C14	1.366 (3)

C2—C3	1.377 (3)	C13—H13	0.9300
C2—H2	0.9300	C14—H14	0.9300
C3—C4	1.385 (3)	C15—H15A	0.9600
C3—H3A	0.9300	C15—H15B	0.9600
C4—C5	1.380 (3)	C15—H15C	0.9600
C5—C6	1.374 (3)		
C7—N1—N2	116.04 (18)	O2—C8—N2	120.15 (18)
C8—N2—N1	117.96 (17)	O2—C8—C9	121.28 (18)
C8—N2—H2A	121.0	N2—C8—C9	118.56 (18)
N1—N2—H2A	121.0	C10—C9—C14	118.51 (19)
C4—O1—H1	109.5	C10—C9—C8	118.92 (18)
C10—O3—H3	109.5	C14—C9—C8	122.55 (18)
C6—C1—C2	117.84 (18)	O3—C10—C9	122.45 (19)
C6—C1—C7	120.81 (18)	O3—C10—C11	116.04 (19)
C2—C1—C7	121.34 (18)	C9—C10—C11	121.5 (2)
C3—C2—C1	120.93 (18)	C12—C11—C10	117.6 (2)
C3—C2—H2	119.5	C12—C11—C15	123.2 (2)
C1—C2—H2	119.5	C10—C11—C15	119.2 (2)
C2—C3—C4	120.13 (19)	C11—C12—C13	121.8 (2)
C2—C3—H3A	119.9	C11—C12—H12	119.1
C4—C3—H3A	119.9	C13—C12—H12	119.1
O1—C4—C5	118.58 (18)	C14—C13—C12	120.0 (2)
O1—C4—C3	121.69 (18)	C14—C13—H13	120.0
C5—C4—C3	119.73 (18)	C12—C13—H13	120.0
C6—C5—C4	119.64 (19)	C13—C14—C9	120.6 (2)
C6—C5—H5	120.2	C13—C14—H14	119.7
C4—C5—H5	120.2	C9—C14—H14	119.7
C5—C6—C1	121.70 (19)	C11—C15—H15A	109.5
C5—C6—H6	119.1	C11—C15—H15B	109.5
C1—C6—H6	119.1	H15A—C15—H15B	109.5
N1—C7—C1	120.85 (19)	C11—C15—H15C	109.5
N1—C7—H7	119.6	H15A—C15—H15C	109.5
C1—C7—H7	119.6	H15B—C15—H15C	109.5

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O3—H3 \cdots O2	0.82	1.86	2.575 (2)	146
O1—H1 \cdots O2 ⁱ	0.82	2.00	2.809 (2)	167
N2—H2A \cdots O1 ⁱⁱ	0.86	2.36	3.067 (2)	139
C3—H3A \cdots O2 ⁱ	0.93	2.51	3.208 (2)	132

Symmetry codes: (i) $-x, y+1/2, -z+3/2$; (ii) $-x+1/2, -y+2, z-1/2$.