

(E)-1-(2-Hydroxy-5-methoxybenzylidene)thiosemicarbazide

Amir Adabi Ardakani,^{a*} Hadi Kargar,^b Reza Kia^{c,d} and Muhammad Nawaz Tahir^{e*}

^aArdakan Branch, Islamic Azad University, Ardakan, Iran, ^bDepartment of Chemistry, Payame Noor University, PO BOX 19395-3697 Tehran, I. R. of IRAN, ^cX-ray Crystallography Lab., Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Tehran, IRAN, ^dDepartment of Chemistry, Science and Research Branch, Islamic Azad University, Tehran, Iran, and ^eDepartment of Physics, University of Sargodha, Punjab, Pakistan
Correspondence e-mail: A.Adabi@iauardakan.ac.ir, dmntahir_uos@yahoo.com


Received 28 December 2011; accepted 29 December 2011

Key indicators: single-crystal X-ray study; $T = 291\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.035; wR factor = 0.110; data-to-parameter ratio = 19.4.

In the title molecule, $\text{C}_9\text{H}_{11}\text{N}_3\text{O}_2\text{S}$, an intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bond generates an $S(6)$ ring motif. In the crystal, molecules are linked via pairs of $\text{N}-\text{H}\cdots\text{S}$ interactions, forming inversion dimers with $R_2^2(8)$ ring motifs. These dimers are further linked via $\text{N}-\text{H}\cdots\text{S}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a two-dimensional network lying parallel to (100). The crystal structure is further stabilized by intermolecular $\pi-\pi$ interactions [centroid–centroid distance = 3.7972 (9) \AA].

Related literature

For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For background to thiosemicarbazones in coordination chemistry, see: Casas *et al.* (2000). For their biological applications, see: Maccioni *et al.* (2003); Ferrari *et al.* (2000). For related structures, see: Kargar *et al.* (2010a,b); Adabi Ardakani *et al.* (2012).



Experimental

Crystal data

$\text{C}_9\text{H}_{11}\text{N}_3\text{O}_2\text{S}$	$V = 1074.55 (4)\text{ \AA}^3$
$M_r = 225.27$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 7.4878 (2)\text{ \AA}$	$\mu = 0.29\text{ mm}^{-1}$
$b = 9.9880 (2)\text{ \AA}$	$T = 291\text{ K}$
$c = 14.3754 (3)\text{ \AA}$	$0.24 \times 0.14 \times 0.08\text{ mm}$
$\beta = 91.846 (1)^\circ$	

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	10176 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	2673 independent reflections
$T_{\min} = 0.800$, $T_{\max} = 0.926$	2365 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.016$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$	138 parameters
$wR(F^2) = 0.110$	H-atom parameters constrained
$S = 1.07$	$\Delta\rho_{\text{max}} = 0.31\text{ e \AA}^{-3}$
2673 reflections	$\Delta\rho_{\text{min}} = -0.24\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots N1	0.82	1.97	2.6844 (15)	146
N2—H2 \cdots Si ⁱ	0.86	2.61	3.3706 (12)	148
N3—H3A \cdots Si ⁱⁱ	0.86	2.66	3.2706 (12)	129
N3—H3B \cdots O1 ⁱⁱⁱ	0.86	2.11	2.9604 (17)	172

Symmetry codes: (i) $-x, -y, -z + 2$; (ii) $-x, y + \frac{1}{2}, -z + \frac{5}{2}$; (iii) $-x, y - \frac{1}{2}, -z + \frac{5}{2}$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

AAA thanks the Islamic Azad University, Ardakan Branch (this paper was extracted from the research project). HK thanks PNU for financial support. MNT thanks Sargodha University for research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2361).

References

- Adabi Ardakani, A., Kargar, H., Kia, R. & Tahir, M. N. (2012). *Acta Cryst. E68*, o340–o341.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Casas, J. S., Garcia-Tasende, M. S. & Sordo, J. (2000). *Coord. Chem. Rev.* **209**, 197–261.
- Ferrari, M. B., Capacchi, S., Reffo, G., Pelosi, G., Tarasconi, P., Albertini, R., Pinelli, S. & Lunghi, P. (2000). *J. Inorg. Biochem.* **81**, 89–97.
- Kargar, H., Kia, R., Akkurt, M. & Büyükgüngör, O. (2010a). *Acta Cryst. E66*, o2999.

- Kargar, H., Kia, R., Akkurt, M. & Büyükgüngör, O. (2010b). *Acta Cryst. E*66, o2981.
- Maccioni, E., Cardia, M. C., Distinto, S., Bonsignore, L. & De Logu, A. (2003). *Farmacol* 58, 951–959.
- Sheldrick, G. M. (2008). *Acta Cryst. A*64, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D*65, 148–155.

supporting information

Acta Cryst. (2012). E68, o324–o325 [doi:10.1107/S1600536811056182]

(E)-1-(2-Hydroxy-5-methoxybenzylidene)thiosemicarbazide

Amir Adabi Ardakani, Hadi Kargar, Reza Kia and Muhammad Nawaz Tahir

S1. Comment

Thiosemicarbazones constitute an important class of N,S donor ligands due to their propensity to react with a wide range of metals (Casas *et al.*, 2000). Thiosemicarbazones exhibit various biological activities and have therefore attracted considerable pharmaceutical interest (Maccioni *et al.*, 2003; Ferrari *et al.*, 2000). We report herein on the synthesis and crystal structure of the title a hydrazone Schiff base compound.

The asymmetric unit of the title compound, Fig. 1, comprises a hydrazone Schiff base ligand. The bond lengths and angles are within the normal ranges and are comparable to those reported for related structures (Kargar *et al.*, 2010*a,b*; Adabi *et al.*, 2012). An intramolecular O—H···N hydrogen bond (Table 1) generates an S(6) ring motif (Bernstein *et al.*, 1995).


In the crystal, pairs of intermolecular N—H···S hydrogen bonds make inversion dimers with an R₂(8) ring motif. Intermolecular N—H···S and N—H···O hydrogen bonds link neighbouring molecules into a two-dimensional extended network parallel to (1 0 0). For details of the hydrogen bonding see Table 1, and Fig. 2. The crystal structure is further stabilized by an intermolecular $\pi\cdots\pi$ interaction [$Cg1\cdots Cg1^i = 3.7972(9)\text{\AA}$; (i) $-x+1, -y+1, -z+2$; $Cg1$ is the centroid of ring (C1–C6)].

S2. Experimental


A mixture of 5-methoxysalicylaldehyde (0.01 mol) and hydrazinecarbothioamide (0.01 mol) in 20 ml of ethanol was refluxed for about 2 h. On cooling, the solid separated was filtered and recrystallized from ethanol. Colourless plate-like crystals of the title compound, suitable for X-ray diffraction, were obtained by slow evaporation of a solution in ethanol.

S3. Refinement

O- and N-bound H atoms were located in a difference Fourier map and were constrained to ride on their parent atoms: O—H = 0.82 Å, N—H = 0.86 Å, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ and $1.2U_{\text{eq}}(\text{N})$. The C-bound H-atoms were included in calculated positions and treated as riding atoms: C—H = 0.93 and 0.96 Å for CH and CH₃ H-atoms, respectively, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C})$, where $k = 1.5$ for CH₃ H-atoms, and $k = 1.2$ for all other H-atoms.

**Figure 1**

The molecular structure of the title molecule, showing 40% probability displacement ellipsoids and the atomic numbering. The intramolecular O-H \cdots N hydrogen bond is drawn as a dashed line.

**Figure 2**

The crystal packing of the title compound, viewed along the a -axis, showing the two-dimensional extended network parallel to $(1\ 0\ 0)$. The $\text{N}-\text{H}\cdots\text{S}$ and $\text{N}-\text{H}\cdots\text{O}$, hydrogen bonds are shown as dashed lines [see Table 1 for details; only the H atoms involved in these interactions are shown].

(E)-1-(2-Hydroxy-5-methoxybenzylidene)thiosemicarbazide

Crystal data

$\text{C}_9\text{H}_{11}\text{N}_3\text{O}_2\text{S}$
 $M_r = 225.27$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 7.4878 (2)$ Å
 $b = 9.9880 (2)$ Å
 $c = 14.3754 (3)$ Å
 $\beta = 91.846 (1)^\circ$
 $V = 1074.55 (4)$ Å³
 $Z = 4$

$F(000) = 472$
 $D_x = 1.392 \text{ Mg m}^{-3}$
 $\text{Mo } K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 2750 reflections
 $\theta = 2.4\text{--}27.5^\circ$
 $\mu = 0.29 \text{ mm}^{-1}$
 $T = 291 \text{ K}$
Plate, colourless
 $0.24 \times 0.14 \times 0.08$ mm

Data collection

Bruker SMART APEXII CCD area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2005)
 $T_{\min} = 0.800$, $T_{\max} = 0.926$
10176 measured reflections
2673 independent reflections

2365 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$
 $\theta_{\text{max}} = 28.4^\circ, \theta_{\text{min}} = 2.5^\circ$

$h = -9 \rightarrow 9$
 $k = -13 \rightarrow 13$
 $l = -17 \rightarrow 19$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.110$
 $S = 1.07$
2673 reflections
138 parameters
0 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0594P)^2 + 0.2733P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.31 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.24 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.26170 (16)	0.48062 (13)	0.97372 (9)	0.0346 (3)
C2	0.24057 (17)	0.58273 (14)	1.03826 (10)	0.0386 (3)
C3	0.2907 (2)	0.71257 (15)	1.01635 (13)	0.0510 (4)
H3	0.2758	0.7812	1.0591	0.061*
C4	0.3626 (2)	0.73967 (16)	0.93147 (14)	0.0573 (4)
H4	0.3962	0.8268	0.9175	0.069*
C5	0.3857 (2)	0.63922 (17)	0.86653 (12)	0.0512 (4)
C6	0.33436 (19)	0.51057 (16)	0.88716 (10)	0.0438 (3)
H6	0.3478	0.4429	0.8435	0.053*
C7	0.4842 (3)	0.5753 (3)	0.71703 (14)	0.0797 (7)
H7A	0.5574	0.5054	0.7437	0.120*
H7B	0.5415	0.6124	0.6641	0.120*
H7C	0.3701	0.5392	0.6978	0.120*
C8	0.21137 (17)	0.34293 (13)	0.99115 (9)	0.0365 (3)
H8	0.2270	0.2798	0.9445	0.044*
C9	0.02639 (18)	0.11834 (12)	1.14708 (9)	0.0353 (3)
N1	0.14624 (14)	0.30443 (10)	1.06812 (8)	0.0349 (2)
N2	0.10299 (16)	0.17032 (11)	1.07211 (8)	0.0388 (3)
H2	0.1255	0.1192	1.0258	0.047*
N3	0.0098 (2)	0.19397 (12)	1.22149 (8)	0.0507 (3)
H3A	0.0478	0.2752	1.2214	0.061*
H3B	-0.0390	0.1620	1.2701	0.061*

O1	0.17090 (17)	0.56061 (11)	1.12351 (8)	0.0514 (3)
H1	0.1484	0.4807	1.1292	0.077*
O2	0.4601 (2)	0.67718 (16)	0.78420 (11)	0.0777 (4)
S1	-0.04238 (6)	-0.04281 (3)	1.14369 (3)	0.04739 (14)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0324 (6)	0.0336 (6)	0.0378 (6)	-0.0011 (5)	0.0011 (5)	0.0040 (5)
C2	0.0356 (6)	0.0346 (6)	0.0455 (7)	0.0003 (5)	0.0019 (5)	0.0011 (5)
C3	0.0486 (8)	0.0323 (7)	0.0722 (10)	-0.0006 (6)	0.0028 (7)	-0.0008 (7)
C4	0.0489 (8)	0.0387 (7)	0.0844 (12)	-0.0041 (6)	0.0022 (8)	0.0213 (8)
C5	0.0422 (7)	0.0543 (9)	0.0573 (9)	-0.0021 (6)	0.0051 (6)	0.0232 (7)
C6	0.0430 (7)	0.0474 (8)	0.0411 (7)	-0.0017 (6)	0.0048 (5)	0.0072 (6)
C7	0.0654 (11)	0.123 (2)	0.0515 (10)	0.0004 (12)	0.0163 (9)	0.0260 (12)
C8	0.0411 (6)	0.0329 (6)	0.0356 (6)	-0.0024 (5)	0.0041 (5)	-0.0009 (5)
C9	0.0440 (6)	0.0293 (6)	0.0326 (6)	0.0004 (5)	0.0021 (5)	0.0016 (5)
N1	0.0401 (5)	0.0285 (5)	0.0362 (5)	-0.0025 (4)	0.0027 (4)	0.0010 (4)
N2	0.0540 (6)	0.0282 (5)	0.0346 (5)	-0.0040 (4)	0.0091 (5)	-0.0012 (4)
N3	0.0834 (9)	0.0347 (6)	0.0347 (6)	-0.0109 (6)	0.0138 (6)	-0.0034 (5)
O1	0.0678 (7)	0.0401 (5)	0.0473 (6)	-0.0046 (5)	0.0163 (5)	-0.0072 (4)
O2	0.0770 (9)	0.0822 (10)	0.0750 (9)	-0.0103 (7)	0.0193 (7)	0.0389 (8)
S1	0.0754 (3)	0.02908 (19)	0.0382 (2)	-0.00904 (14)	0.01049 (17)	0.00096 (12)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.3912 (19)	C7—H7A	0.9600
C1—C6	1.4063 (18)	C7—H7B	0.9600
C1—C8	1.4499 (17)	C7—H7C	0.9600
C2—O1	1.3653 (17)	C8—N1	1.2824 (16)
C2—C3	1.389 (2)	C8—H8	0.9300
C3—C4	1.376 (2)	C9—N3	1.3187 (17)
C3—H3	0.9300	C9—N2	1.3415 (16)
C4—C5	1.385 (3)	C9—S1	1.6901 (13)
C4—H4	0.9300	N1—N2	1.3797 (14)
C5—C6	1.376 (2)	N2—H2	0.8600
C5—O2	1.3774 (19)	N3—H3A	0.8600
C6—H6	0.9300	N3—H3B	0.8600
C7—O2	1.418 (3)	O1—H1	0.8200
C2—C1—C6		H7A—C7—H7B	109.5
C2—C1—C8		O2—C7—H7C	109.5
C6—C1—C8		H7A—C7—H7C	109.5
O1—C2—C3		H7B—C7—H7C	109.5
O1—C2—C1		N1—C8—C1	122.80 (12)
C3—C2—C1		N1—C8—H8	118.6
C4—C3—C2		C1—C8—H8	118.6
C4—C3—H3		N3—C9—N2	118.93 (12)

C2—C3—H3	120.0	N3—C9—S1	122.17 (10)
C3—C4—C5	121.07 (14)	N2—C9—S1	118.89 (10)
C3—C4—H4	119.5	C8—N1—N2	115.04 (11)
C5—C4—H4	119.5	C9—N2—N1	121.08 (11)
C6—C5—O2	124.45 (17)	C9—N2—H2	119.5
C6—C5—C4	119.29 (14)	N1—N2—H2	119.5
O2—C5—C4	116.26 (15)	C9—N3—H3A	120.0
C5—C6—C1	120.53 (15)	C9—N3—H3B	120.0
C5—C6—H6	119.7	H3A—N3—H3B	120.0
C1—C6—H6	119.7	C2—O1—H1	109.5
O2—C7—H7A	109.5	C5—O2—C7	116.83 (15)
O2—C7—H7B	109.5		
C6—C1—C2—O1	179.97 (12)	C2—C1—C6—C5	0.6 (2)
C8—C1—C2—O1	0.2 (2)	C8—C1—C6—C5	-179.64 (13)
C6—C1—C2—C3	0.2 (2)	C2—C1—C8—N1	-1.2 (2)
C8—C1—C2—C3	-179.54 (13)	C6—C1—C8—N1	179.03 (12)
O1—C2—C3—C4	179.62 (14)	C1—C8—N1—N2	178.93 (11)
C1—C2—C3—C4	-0.6 (2)	N3—C9—N2—N1	-7.2 (2)
C2—C3—C4—C5	0.2 (2)	S1—C9—N2—N1	173.92 (10)
C3—C4—C5—C6	0.6 (2)	C8—N1—N2—C9	-177.01 (12)
C3—C4—C5—O2	-179.49 (14)	C6—C5—O2—C7	0.0 (2)
O2—C5—C6—C1	179.10 (14)	C4—C5—O2—C7	-179.90 (16)
C4—C5—C6—C1	-1.0 (2)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1···N1	0.82	1.97	2.6844 (15)	146
N2—H2···S1 ⁱ	0.86	2.61	3.3706 (12)	148
N3—H3A···S1 ⁱⁱ	0.86	2.66	3.2706 (12)	129
N3—H3B···O1 ⁱⁱⁱ	0.86	2.11	2.9604 (17)	172

Symmetry codes: (i) $-x, -y, -z+2$; (ii) $-x, y+1/2, -z+5/2$; (iii) $-x, y-1/2, -z+5/2$.