

Bis(η^6 -naphthalene)molybdenum(0)**Mikhail E. Minyaev,^{a*} John E. Ellis^b and William J. Wolf^b**

^aInstitute of Organoelement Compounds, 28 Vavilova Str., 117813 Moscow, Russian Federation, and ^bDepartment of Chemistry, University of Minnesota, Minneapolis, 207 Pleasant Str. SE, MN 55455, USA

Correspondence e-mail: mminyaev@mail.ru


Received 22 December 2011; accepted 19 January 2012

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in main residue; R factor = 0.028; wR factor = 0.069; data-to-parameter ratio = 12.7.

The title compound, $[\text{Mo}(\text{C}_{10}\text{H}_8)_2]$, was prepared from the naphthalene radical anion and $\text{MoCl}_4(\text{thf})_2$ (thf is tetrahydrofuran). In the crystal, the molecule is located on an inversion center. The Mo atom is equally disordered over two positions; the range of Mo–C distances is 2.2244 (19)–2.3400 (17) Å for both components of the disorder.

Related literature

For background to transition metal–arene complexes, see: Seydel (2002a,b). For the structures of the isotopic Cr and V complexes, see: Elschenbroich *et al.* (1982); Pomije *et al.* (1997). For the structures of homoleptic naphthalenate ate-complexes, see: Jang & Ellis (1994); Brennessel *et al.* (2002, 2006). For the preparation of the title compound, see: Kündig & Timms (1977); Thi *et al.* (1992); Pomije *et al.* (1997).

**Experimental***Crystal data*

$[\text{Mo}(\text{C}_{10}\text{H}_8)_2]$	$V = 707.47 (15)$ Å ³
$M_r = 352.27$	$Z = 2$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 8.4452 (10)$ Å	$\mu = 0.92$ mm ^{−1}
$b = 8.0716 (10)$ Å	$T = 173$ K
$c = 10.9890 (13)$ Å	$0.60 \times 0.60 \times 0.30$ mm
$\beta = 109.186 (2)$ °	

Data collection

Bruker SMART Platform CCD diffractometer	8208 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2003)	1674 independent reflections
$T_{\min} = 0.610$, $T_{\max} = 0.771$	1420 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$	132 parameters
$wR(F^2) = 0.069$	All H-atom parameters refined
$S = 1.03$	$\Delta\rho_{\max} = 0.27$ e Å ^{−3}
1674 reflections	$\Delta\rho_{\min} = -0.28$ e Å ^{−3}

Data collection: *SMART* (Bruker, 2003); cell refinement: *SAINT* (Bruker, 2003); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors are grateful to the National Science Foundation program (grant CHE-0841014).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK5041).

References

- Brennessel, W. W., Ellis, J. E., Pomije, M. K., Sussman, V. J., Urnezius, E. & Young, V. G. (2002). *J. Am. Chem. Soc.* **124**, 10258–10259.
- Brennessel, W. W., Young, V. G. & Ellis, J. E. (2006). *Angew. Chem. Int. Ed.* **45**, 7268–7271.
- Bruker (2003). *SADABS, SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Elschenbroich, C., Mockel, R., Massa, W., Birkhahn, M. & Zenneck, U. (1982). *Chem. Ber.* **115**, 334–345.
- Jang, M. & Ellis, J. E. (1994). *Angew. Chem. Int. Ed. Engl.* **33**, 1973–1975.
- Kündig, E. P. & Timms, P. L. (1977). *J. Chem. Soc. Chem. Commun.* pp. 912–913.
- Pomije, M. K., Kurth, C. J., Ellis, J. E. & Barybin, M. V. (1997). *Organometallics*, **16**, 3582–3587.
- Seydel, D. (2002a). *Organometallics*, **21**, 1520–1530.
- Seydel, D. (2002b). *Organometallics*, **21**, 2800–2820.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Thi, N. P. D., Spicher, S., Paglia, P., Bernardinelli, G., Kündig, E. P. & Timms, P. L. (1992). *Helv. Chim. Acta*, **75**, 2593–2607.

supporting information

Acta Cryst. (2012). E68, m220 [doi:10.1107/S1600536812002504]

Bis(η^6 -naphthalene)molybdenum(0)

Mikhail E. Minyaev, John E. Ellis and William J. Wolf

S1. Comment

Studies of classical arene complexes of transition elements have had significant impact on understanding metal – organic ligand bonding (Seyferth, 2002*a,b*). However, only a few molecular structures of d-metal neutral homoleptic naphthalene species, i.e. Cr(η^6 -C₁₀H₈)₂ (Elschenbroich *et al.*, 1982), V(η^6 -C₁₀H₈)₂ (Pomije *et al.*, 1997) and homoleptic naphthalene *ate-complexes* [Zr(η^4 -C₁₀H₈)₃]²⁻ (Jang *et al.*, 1994), [Ta(η^4 -C₁₀H₈)₃]⁻ (Brennessel *et al.*, 2002) and [Co(η^4 -C₁₀H₈)₂]⁻ (Brennessel *et al.*, 2006), have been established.

Previous examples of neutral homoleptic naphthalene complexes of transition metals were obtained by metal-atom-ligand-vapor co-condensation synthesis (Kündig *et al.*, 1977). The title compound, Mo(η^6 -C₁₀H₈)₂, was later obtained by reduction of MoCl₅ or MoCl₄(thf)₂ with Mg, followed by evaporation of Mo atoms along with free naphthalene under vacuum (Thi *et al.*, 1992), or by reduction of MoCl₄(thf)₂ or MoCl₃(thf)₃ with lithium naphthalenide (Pomije *et al.*, 1997).

We prepared Mo(C₁₀H₈)₂ from LiC₁₀H₈ and MoCl₄(thf)₂ under argon in THF media in 13% yield. The ¹H NMR spectrum is identical with previously described by Pomije *et al.* (1997). The Mo(C₁₀H₈)₂ crystal structure is isomorphous with previously published structures of Cr(C₁₀H₈)₂ (Elschenbroich *et al.*, 1982) and V(C₁₀H₈)₂ (Pomije *et al.*, 1997). The molecule is located on the inversion center, one half of the molecule is unique. The molybdenum atom is disordered equally between two positions with occupancy of 50%. The C–C distances are similar to those found in Cr(C₁₀H₈)₂ and V(C₁₀H₈)₂. The Mo–centroid(C₆) distances are 1.782 Å (C1–C2–C3–C4–C9–C10) and 1.770 Å (C5–C6–C7–C8–C10–C9) and centroid(C₆)–Mo–centroid(C₆) angle is 178°. The naphthalene ligand is slightly bent along the C9–C10 bond towards molybdenum atom, the angle between flat rings in the naphthalene ligand is 177°.

S2. Experimental

All synthetic manipulations were carried out in atmosphere of purified argon or under vacuum, using Schlenk type glassware and dry box techniques. Freshly cut lithium wire (0.34 g, 49 mmol), sublimed naphthalene (9.15 g, 71.4 mmol) and THF (200 ml) were added to a 1 L round-bottomed flask equipped with a glass covered stirbar. The mixture was stirred for 17 h at ambient temperature, then the resulting deep green solution was cooled to -60°C. A cold (-60°C) suspension of MoCl₄(thf)₂ (4.5 g, 11.8 mmol) and C₁₀H₈ (6.10 g, 47.6 mmol) in THF (100 ml) was added dropwise to the cold solution of lithium naphthalenide *via* large cannula. The reaction mixture was allowed to warm to ambient temperature over 15 h. It was filtered to provide a dark red solution, then THF was removed under vacuum. The resulting solid was stirred in toluene (200 ml), the mixture was filtered, and all toluene was removed from the solution under vacuum. The remaining solid was transferred to a sublimator, and an excess of naphthalene was removed by sublimation under vacuum at 30°C over three days. The remaining purple brown solid was taken up in toluene (100 ml) and the solution was filtered. All but 5 ml of toluene was removed under vacuum, pentane (100 ml) was added. The resulting mixture was cooled to -78°C and stirred. The precipitate was filtered off to give Mo(C₁₀H₈)₂ (0.539 g, 1.53 mmol, 13.0%) as a dark purple microcrystalline solid. ¹H NMR (300 MHz, C₆D₆, 25°C): δ (p.p.m.) = 4.67 (m, 2H), 5.02 (m, 2H), 6.56

(m, 2H), 6.78 (m, 2H) p.p.m..

X-ray quality single crystals were grown at 0°C over a 2 week period by slow diffusion of pentane (100 ml) into a nearly saturated solution of $\text{Mo}(\text{C}_{10}\text{H}_8)_2$ (100 mg) in toluene (c.a. 20 ml).

S3. Refinement

The C—H atoms were refined freely; range of C—H distances = 0.89 (2) to 0.94 (2) Å. The Mo atom is disordered equally between two positions with occupancy of 50% each.




Figure 1

Molecular structure of $\text{Mo}(\text{C}_{10}\text{H}_8)_2$ (50% atomic displacement parameters). The asymmetric unit consists of one half of the molecule. The occupancies of Mo1 and $\text{Mo}1^i$ sites are 50%. Symmetry code: (i) $-x + 1, -y, -z + 2$.

Bis(η^6 -naphthalene)molybdenum(0)

Crystal data

$[\text{Mo}(\text{C}_{10}\text{H}_8)_2]$

$M_r = 352.27$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 8.4452 (10)$ Å

$b = 8.0716 (10)$ Å

$c = 10.9890 (13)$ Å

$\beta = 109.186 (2)^\circ$

$V = 707.47 (15)$ Å³

$Z = 2$

$F(000) = 356$

$D_x = 1.654 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2191 reflections

$\theta = 2.7\text{--}28.3^\circ$

$\mu = 0.92 \text{ mm}^{-1}$

$T = 173$ K

Block, dark-red

$0.60 \times 0.60 \times 0.30$ mm

Data collection

Bruker SMART Platform CCD
diffractometer
Radiation source: normal-focus sealed tube
Graphite monochromator
area detector, ω and φ scans
Absorption correction: multi-scan
(SADABS; Bruker, 2003)
 $T_{\min} = 0.610$, $T_{\max} = 0.771$

8208 measured reflections
1674 independent reflections
1420 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$
 $\theta_{\max} = 27.9^\circ$, $\theta_{\min} = 2.7^\circ$
 $h = -10 \rightarrow 11$
 $k = -10 \rightarrow 10$
 $l = -14 \rightarrow 14$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.069$
 $S = 1.03$
1674 reflections
132 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
All H-atom parameters refined
 $w = 1/[\sigma^2(F_o^2) + (0.0357P)^2 + 0.251P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.27 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Mo1	0.63424 (3)	0.01329 (3)	0.97463 (3)	0.02253 (11)	0.50
C1	0.4807 (3)	0.0582 (3)	0.76370 (18)	0.0382 (4)	
C2	0.6382 (3)	0.1252 (3)	0.7895 (2)	0.0481 (6)	
C3	0.7005 (3)	0.2429 (3)	0.8901 (2)	0.0452 (5)	
C4	0.6048 (2)	0.2901 (2)	0.96470 (19)	0.0358 (4)	
C9	0.4393 (2)	0.22530 (19)	0.93978 (16)	0.0261 (4)	
C5	0.3393 (2)	0.2642 (2)	1.01832 (17)	0.0318 (4)	
C6	0.1831 (3)	0.1938 (2)	0.99537 (19)	0.0358 (4)	
C7	0.1202 (2)	0.0780 (3)	0.89365 (19)	0.0372 (4)	
C8	0.2145 (2)	0.0343 (2)	0.81728 (17)	0.0330 (4)	
C10	0.3753 (2)	0.1064 (2)	0.83615 (15)	0.0277 (4)	
H1	0.442 (3)	-0.021 (2)	0.702 (2)	0.040 (6)*	
H2	0.707 (3)	0.091 (3)	0.744 (2)	0.058 (7)*	
H3	0.806 (3)	0.289 (3)	0.910 (2)	0.058 (7)*	
H4	0.641 (3)	0.363 (3)	1.032 (2)	0.037 (6)*	
H5	0.380 (2)	0.337 (2)	1.086 (2)	0.034 (5)*	

H6	0.118 (3)	0.218 (3)	1.048 (2)	0.047 (6)*
H7	0.021 (3)	0.024 (3)	0.881 (2)	0.049 (7)*
H8	0.179 (3)	-0.044 (3)	0.758 (2)	0.034 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mo1	0.02548 (16)	0.02166 (16)	0.01930 (16)	0.00090 (11)	0.00579 (11)	0.00237 (10)
C1	0.0496 (12)	0.0428 (11)	0.0217 (9)	0.0128 (9)	0.0111 (8)	0.0065 (8)
C2	0.0527 (13)	0.0604 (14)	0.0398 (12)	0.0210 (11)	0.0268 (11)	0.0232 (10)
C3	0.0342 (11)	0.0442 (12)	0.0579 (14)	0.0024 (9)	0.0160 (10)	0.0263 (10)
C4	0.0372 (11)	0.0238 (9)	0.0396 (11)	-0.0031 (7)	0.0036 (9)	0.0086 (8)
C9	0.0314 (9)	0.0194 (8)	0.0246 (8)	0.0017 (6)	0.0054 (7)	0.0048 (6)
C5	0.0420 (11)	0.0229 (8)	0.0276 (9)	0.0049 (7)	0.0078 (8)	-0.0004 (7)
C6	0.0368 (10)	0.0364 (10)	0.0359 (10)	0.0079 (8)	0.0143 (9)	0.0045 (8)
C7	0.0286 (10)	0.0366 (10)	0.0410 (11)	-0.0017 (8)	0.0043 (8)	0.0086 (8)
C8	0.0390 (10)	0.0259 (9)	0.0245 (9)	-0.0003 (7)	-0.0025 (8)	0.0005 (7)
C10	0.0350 (9)	0.0248 (8)	0.0194 (8)	0.0050 (7)	0.0038 (7)	0.0056 (6)

Geometric parameters (\AA , ^\circ)

Mo1—C1	2.2826 (19)	C4—C9	1.432 (3)
Mo1—C2	2.236 (2)	C9—C5	1.427 (3)
Mo1—C3	2.225 (2)	C5—C6	1.381 (3)
Mo1—C4	2.2466 (19)	C6—C7	1.419 (3)
Mo1—C9	2.3176 (16)	C7—C8	1.379 (3)
Mo1—C10	2.3400 (17)	C8—C10	1.429 (3)
Mo1—C5 ⁱ	2.2495 (18)	C9—C10	1.451 (2)
Mo1—C6 ⁱ	2.2244 (19)	C1—H1	0.91 (2)
Mo1—C7 ⁱ	2.2284 (19)	C2—H2	0.93 (2)
Mo1—C8 ⁱ	2.2557 (18)	C3—H3	0.92 (3)
Mo1—C9 ⁱ	2.3158 (16)	C4—H4	0.91 (2)
Mo1—C10 ⁱ	2.3182 (16)	C5—H5	0.92 (2)
C1—C10	1.429 (3)	C6—H6	0.94 (2)
C1—C2	1.377 (3)	C7—H7	0.91 (3)
C2—C3	1.421 (3)	C8—H8	0.89 (2)
C3—C4	1.380 (3)		
C6 ⁱ —Mo1—C3	115.50 (8)	C2—C1—Mo1	70.43 (12)
C6 ⁱ —Mo1—C7 ⁱ	37.18 (7)	C10—C1—Mo1	74.19 (10)
C3—Mo1—C7 ⁱ	103.80 (8)	C1—C2—C3	120.6 (2)
C6 ⁱ —Mo1—C2	102.80 (8)	C1—C2—Mo1	74.12 (12)
C3—Mo1—C2	37.15 (9)	C3—C2—Mo1	71.01 (12)
C7 ⁱ —Mo1—C2	115.83 (8)	C4—C3—C2	120.5 (2)
C6 ⁱ —Mo1—C4	144.57 (7)	C4—C3—Mo1	72.87 (11)
C3—Mo1—C4	35.95 (8)	C2—C3—Mo1	71.84 (12)
C7 ⁱ —Mo1—C4	115.15 (7)	C3—C4—C9	120.73 (19)
C2—Mo1—C4	65.70 (8)	C3—C4—Mo1	71.18 (12)

C6 ⁱ —Mo1—C5 ⁱ	35.95 (7)	C9—C4—Mo1	74.44 (10)
C3—Mo1—C5 ⁱ	143.88 (8)	C5—C9—C4	122.59 (17)
C7 ⁱ —Mo1—C5 ⁱ	65.64 (7)	C5—C9—C10	118.58 (16)
C2—Mo1—C5 ⁱ	113.74 (8)	C4—C9—C10	118.68 (17)
C4—Mo1—C5 ⁱ	179.14 (7)	C5—C9—Mo1 ⁱ	69.28 (9)
C6 ⁱ —Mo1—C8 ⁱ	65.65 (7)	C4—C9—Mo1 ⁱ	126.95 (11)
C3—Mo1—C8 ⁱ	115.38 (8)	C10—C9—Mo1 ⁱ	71.85 (9)
C7 ⁱ —Mo1—C8 ⁱ	35.82 (7)	C5—C9—Mo1	126.30 (11)
C2—Mo1—C8 ⁱ	144.91 (9)	C4—C9—Mo1	69.04 (10)
C4—Mo1—C8 ⁱ	103.61 (7)	C10—C9—Mo1	72.69 (9)
C5 ⁱ —Mo1—C8 ⁱ	77.22 (7)	Mo1 ⁱ —C9—Mo1	65.77 (4)
C6 ⁱ —Mo1—C1	113.87 (7)	C6—C5—C9	121.25 (17)
C3—Mo1—C1	65.22 (9)	C6—C5—Mo1 ⁱ	71.04 (11)
C7 ⁱ —Mo1—C1	144.00 (8)	C9—C5—Mo1 ⁱ	74.33 (9)
C2—Mo1—C1	35.45 (8)	C5—C6—C7	120.16 (19)
C4—Mo1—C1	77.05 (7)	C5—C6—Mo1 ⁱ	73.01 (11)
C5 ⁱ —Mo1—C1	102.12 (7)	C7—C6—Mo1 ⁱ	71.56 (11)
C8 ⁱ —Mo1—C1	179.33 (7)	C8—C7—C6	120.40 (19)
C6 ⁱ —Mo1—C9 ⁱ	65.18 (7)	C8—C7—Mo1 ⁱ	73.17 (11)
C3—Mo1—C9 ⁱ	179.04 (7)	C6—C7—Mo1 ⁱ	71.26 (11)
C7 ⁱ —Mo1—C9 ⁱ	77.15 (7)	C7—C8—C10	121.25 (17)
C2—Mo1—C9 ⁱ	142.34 (8)	C7—C8—Mo1 ⁱ	71.01 (11)
C4—Mo1—C9 ⁱ	143.76 (7)	C10—C8—Mo1 ⁱ	74.20 (10)
C5 ⁱ —Mo1—C9 ⁱ	36.39 (6)	C8—C10—C1	122.96 (17)
C8 ⁱ —Mo1—C9 ⁱ	65.46 (6)	C8—C10—C9	118.33 (16)
C1—Mo1—C9 ⁱ	113.94 (7)	C1—C10—C9	118.56 (17)
C6 ⁱ —Mo1—C9	178.65 (7)	C8—C10—Mo1 ⁱ	69.43 (10)
C3—Mo1—C9	65.07 (7)	C1—C10—Mo1 ⁱ	126.69 (12)
C7 ⁱ —Mo1—C9	144.14 (7)	C9—C10—Mo1 ⁱ	71.66 (9)
C2—Mo1—C9	76.91 (7)	C8—C10—Mo1	127.15 (11)
C4—Mo1—C9	36.52 (6)	C1—C10—Mo1	69.81 (10)
C5 ⁱ —Mo1—C9	142.94 (7)	C9—C10—Mo1	71.02 (9)
C8 ⁱ —Mo1—C9	115.35 (7)	Mo1 ⁱ —C10—Mo1	65.37 (4)
C1—Mo1—C9	65.12 (7)	C2—C1—H1	120.8 (14)
C9 ⁱ —Mo1—C9	114.23 (4)	C10—C1—H1	118.2 (14)
C6 ⁱ —Mo1—C10 ⁱ	77.33 (7)	Mo1—C1—H1	126.2 (13)
C3—Mo1—C10 ⁱ	144.01 (8)	C1—C2—H2	120.5 (16)
C7 ⁱ —Mo1—C10 ⁱ	65.09 (7)	C3—C2—H2	118.9 (16)
C2—Mo1—C10 ⁱ	178.65 (8)	Mo1—C2—H2	125.0 (15)
C4—Mo1—C10 ⁱ	114.97 (7)	C4—C3—H3	117.5 (15)
C5 ⁱ —Mo1—C10 ⁱ	65.57 (6)	C2—C3—H3	122.1 (15)
C8 ⁱ —Mo1—C10 ⁱ	36.37 (7)	Mo1—C3—H3	126.5 (15)
C1—Mo1—C10 ⁱ	143.26 (7)	C3—C4—H4	123.6 (13)
C9 ⁱ —Mo1—C10 ⁱ	36.49 (6)	C9—C4—H4	115.7 (13)
C9—Mo1—C10 ⁱ	102.93 (6)	Mo1—C4—H4	126.8 (13)
C6 ⁱ —Mo1—C10	142.39 (7)	C6—C5—H5	119.4 (12)
C3—Mo1—C10	76.73 (7)	C9—C5—H5	119.4 (12)
C7 ⁱ —Mo1—C10	179.42 (7)	Mo1 ⁱ —C5—H5	127.2 (12)

C2—Mo1—C10	64.45 (7)	C5—C6—H6	121.2 (14)
C4—Mo1—C10	65.41 (6)	C7—C6—H6	118.6 (14)
C5 ⁱ —Mo1—C10	113.79 (6)	Mo1 ⁱ —C6—H6	125.7 (14)
C8 ⁱ —Mo1—C10	144.17 (7)	C8—C7—H7	118.3 (15)
C1—Mo1—C10	35.99 (6)	C6—C7—H7	121.1 (15)
C9 ⁱ —Mo1—C10	102.32 (6)	Mo1 ⁱ —C7—H7	123.6 (14)
C9—Mo1—C10	36.29 (6)	C7—C8—H8	120.6 (13)
C10 ⁱ —Mo1—C10	114.63 (4)	C10—C8—H8	118.1 (13)
C2—C1—C10	121.0 (2)	Mo1 ⁱ —C8—H8	124.5 (13)
C6 ⁱ —Mo1—C1—C2	-78.24 (15)	C9 ⁱ —Mo1—C9—C4	151.44 (12)
C3—Mo1—C1—C2	30.08 (13)	C10 ⁱ —Mo1—C9—C4	114.69 (11)
C7 ⁱ —Mo1—C1—C2	-49.1 (2)	C10—Mo1—C9—C4	-131.12 (15)
C4—Mo1—C1—C2	66.11 (14)	C3—Mo1—C9—C10	101.84 (12)
C5 ⁱ —Mo1—C1—C2	-114.10 (14)	C7 ⁱ —Mo1—C9—C10	-179.35 (11)
C9 ⁱ —Mo1—C1—C2	-150.44 (13)	C2—Mo1—C9—C10	64.44 (11)
C9—Mo1—C1—C2	102.75 (15)	C4—Mo1—C9—C10	131.12 (15)
C10 ⁱ —Mo1—C1—C2	-179.30 (12)	C5 ⁱ —Mo1—C9—C10	-47.76 (15)
C10—Mo1—C1—C2	131.9 (2)	C8 ⁱ —Mo1—C9—C10	-150.53 (10)
C6 ⁱ —Mo1—C1—C10	149.85 (11)	C1—Mo1—C9—C10	28.93 (10)
C3—Mo1—C1—C10	-101.83 (13)	C9 ⁱ —Mo1—C9—C10	-77.43 (9)
C7 ⁱ —Mo1—C1—C10	179.02 (11)	C10 ⁱ —Mo1—C9—C10	-114.18 (8)
C2—Mo1—C1—C10	-131.9 (2)	C3—Mo1—C9—Mo1 ⁱ	179.28 (8)
C4—Mo1—C1—C10	-65.80 (12)	C7 ⁱ —Mo1—C9—Mo1 ⁱ	-101.91 (11)
C5 ⁱ —Mo1—C1—C10	113.99 (12)	C2—Mo1—C9—Mo1 ⁱ	141.87 (8)
C9 ⁱ —Mo1—C1—C10	77.65 (12)	C4—Mo1—C9—Mo1 ⁱ	-151.44 (12)
C9—Mo1—C1—C10	-29.17 (10)	C5 ⁱ —Mo1—C9—Mo1 ⁱ	29.67 (11)
C10 ⁱ —Mo1—C1—C10	48.78 (19)	C8 ⁱ —Mo1—C9—Mo1 ⁱ	-73.10 (7)
C10—C1—C2—C3	0.7 (3)	C1—Mo1—C9—Mo1 ⁱ	106.36 (7)
Mo1—C1—C2—C3	-55.86 (17)	C9 ⁱ —Mo1—C9—Mo1 ⁱ	0.0
C10—C1—C2—Mo1	56.59 (17)	C10 ⁱ —Mo1—C9—Mo1 ⁱ	-36.75 (6)
C6 ⁱ —Mo1—C2—C1	113.35 (13)	C10—Mo1—C9—Mo1 ⁱ	77.43 (9)
C3—Mo1—C2—C1	-131.1 (2)	C4—C9—C5—C6	-177.00 (16)
C7 ⁱ —Mo1—C2—C1	150.43 (12)	C10—C9—C5—C6	-1.4 (2)
C4—Mo1—C2—C1	-102.13 (14)	Mo1 ⁱ —C9—C5—C6	-55.60 (15)
C5 ⁱ —Mo1—C2—C1	77.16 (14)	Mo1—C9—C5—C6	-90.3 (2)
C8 ⁱ —Mo1—C2—C1	179.02 (12)	C4—C9—C5—Mo1 ⁱ	-121.39 (15)
C9 ⁱ —Mo1—C2—C1	47.56 (19)	C10—C9—C5—Mo1 ⁱ	54.17 (13)
C9—Mo1—C2—C1	-65.29 (13)	Mo1—C9—C5—Mo1 ⁱ	-34.68 (12)
C10—Mo1—C2—C1	-29.00 (12)	C9—C5—C6—C7	1.2 (3)
C6 ⁱ —Mo1—C2—C3	-115.54 (13)	Mo1 ⁱ —C5—C6—C7	-55.95 (16)
C7 ⁱ —Mo1—C2—C3	-78.46 (14)	C9—C5—C6—Mo1 ⁱ	57.15 (15)
C4—Mo1—C2—C3	28.97 (12)	C5—C6—C7—C8	0.3 (3)
C5 ⁱ —Mo1—C2—C3	-151.74 (12)	Mo1 ⁱ —C6—C7—C8	-56.39 (16)
C8 ⁱ —Mo1—C2—C3	-49.88 (19)	C5—C6—C7—Mo1 ⁱ	56.64 (16)
C1—Mo1—C2—C3	131.10 (19)	C6—C7—C8—C10	-1.4 (3)
C9 ⁱ —Mo1—C2—C3	178.66 (11)	Mo1 ⁱ —C7—C8—C10	-56.93 (15)
C9—Mo1—C2—C3	65.81 (13)	C6—C7—C8—Mo1 ⁱ	55.48 (16)

C10—Mo1—C2—C3	102.10 (13)	C7—C8—C10—C1	176.59 (16)
C1—C2—C3—C4	0.9 (3)	Mo1 ⁱ —C8—C10—C1	121.15 (16)
Mo1—C2—C3—C4	-56.46 (16)	C7—C8—C10—C9	1.2 (2)
C1—C2—C3—Mo1	57.35 (17)	Mo1 ⁱ —C8—C10—C9	-54.27 (13)
C6 ⁱ —Mo1—C3—C4	-151.64 (12)	C7—C8—C10—Mo1 ⁱ	55.44 (16)
C7 ⁱ —Mo1—C3—C4	-114.00 (13)	C7—C8—C10—Mo1	88.0 (2)
C2—Mo1—C3—C4	131.25 (19)	Mo1 ⁱ —C8—C10—Mo1	32.57 (12)
C5 ⁱ —Mo1—C3—C4	178.57 (12)	C2—C1—C10—C8	-176.75 (16)
C8 ⁱ —Mo1—C3—C4	-77.86 (14)	Mo1—C1—C10—C8	-121.91 (16)
C1—Mo1—C3—C4	102.48 (14)	C2—C1—C10—C9	-1.3 (3)
C9—Mo1—C3—C4	29.72 (11)	Mo1—C1—C10—C9	53.51 (13)
C10 ⁱ —Mo1—C3—C4	-47.57 (18)	C2—C1—C10—Mo1 ⁱ	-89.0 (2)
C10—Mo1—C3—C4	66.25 (12)	Mo1—C1—C10—Mo1 ⁱ	-34.14 (14)
C6 ⁱ —Mo1—C3—C2	77.11 (14)	C2—C1—C10—Mo1	-54.85 (17)
C7 ⁱ —Mo1—C3—C2	114.75 (13)	C5—C9—C10—C8	0.3 (2)
C4—Mo1—C3—C2	-131.25 (19)	C4—C9—C10—C8	176.00 (15)
C5 ⁱ —Mo1—C3—C2	47.32 (19)	Mo1 ⁱ —C9—C10—C8	53.20 (13)
C8 ⁱ —Mo1—C3—C2	150.89 (12)	Mo1—C9—C10—C8	122.69 (14)
C1—Mo1—C3—C2	-28.78 (12)	C5—C9—C10—C1	-175.37 (15)
C9—Mo1—C3—C2	-101.53 (14)	C4—C9—C10—C1	0.4 (2)
C10 ⁱ —Mo1—C3—C2	-178.82 (12)	Mo1 ⁱ —C9—C10—C1	-122.43 (15)
C10—Mo1—C3—C2	-65.01 (13)	Mo1—C9—C10—C1	-52.93 (14)
C2—C3—C4—C9	-1.9 (3)	C5—C9—C10—Mo1 ⁱ	-52.94 (13)
Mo1—C3—C4—C9	-57.84 (15)	C4—C9—C10—Mo1 ⁱ	122.80 (14)
C2—C3—C4—Mo1	55.97 (17)	Mo1—C9—C10—Mo1 ⁱ	69.50 (4)
C6 ⁱ —Mo1—C4—C3	47.69 (19)	C5—C9—C10—Mo1	-122.43 (14)
C7 ⁱ —Mo1—C4—C3	78.55 (14)	C4—C9—C10—Mo1	53.30 (13)
C2—Mo1—C4—C3	-29.87 (13)	Mo1 ⁱ —C9—C10—Mo1	-69.50 (4)
C8 ⁱ —Mo1—C4—C3	114.66 (13)	C6 ⁱ —Mo1—C10—C8	67.8 (2)
C1—Mo1—C4—C3	-65.46 (13)	C3—Mo1—C10—C8	-177.41 (17)
C9 ⁱ —Mo1—C4—C3	-178.45 (12)	C2—Mo1—C10—C8	145.24 (18)
C9—Mo1—C4—C3	-130.94 (18)	C4—Mo1—C10—C8	-141.19 (18)
C10 ⁱ —Mo1—C4—C3	151.41 (12)	C5 ⁱ —Mo1—C10—C8	39.17 (17)
C10—Mo1—C4—C3	-101.58 (14)	C8 ⁱ —Mo1—C10—C8	-62.2 (2)
C6 ⁱ —Mo1—C4—C9	178.64 (11)	C1—Mo1—C10—C8	116.7 (2)
C3—Mo1—C4—C9	130.94 (18)	C9 ⁱ —Mo1—C10—C8	2.70 (17)
C7 ⁱ —Mo1—C4—C9	-150.51 (11)	C9—Mo1—C10—C8	-111.65 (19)
C2—Mo1—C4—C9	101.07 (12)	C10 ⁱ —Mo1—C10—C8	-33.67 (14)
C8 ⁱ —Mo1—C4—C9	-114.40 (11)	C6 ⁱ —Mo1—C10—C1	-48.81 (17)
C1—Mo1—C4—C9	65.49 (11)	C3—Mo1—C10—C1	65.93 (13)
C9 ⁱ —Mo1—C4—C9	-47.50 (18)	C2—Mo1—C10—C1	28.59 (13)
C10 ⁱ —Mo1—C4—C9	-77.64 (12)	C4—Mo1—C10—C1	102.16 (13)
C10—Mo1—C4—C9	29.36 (10)	C5 ⁱ —Mo1—C10—C1	-77.49 (13)
C3—C4—C9—C5	176.78 (16)	C8 ⁱ —Mo1—C10—C1	-178.89 (12)
Mo1—C4—C9—C5	120.50 (15)	C9 ⁱ —Mo1—C10—C1	-113.95 (12)
C3—C4—C9—C10	1.2 (2)	C9—Mo1—C10—C1	131.69 (16)
Mo1—C4—C9—C10	-55.06 (13)	C10 ⁱ —Mo1—C10—C1	-150.33 (13)
C3—C4—C9—Mo1 ⁱ	89.3 (2)	C6 ⁱ —Mo1—C10—C9	179.49 (11)

Mo1—C4—C9—Mo1 ⁱ	33.05 (13)	C3—Mo1—C10—C9	−65.76 (11)
C3—C4—C9—Mo1	56.28 (16)	C2—Mo1—C10—C9	−103.11 (12)
C3—Mo1—C9—C5	−145.02 (17)	C4—Mo1—C10—C9	−29.54 (10)
C7 ⁱ —Mo1—C9—C5	−66.2 (2)	C5 ⁱ —Mo1—C10—C9	150.82 (10)
C2—Mo1—C9—C5	177.57 (17)	C8 ⁱ —Mo1—C10—C9	49.42 (15)
C4—Mo1—C9—C5	−115.7 (2)	C1—Mo1—C10—C9	−131.69 (16)
C5 ⁱ —Mo1—C9—C5	65.4 (2)	C9 ⁱ —Mo1—C10—C9	114.35 (8)
C8 ⁱ —Mo1—C9—C5	−37.40 (17)	C10 ⁱ —Mo1—C10—C9	77.98 (9)
C1—Mo1—C9—C5	142.06 (17)	C6 ⁱ —Mo1—C10—Mo1 ⁱ	101.52 (11)
C9 ⁱ —Mo1—C9—C5	35.70 (13)	C3—Mo1—C10—Mo1 ⁱ	−143.74 (8)
C10 ⁱ —Mo1—C9—C5	−1.05 (16)	C2—Mo1—C10—Mo1 ⁱ	178.91 (8)
C10—Mo1—C9—C5	113.13 (19)	C4—Mo1—C10—Mo1 ⁱ	−107.52 (7)
C3—Mo1—C9—C4	−29.28 (12)	C5 ⁱ —Mo1—C10—Mo1 ⁱ	72.84 (7)
C7 ⁱ —Mo1—C9—C4	49.53 (16)	C8 ⁱ —Mo1—C10—Mo1 ⁱ	−28.56 (11)
C2—Mo1—C9—C4	−66.68 (12)	C1—Mo1—C10—Mo1 ⁱ	150.33 (13)
C5 ⁱ —Mo1—C9—C4	−178.88 (12)	C9 ⁱ —Mo1—C10—Mo1 ⁱ	36.37 (6)
C8 ⁱ —Mo1—C9—C4	78.35 (12)	C9—Mo1—C10—Mo1 ⁱ	−77.98 (9)
C1—Mo1—C9—C4	−102.19 (13)	C10 ⁱ —Mo1—C10—Mo1 ⁱ	0.0

Symmetry code: (i) $-x+1, -y, -z+2$.