

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(1*E*,2*E*)-1,2-Bis[1-(3-nitrophenyl)ethylidene]hydrazine

Safra Izuani Jama Asik,^a Hoong-Kun Fun,^{a*‡}
Ibrahim Abdul Razak,^a Patcharaporn Jansrisewangwong^b
and Suchada Chantrapromma^{b§}

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand

Correspondence e-mail: hkfun@usm.my


Received 31 January 2012; accepted 3 February 2012

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.047; wR factor = 0.146; data-to-parameter ratio = 20.5.

The asymmetric unit of the title compound, $\text{C}_{16}\text{H}_{14}\text{N}_4\text{O}_4$, contains one half-molecule of (nitrophenyl)ethanimine and the complete molecule is generated by a crystallographic inversion centre. The molecule has an *E* conformation with respect to each $\text{C}=\text{N}$ double bond. The central $\text{C}=\text{N}-\text{N}=\text{C}$ plane is twisted from the benzene rings with a dihedral angle of $24.76(11)^\circ$. In the crystal, $\text{C}-\text{H}\cdots\text{O}$ interactions link the molecules to form sheets that lie parallel to $(10\bar{4})$.

Related literature

For the biological activity of hydrazones, see: Khanmohammadi *et al.* (2008); Luboch *et al.* (2009). For related structures, see: Chantrapromma *et al.* (2011); Fun, Jansrisewangwong *et al.* (2011); Fun, Nilwanna *et al.* (2011); Jansrisewangwong *et al.* (2010); Nilwanna *et al.* (2011). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{14}\text{N}_4\text{O}_4$
 $M_r = 326.31$
Monoclinic, $P2_1/c$
 $a = 3.9296(3)$ Å

$b = 7.4448(5)$ Å
 $c = 26.3979(19)$ Å
 $\beta = 94.022(1)^\circ$
 $V = 770.37(10)$ Å³

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹

$T = 296$ K
 $0.34 \times 0.17 \times 0.10$ mm

Data collection

Bruker APEX DUO CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.966$, $T_{\max} = 0.990$

15392 measured reflections
2254 independent reflections
1686 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.146$
 $S = 1.06$
2254 reflections

110 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.21$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.18$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C3}-\text{H3A}\cdots\text{O1}^i$	0.93	2.57	3.239 (2)	129

Symmetry code: (i) $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HKF, IAR and SIJA thank the Malaysian Government and Universiti Sains Malaysia for Research University grants (Nos. 1001/PFIZIK/811160 and 1001/PFIZIK/811151). PJ and SC thank the Prince of Songkla University for financial support through the Crystal Materials Research Unit.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5067).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chantrapromma, S., Jansrisewangwong, P., Chanawanno, K. & Fun, H.-K. (2011). *Acta Cryst. E67*, o2221–o2222.
- Fun, H.-K., Jansrisewangwong, P., Karalai, C. & Chantrapromma, S. (2011). *Acta Cryst. E67*, o3424.
- Fun, H.-K., Nilwanna, B., Jansrisewangwong, P., Kobkeattawin, T. & Chantrapromma, S. (2011). *Acta Cryst. E67*, o3202–o3203.
- Jansrisewangwong, P., Chantrapromma, S. & Fun, H.-K. (2010). *Acta Cryst. E66*, o2170.
- Khanmohammadi, H., Abnosi, M. H., Hosseinzadeh, A. & Erfantalab, M. (2008). *Spectrochim. Acta A*, **71**, 1474–1480.
- Luboch, E., Wagner-Wysiecka, E. & Rzymowski, T. (2009). *Tetrahedron*, **65**, 10671–10678.
- Nilwanna, B., Chantrapromma, S., Jansrisewangwong, P. & Fun, H.-K. (2011). *Acta Cryst. E67*, o3084–o3085.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

‡ Thomson Reuters ResearcherID: A-3561-2009.

§ Thomson Reuters ResearcherID: A-5085-2009. Additional correspondence author e-mail: suchada.c@psu.ac.th.

supporting information

Acta Cryst. (2012). E68, o643 [doi:10.1107/S1600536812004722]

(1*E*,2*E*)-1,2-Bis[1-(3-nitrophenyl)ethylidene]hydrazine

Safra Izuani Jama Asik, Hoong-Kun Fun, Ibrahim Abdul Razak, Patcharaporn Jansrisewangwong and Suchada Chantraproma

S1. Comment

Hydrazones have been intensively investigated mostly because of their potential applications as antibacterial and antifungal drugs (Khanmohammadi *et al.*, 2008) and as fluorescent chemosensors for metal ions (Luboch *et al.*, 2009). Owing to our medicinal chemistry research on hydrazones, we previously reported the synthesis and crystal structures of some hydrazone derivatives (Chantrapromma *et al.*, 2011; Fun, Jansrisewangwong *et al.*, 2011; Fun, Nilwanna *et al.*, 2011; Jansrisewangwong *et al.*, 2010; Nilwanna *et al.*, 2011). The title compound was synthesized to study for antibacterial activity and fluorescence properties in order to get more detail on the structural activity relationship through comparing with other closely related compounds.

The asymmetric unit of (I) (Fig. 1), contains one half-molecule of (nitrophenyl)ethanimine and the complete molecule is generated by a crystallographic inversion centre (-x,-y + 1,-z). The molecule is in an *E* configuration with respect to C7=N1 double bond [1.2803 (17) Å] with the torsion angle N1A—N1—C7—C1 = 179.92 (10)°. The methyl groups are twisted from the planes of benzene (C1—C6 and C1A—C6A) rings and their orientations can be indicated by the torsion angles (C1—C6—C7—C8 and C1A—C6A—C7A—C8A) = 24.91 (19)°. The bond lengths are within the normal range (Allen *et al.*, 1987) and are comparable with the related structures (Chantrapromma *et al.*, 2011; Fun, Jansrisewangwong *et al.*, 2011; Fun, Nilwanna *et al.*, 2011; Jansrisewangwong *et al.*, 2010; Nilwanna *et al.*, 2011). In the crystal structure (Fig. 2), the C3—H3A⋯O1 interaction links the molecules into two-dimensional layers parallel to the (1 0 $\bar{4}$) plane.

S2. Experimental

The title compound was synthesized by mixing a solution (1:2 molar ratio) of hydrazine hydrate (0.10 ml, 2 mmol) and 3-nitroacetophenone (0.66 g, 4 mmol) in ethanol (20 ml). The resulting solution was refluxed for 4 h, yielding the yellow crystalline solid. The resultant solid was filtered off and washed with methanol. Yellow block-shaped single crystals of the title compound suitable for X-ray structure determination were recrystallized from acetone by slow evaporation of the solvent at room temperature over several days (m.p. 469–471 K).

S3. Refinement

All the H atoms were positioned geometrically and refined using a riding model with C—H = 0.93 and 0.96 Å. The U_{iso} values were constrained to be $1.5U_{eq}$ of the carrier atom for methyl H atoms and $1.2U_{eq}$ for the remaining H atoms. A rotating group model was used for the methyl groups.




Figure 1

The structure of the title compound, showing 50% probability displacement ellipsoids. H atoms are shown as spheres of arbitrary radius.




Figure 2

The crystal packing, viewed along the *b*-axis, showing the molecules linked into two-dimensional layers parallel to the $(10\bar{4})$ plane. H atoms that not involved in hydrogen bonding (dashed lines) are omitted for clarity.

(1*E*,2*E*)-1,2-Bis[1-(3-nitrophenyl)ethylidene]hydrazine

Crystal data

$C_{16}H_{14}N_4O_4$

$M_r = 326.31$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 3.9296(3)\ \text{\AA}$

$b = 7.4448(5)\ \text{\AA}$

$c = 26.3979(19)\ \text{\AA}$

$\beta = 94.022(1)^\circ$

$V = 770.37 (10) \text{ \AA}^3$
 $Z = 2$
 $F(000) = 340$
 $D_x = 1.407 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 4585 reflections

$\theta = 2.8\text{--}28.3^\circ$
 $\mu = 0.10 \text{ mm}^{-1}$
 $T = 296 \text{ K}$
 Block, yellow
 $0.34 \times 0.17 \times 0.10 \text{ mm}$

Data collection

Bruker APEX DUO CCD area-detector diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.966$, $T_{\max} = 0.990$

15392 measured reflections
 2254 independent reflections
 1686 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$
 $\theta_{\max} = 30.1^\circ$, $\theta_{\min} = 1.6^\circ$
 $h = -5 \rightarrow 5$
 $k = -10 \rightarrow 10$
 $l = -37 \rightarrow 37$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.146$
 $S = 1.06$
 2254 reflections
 110 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.069P)^2 + 0.1466P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.21 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.18 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.9642 (5)	0.1248 (2)	0.20045 (5)	0.1005 (6)
O2	0.8102 (4)	0.06399 (18)	0.12349 (5)	0.0867 (5)
N1	0.0796 (3)	0.47453 (16)	0.02358 (4)	0.0461 (3)
N2	0.8220 (3)	0.16404 (18)	0.15972 (5)	0.0572 (3)
C1	0.3995 (4)	0.6788 (2)	0.14058 (5)	0.0507 (3)
H1A	0.3117	0.7942	0.1362	0.061*
C2	0.5585 (4)	0.6293 (2)	0.18705 (6)	0.0583 (4)
H2A	0.5726	0.7113	0.2137	0.070*
C3	0.6956 (4)	0.4607 (2)	0.19421 (5)	0.0525 (4)
H3A	0.8039	0.4272	0.2252	0.063*
C4	0.6669 (3)	0.34261 (18)	0.15379 (5)	0.0429 (3)

C5	0.5072 (3)	0.38640 (17)	0.10740 (4)	0.0400 (3)
H5A	0.4914	0.3029	0.0811	0.048*
C6	0.3697 (3)	0.55758 (17)	0.10032 (4)	0.0390 (3)
C7	0.1991 (3)	0.60690 (18)	0.05020 (5)	0.0399 (3)
C8	0.1829 (5)	0.7991 (2)	0.03461 (6)	0.0643 (4)
H8A	0.1410	0.8068	-0.0016	0.096*
H8B	0.3956	0.8567	0.0447	0.096*
H8C	0.0016	0.8578	0.0507	0.096*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.1420 (14)	0.0814 (10)	0.0708 (9)	0.0246 (9)	-0.0438 (9)	0.0216 (7)
O2	0.1242 (12)	0.0551 (7)	0.0763 (9)	0.0281 (7)	-0.0242 (8)	-0.0045 (6)
N1	0.0550 (6)	0.0460 (6)	0.0357 (5)	0.0002 (5)	-0.0080 (4)	0.0071 (4)
N2	0.0638 (8)	0.0518 (7)	0.0535 (7)	0.0030 (6)	-0.0133 (6)	0.0126 (6)
C1	0.0561 (8)	0.0465 (7)	0.0488 (8)	0.0046 (6)	-0.0020 (6)	-0.0061 (6)
C2	0.0696 (10)	0.0617 (9)	0.0424 (7)	-0.0022 (7)	-0.0053 (6)	-0.0130 (6)
C3	0.0576 (8)	0.0632 (9)	0.0350 (6)	-0.0071 (7)	-0.0088 (5)	0.0015 (6)
C4	0.0440 (6)	0.0450 (7)	0.0388 (6)	-0.0030 (5)	-0.0044 (5)	0.0058 (5)
C5	0.0430 (6)	0.0426 (7)	0.0338 (6)	-0.0013 (5)	-0.0024 (4)	0.0006 (5)
C6	0.0383 (6)	0.0425 (6)	0.0357 (6)	0.0002 (5)	-0.0007 (4)	0.0015 (5)
C7	0.0391 (6)	0.0426 (7)	0.0376 (6)	0.0045 (5)	0.0003 (4)	0.0042 (5)
C8	0.0847 (11)	0.0450 (8)	0.0600 (9)	0.0046 (8)	-0.0171 (8)	0.0064 (7)

Geometric parameters (Å, °)

O1—N2	1.2121 (16)	C3—C4	1.3808 (19)
O2—N2	1.2105 (19)	C3—H3A	0.9300
N1—C7	1.2803 (17)	C4—C5	1.3756 (17)
N1—N1 ⁱ	1.406 (2)	C5—C6	1.3916 (18)
N2—C4	1.4663 (19)	C5—H5A	0.9300
C1—C2	1.387 (2)	C6—C7	1.4867 (17)
C1—C6	1.3930 (19)	C7—C8	1.489 (2)
C1—H1A	0.9300	C8—H8A	0.9600
C2—C3	1.374 (2)	C8—H8B	0.9600
C2—H2A	0.9300	C8—H8C	0.9600
C7—N1—N1 ⁱ	113.69 (14)	C4—C5—C6	119.02 (12)
O2—N2—O1	122.84 (15)	C4—C5—H5A	120.5
O2—N2—C4	118.75 (12)	C6—C5—H5A	120.5
O1—N2—C4	118.38 (14)	C5—C6—C1	118.67 (11)
C2—C1—C6	120.76 (14)	C5—C6—C7	119.49 (11)
C2—C1—H1A	119.6	C1—C6—C7	121.84 (12)
C6—C1—H1A	119.6	N1—C7—C6	115.05 (11)
C3—C2—C1	120.81 (14)	N1—C7—C8	125.53 (12)
C3—C2—H2A	119.6	C6—C7—C8	119.41 (12)
C1—C2—H2A	119.6	C7—C8—H8A	109.5

C2—C3—C4	117.74 (12)	C7—C8—H8B	109.5
C2—C3—H3A	121.1	H8A—C8—H8B	109.5
C4—C3—H3A	121.1	C7—C8—H8C	109.5
C5—C4—C3	122.99 (13)	H8A—C8—H8C	109.5
C5—C4—N2	118.04 (12)	H8B—C8—H8C	109.5
C3—C4—N2	118.94 (12)		
C6—C1—C2—C3	1.1 (2)	C4—C5—C6—C1	0.12 (19)
C1—C2—C3—C4	-0.5 (2)	C4—C5—C6—C7	179.49 (11)
C2—C3—C4—C5	-0.3 (2)	C2—C1—C6—C5	-0.9 (2)
C2—C3—C4—N2	177.71 (14)	C2—C1—C6—C7	179.72 (13)
O2—N2—C4—C5	1.6 (2)	N1 ⁱ —N1—C7—C6	-179.90 (13)
O1—N2—C4—C5	179.73 (15)	N1 ⁱ —N1—C7—C8	-0.7 (2)
O2—N2—C4—C3	-176.58 (16)	C5—C6—C7—N1	24.84 (17)
O1—N2—C4—C3	1.6 (2)	C1—C6—C7—N1	-155.80 (13)
C3—C4—C5—C6	0.5 (2)	C5—C6—C7—C8	-154.43 (14)
N2—C4—C5—C6	-177.55 (11)	C1—C6—C7—C8	24.9 (2)

Symmetry code: (i) $-x, -y+1, -z$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C3—H3A...O1 ⁱⁱ	0.93	2.57	3.239 (2)	129

Symmetry code: (ii) $-x+2, y+1/2, -z+1/2$.