

# 6-[(2-Chloropyridin-5-ylmethyl)(ethyl)azanyl]-4-(2-fluorophenyl)-1-methyl-5-nitro-1,2,3,4-tetrahydropyridin-2-one

Chuan-Wen Sun,\* Jing Wang and Ying Wu

Department of Chemistry, College of Life and Environmental Science, Shanghai Normal University, Shanghai 200234, People's Republic of China  
Correspondence e-mail: jingwangchemistry@163.com


Received 5 January 2012; accepted 9 February 2012

Key indicators: single-crystal X-ray study;  $T = 298\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$ ;  $R$  factor = 0.070;  $wR$  factor = 0.151; data-to-parameter ratio = 12.8.

In the title compound,  $\text{C}_{20}\text{H}_{20}\text{ClFN}_4\text{O}_3$ , the tetrahydropyridone ring adopts a skew boat conformation. The dihedral angle between the mean planes of the benzene and pyridine rings is  $80.7(3)^\circ$ . In the crystal, weak  $\text{C}-\text{H}\cdots\text{O}$  interactions are observed.

## Related literature

For general background to neonicotinoid compounds and their application as insecticides, see: Jeschke & Nauen (2008); Kagabu & Matsuno (1997); Ohno *et al.* (2009); Shao *et al.* (2008); Tian *et al.* (2007); Tomizawa & Casida (2009). For the synthesis of the title compound, see: Zhang *et al.* (2010). For puckering parameters, see Cremer & Pople (1975).



## Experimental

### Crystal data

$\text{C}_{20}\text{H}_{20}\text{ClFN}_4\text{O}_3$   
 $M_r = 418.85$   
Triclinic,  $P\bar{1}$   
 $a = 6.750(2)\text{ \AA}$   
 $b = 8.262(3)\text{ \AA}$   
 $c = 17.853(6)\text{ \AA}$

$\alpha = 94.981(6)^\circ$   
 $\beta = 91.302(7)^\circ$   
 $\gamma = 100.593(7)^\circ$   
 $V = 974.2(6)\text{ \AA}^3$   
 $Z = 2$   
Mo  $K\alpha$  radiation

$\mu = 0.24\text{ mm}^{-1}$   
 $T = 298\text{ K}$

$0.23 \times 0.20 \times 0.20\text{ mm}$

### Data collection

Bruker SMART CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2003)  
 $T_{\min} = 0.938$ ,  $T_{\max} = 0.954$

5691 measured reflections  
3391 independent reflections  
1943 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.073$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.070$   
 $wR(F^2) = 0.151$   
 $S = 0.96$   
3391 reflections

264 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.23\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C8—H8C $\cdots$ O1 <sup>i</sup>	0.96	2.46	3.383 (5)	161
C6—H6A $\cdots$ O3 <sup>ii</sup>	0.97	2.57	3.506 (5)	161
C3—H3 $\cdots$ O1	0.93	2.53	3.320 (5)	143
C2—H2 $\cdots$ O3 <sup>iii</sup>	0.93	2.46	3.344 (5)	159

Symmetry codes: (i)  $x + 1, y, z$ ; (ii)  $x, y + 1, z$ ; (iii)  $-x + 1, -y, -z + 2$ .

Data collection: *SMART* (Bruker, 2001); cell refinement: *SMART*; data reduction: *SAINT* (Bruker, 2001); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported mostly by the National Natural Science Foundation of China (grant Nos 21042010, 21102092 and 30870560).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JJ2119).

## References

- Bruker (2001). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Jeschke, P. & Nauen, R. (2008). *Pest Manage. Sci.* **64**, 1084–1098.
- Kagabu, S. & Matsuno, H. (1997). *J. Agric. Food Chem.* **45**, 276–281.
- Ohno, I., Tomizawa, M., Durkin, K. A., Naruse, Y., Casida, J. E. & Kagabu, S. (2009). *Chem. Res. Toxicol.* **22**, 476–482.
- Shao, X. S., Zhang, W. W., Peng, Y. Q., Li, Z., Tian, Z. Z. & Qian, X. H. (2008). *Bioorg. Med. Chem. Lett.* **18**, 6513–6516.
- Sheldrick, G. M. (2003). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Tian, Z. Z., Shao, X. S., Li, Z., Qian, X. H. & Huang, Q. C. (2007). *J. Agric. Food Chem.* **55**, 2288–2292.
- Tomizawa, M. & Casida, J. E. (2009). *Acc. Chem. Res.* **42**, 260–269.
- Zhang, W. W., Yang, X. B., Chen, W. D., Xu, X. Y., Li, L., Zhai, H. B. & Li, Z. (2010). *J. Agric. Food Chem.* **58**, 2741–2745.

# supporting information

*Acta Cryst.* (2012). E68, o753 [doi:10.1107/S1600536812005776]

## **6-[(2-Chloropyridin-5-ylmethyl)(ethyl)azanyl]-4-(2-fluorophenyl)-1-methyl-5-nitro-1,2,3,4-tetrahydropyridin-2-one**

**Chuan-Wen Sun, Jing Wang and Ying Wu**

### **S1. Comment**

In recent years, neonicotinoid insecticide compounds have rapidly grown and become a new chemical class of insecticides because of their novel structure and mode of action (Jeschke & Nauen, 2008; Kagabu & Matsuno, 1997; Ohno *et al.*, 2009; Shao *et al.*, 2008; Tian *et al.*, 2007; Tomizawa & Casida, 2009). We report here the synthesis and crystal structure of one of these compounds,  $C_{20}H_{20}ClFN_4O_3$ , the title compound, (I).


In (I) the tetrahydropyridone ring adopts a skew boat conformation with puckering parameters (Cremer & Pople, 1975) Q,  $\theta$  and  $\varphi$  of  $0.526(4)\text{\AA}$ ,  $113.2(4)^\circ$  and  $22.0(4)^\circ$  and a N2/C9/C10/N4 dihedral angle of  $-21.6(6)^\circ$  (Fig. 1). In the tetrahydropyridone moiety, C11—C10, C12—C11 and C13—C12 bond lengths ( $1.499(5)\text{\AA}$ ,  $1.527(5)\text{\AA}$  and  $1.503(5)\text{\AA}$ ) are slightly shorter than normal. The C13—N3, C9—N3, and C10—N4 bond lengths ( $1.391(4)\text{\AA}$ ,  $1.417(4)\text{\AA}$ , and  $1.422(4)\text{\AA}$ , respectively) are slightly shorter than normal while the C9=C10 bond length ( $1.360(4)\text{\AA}$ ) is slightly longer than normal. The dihedral angle between the mean planes of the benzene and pyridine rings is  $80.7(3)^\circ$ . In the crystal structure weak C2—H2 $\cdots$ O3, C8—H8C $\cdots$ O1 and C6—H6A $\cdots$ O3 intermolecular interactions are observed which may influence crystal packing (Fig. 2).

### **S2. Experimental**


In the preparation of the title compound, a solution of 2-fluorobenzaldehyde (15 mmol), Meldrum's acid (15 mmol) in ethanol (30 ml), with piperidine (0.1 mmol) used as catalyst, was added dropwise and the solution was stirred at room temperature for 2 h. Nitenpyram (2.75 g, 10 mmol) was added to the reaction mixture, heated to  $65\text{ }^\circ\text{C}$  for 6 h and then cooled to room temperature. The reaction mixture was concentrated under reduced pressure and treated with 20 ml of water whereby the solution was extracted three times with  $\text{CH}_2\text{Cl}_2$ , and the combined extracts were dried over  $\text{MgSO}_4$ . The organic phase was evaporated under reduced pressure and the crude product was subjected to flash chromatography on silica gel, eluting with ethyl acetate /petroleum ether to afford pure yellow crystals (yield 81%). Anal. calcd. for  $C_{20}H_{20}ClFN_4O_3$  C 57.33, H 4.80, N 13.36% found, C 57.35, H 4.81, N 13.38%.

### **S3. Refinement**

All H atoms bonded to C were positioned geometrically [C—H =  $0.93\text{ \AA}$  (aromatic),  $0.97\text{ \AA}$  (methylene) and  $0.96\text{ \AA}$  (methyl)] and refined in the riding mode with [ $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{aromatic and methylene C})$  and  $1.5U_{\text{eq}}(\text{methyl C})$ ].

**Figure 1**

The molecular structure of (I), showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

**Figure 2**

Packing of the title compound, (I), viewed along the  $a$  axis. Weak C2—H2···O3, C8—H8C···O1 and C6—H6A···O3 intermolecular interactions are shown as dashed lines. H atoms not involved in hydrogen bonding have been omitted for clarity.

### **6-[(2-Chloropyridin-5-ylmethyl)(ethyl)azanyl]-4-(2-fluorophenyl)-1-methyl-5-nitro-1,2,3,4-tetrahydropyridin-2-one**

#### *Crystal data*



$M_r = 418.85$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a = 6.750(2)$  Å

$b = 8.262(3)$  Å

$c = 17.853(6)$  Å

$\alpha = 94.981(6)^\circ$

$\beta = 91.302(7)^\circ$

$\gamma = 100.593(7)^\circ$

$V = 974.2(6)$  Å<sup>3</sup>

$Z = 2$

$F(000) = 436$

$D_x = 1.428 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 906 reflections

$\theta = 2.3\text{--}19.9^\circ$

$\mu = 0.24 \text{ mm}^{-1}$

$T = 298$  K

Block, yellow

$0.23 \times 0.20 \times 0.20$  mm

#### *Data collection*

Bruker SMART CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and  $\omega$  scans

Absorption correction: multi-scan  
(SADABS; Sheldrick, 2003)

$T_{\min} = 0.938$ ,  $T_{\max} = 0.954$

5691 measured reflections

3391 independent reflections

1943 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.073$   
 $\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 2.3^\circ$   
 $h = -7 \rightarrow 8$

$k = -9 \rightarrow 9$   
 $l = -21 \rightarrow 16$

### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.070$   
 $wR(F^2) = 0.151$   
 $S = 0.96$   
3391 reflections  
264 parameters  
0 restraints  
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map  
Hydrogen site location: inferred from neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0497P)^2]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} < 0.001$   
 $\Delta\rho_{\text{max}} = 0.23 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.25 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	-0.0482 (2)	0.19595 (18)	1.06337 (7)	0.0963 (5)
F1	0.0556 (4)	-0.3141 (3)	0.57751 (14)	0.0768 (8)
O1	0.1183 (4)	0.3810 (3)	0.70839 (17)	0.0579 (8)
O2	0.0164 (4)	0.1954 (4)	0.61628 (17)	0.0745 (10)
O3	0.4797 (4)	-0.1603 (3)	0.82425 (15)	0.0613 (8)
N1	0.0698 (5)	0.3252 (5)	0.9407 (2)	0.0677 (11)
N2	0.5075 (4)	0.3694 (3)	0.76441 (15)	0.0358 (7)
N3	0.5073 (4)	0.0931 (3)	0.78309 (15)	0.0367 (7)
N4	0.1287 (5)	0.2474 (4)	0.67181 (19)	0.0457 (8)
C1	0.1333 (7)	0.2582 (5)	0.9985 (2)	0.0579 (12)
C2	0.3272 (8)	0.2367 (5)	1.0112 (2)	0.0620 (13)
H2	0.3632	0.1866	1.0528	0.074*
C3	0.2101 (7)	0.3769 (5)	0.8917 (2)	0.0595 (12)
H3	0.1690	0.4245	0.8501	0.071*
C4	0.4110 (6)	0.3647 (4)	0.8982 (2)	0.0431 (10)
C5	0.4664 (6)	0.2921 (5)	0.9599 (2)	0.0539 (11)
H5	0.5998	0.2804	0.9669	0.065*
C6	0.5569 (6)	0.4366 (4)	0.8427 (2)	0.0475 (11)
H6A	0.5631	0.5553	0.8461	0.057*
H6B	0.6899	0.4176	0.8567	0.057*
C7	0.5241 (6)	0.4907 (4)	0.7083 (2)	0.0468 (10)
H7A	0.4291	0.5638	0.7190	0.056*

H7B	0.4883	0.4331	0.6588	0.056*
C8	0.7339 (6)	0.5922 (5)	0.7079 (3)	0.0664 (13)
H8A	0.7682	0.6524	0.7563	0.100*
H8B	0.7388	0.6686	0.6701	0.100*
H8C	0.8283	0.5204	0.6970	0.100*
C9	0.4199 (5)	0.2097 (4)	0.74628 (18)	0.0333 (9)
C10	0.2653 (5)	0.1481 (4)	0.6955 (2)	0.0360 (9)
C11	0.2062 (5)	-0.0340 (4)	0.6735 (2)	0.0387 (9)
H11	0.0674	-0.0550	0.6527	0.046*
C12	0.2044 (5)	-0.1165 (4)	0.7469 (2)	0.0444 (10)
H12A	0.0999	-0.0843	0.7782	0.053*
H12B	0.1738	-0.2357	0.7359	0.053*
C13	0.4050 (6)	-0.0674 (5)	0.78885 (19)	0.0401 (9)
C14	0.7233 (5)	0.1245 (5)	0.8024 (2)	0.0539 (11)
H14A	0.7442	0.1457	0.8560	0.081*
H14B	0.7887	0.2190	0.7787	0.081*
H14C	0.7789	0.0297	0.7852	0.081*
C15	0.3356 (5)	-0.1057 (4)	0.6151 (2)	0.0371 (9)
C16	0.2525 (6)	-0.2468 (5)	0.5697 (2)	0.0491 (11)
C17	0.3578 (7)	-0.3229 (5)	0.5167 (2)	0.0621 (13)
H17	0.2958	-0.4175	0.4870	0.074*
C18	0.5569 (7)	-0.2557 (5)	0.5088 (2)	0.0590 (12)
H18	0.6323	-0.3060	0.4740	0.071*
C19	0.6450 (6)	-0.1144 (5)	0.5521 (2)	0.0500 (11)
H19	0.7796	-0.0685	0.5462	0.060*
C20	0.5345 (6)	-0.0401 (5)	0.6044 (2)	0.0418 (10)
H20	0.5957	0.0563	0.6330	0.050*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.1069 (11)	0.1166 (12)	0.0622 (9)	0.0018 (9)	0.0239 (8)	0.0258 (8)
F1	0.0722 (17)	0.0626 (16)	0.0813 (18)	-0.0162 (13)	0.0080 (14)	-0.0141 (14)
O1	0.0526 (17)	0.0505 (18)	0.077 (2)	0.0280 (15)	-0.0045 (15)	0.0027 (16)
O2	0.074 (2)	0.072 (2)	0.077 (2)	0.0232 (17)	-0.0420 (18)	-0.0028 (17)
O3	0.087 (2)	0.0429 (17)	0.0585 (19)	0.0198 (16)	-0.0108 (16)	0.0156 (15)
N1	0.062 (2)	0.098 (3)	0.044 (2)	0.014 (2)	0.0000 (19)	0.013 (2)
N2	0.0406 (18)	0.0324 (17)	0.0344 (17)	0.0059 (14)	0.0014 (14)	0.0049 (14)
N3	0.0360 (17)	0.0358 (17)	0.0382 (18)	0.0071 (14)	-0.0067 (14)	0.0046 (14)
N4	0.0395 (19)	0.045 (2)	0.054 (2)	0.0081 (16)	-0.0040 (17)	0.0101 (17)
C1	0.073 (3)	0.059 (3)	0.038 (3)	0.006 (2)	0.003 (2)	-0.001 (2)
C2	0.089 (4)	0.059 (3)	0.038 (3)	0.010 (3)	-0.007 (2)	0.010 (2)
C3	0.067 (3)	0.077 (3)	0.037 (2)	0.015 (3)	-0.002 (2)	0.013 (2)
C4	0.050 (3)	0.040 (2)	0.036 (2)	0.008 (2)	-0.0077 (19)	-0.0111 (18)
C5	0.064 (3)	0.056 (3)	0.042 (3)	0.015 (2)	-0.012 (2)	0.004 (2)
C6	0.055 (3)	0.037 (2)	0.046 (2)	0.0004 (19)	-0.008 (2)	-0.0034 (18)
C7	0.056 (3)	0.037 (2)	0.050 (2)	0.0093 (19)	0.004 (2)	0.0132 (19)
C8	0.059 (3)	0.054 (3)	0.089 (4)	0.007 (2)	0.017 (3)	0.028 (3)

C9	0.036 (2)	0.035 (2)	0.031 (2)	0.0089 (17)	0.0030 (17)	0.0075 (16)
C10	0.033 (2)	0.035 (2)	0.041 (2)	0.0104 (17)	-0.0037 (17)	0.0038 (17)
C11	0.032 (2)	0.040 (2)	0.042 (2)	0.0021 (17)	-0.0060 (17)	0.0020 (18)
C12	0.046 (2)	0.040 (2)	0.045 (2)	0.0019 (18)	0.0107 (19)	0.0051 (18)
C13	0.056 (3)	0.039 (2)	0.026 (2)	0.012 (2)	0.0009 (18)	0.0049 (17)
C14	0.043 (2)	0.052 (3)	0.068 (3)	0.014 (2)	-0.010 (2)	0.003 (2)
C15	0.042 (2)	0.035 (2)	0.035 (2)	0.0100 (18)	-0.0018 (17)	0.0062 (17)
C16	0.054 (3)	0.041 (2)	0.048 (3)	-0.003 (2)	-0.001 (2)	0.006 (2)
C17	0.089 (4)	0.046 (3)	0.049 (3)	0.013 (3)	0.011 (3)	-0.006 (2)
C18	0.076 (3)	0.065 (3)	0.044 (3)	0.031 (3)	0.009 (2)	0.010 (2)
C19	0.054 (3)	0.064 (3)	0.037 (2)	0.022 (2)	0.001 (2)	0.009 (2)
C20	0.049 (2)	0.046 (2)	0.032 (2)	0.011 (2)	-0.0010 (18)	0.0057 (18)

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

C11—C1	1.744 (4)	C7—H7B	0.9700
F1—C16	1.358 (4)	C8—H8A	0.9600
O1—N4	1.247 (4)	C8—H8B	0.9600
O2—N4	1.231 (4)	C8—H8C	0.9600
O3—C13	1.206 (4)	C9—C10	1.360 (4)
N1—C1	1.312 (5)	C10—C11	1.499 (5)
N1—C3	1.342 (5)	C11—C15	1.526 (5)
N2—C9	1.352 (4)	C11—C12	1.527 (5)
N2—C6	1.467 (4)	C11—H11	0.9800
N2—C7	1.469 (4)	C12—C13	1.503 (5)
N3—C13	1.391 (4)	C12—H12A	0.9700
N3—C9	1.417 (4)	C12—H12B	0.9700
N3—C14	1.461 (4)	C14—H14A	0.9600
N4—C10	1.422 (4)	C14—H14B	0.9600
C1—C2	1.370 (6)	C14—H14C	0.9600
C2—C5	1.373 (5)	C15—C20	1.376 (5)
C2—H2	0.9300	C15—C16	1.381 (5)
C3—C4	1.381 (5)	C16—C17	1.373 (5)
C3—H3	0.9300	C17—C18	1.372 (5)
C4—C5	1.374 (5)	C17—H17	0.9300
C4—C6	1.495 (5)	C18—C19	1.371 (5)
C5—H5	0.9300	C18—H18	0.9300
C6—H6A	0.9700	C19—C20	1.381 (5)
C6—H6B	0.9700	C19—H19	0.9300
C7—C8	1.508 (5)	C20—H20	0.9300
C7—H7A	0.9700		
C1—N1—C3	115.8 (4)	N2—C9—N3	115.0 (3)
C9—N2—C6	121.8 (3)	C10—C9—N3	116.8 (3)
C9—N2—C7	121.2 (3)	C9—C10—N4	121.7 (3)
C6—N2—C7	116.1 (3)	C9—C10—C11	121.2 (3)
C13—N3—C9	122.3 (3)	N4—C10—C11	116.0 (3)
C13—N3—C14	115.9 (3)	C10—C11—C15	115.7 (3)

C9—N3—C14	120.6 (3)	C10—C11—C12	105.8 (3)
O2—N4—O1	121.0 (3)	C15—C11—C12	112.4 (3)
O2—N4—C10	118.5 (3)	C10—C11—H11	107.5
O1—N4—C10	120.4 (3)	C15—C11—H11	107.5
N1—C1—C2	125.1 (4)	C12—C11—H11	107.5
N1—C1—Cl1	115.6 (4)	C13—C12—C11	110.8 (3)
C2—C1—Cl1	119.3 (4)	C13—C12—H12A	109.5
C1—C2—C5	117.3 (4)	C11—C12—H12A	109.5
C1—C2—H2	121.4	C13—C12—H12B	109.5
C5—C2—H2	121.4	C11—C12—H12B	109.5
N1—C3—C4	125.0 (4)	H12A—C12—H12B	108.1
N1—C3—H3	117.5	O3—C13—N3	120.4 (3)
C4—C3—H3	117.5	O3—C13—C12	123.4 (3)
C5—C4—C3	116.0 (4)	N3—C13—C12	116.2 (3)
C5—C4—C6	123.5 (4)	N3—C14—H14A	109.5
C3—C4—C6	120.3 (4)	N3—C14—H14B	109.5
C2—C5—C4	120.8 (4)	H14A—C14—H14B	109.5
C2—C5—H5	119.6	N3—C14—H14C	109.5
C4—C5—H5	119.6	H14A—C14—H14C	109.5
N2—C6—C4	114.6 (3)	H14B—C14—H14C	109.5
N2—C6—H6A	108.6	C20—C15—C16	116.4 (4)
C4—C6—H6A	108.6	C20—C15—C11	124.4 (3)
N2—C6—H6B	108.6	C16—C15—C11	119.2 (3)
C4—C6—H6B	108.6	F1—C16—C17	118.1 (4)
H6A—C6—H6B	107.6	F1—C16—C15	118.4 (4)
N2—C7—C8	112.1 (3)	C17—C16—C15	123.6 (4)
N2—C7—H7A	109.2	C18—C17—C16	118.3 (4)
C8—C7—H7A	109.2	C18—C17—H17	120.9
N2—C7—H7B	109.2	C16—C17—H17	120.9
C8—C7—H7B	109.2	C19—C18—C17	120.1 (4)
H7A—C7—H7B	107.9	C19—C18—H18	119.9
C7—C8—H8A	109.5	C17—C18—H18	119.9
C7—C8—H8B	109.5	C18—C19—C20	120.2 (4)
H8A—C8—H8B	109.5	C18—C19—H19	119.9
C7—C8—H8C	109.5	C20—C19—H19	119.9
H8A—C8—H8C	109.5	C15—C20—C19	121.4 (4)
H8B—C8—H8C	109.5	C15—C20—H20	119.3
N2—C9—C10	128.1 (3)	C19—C20—H20	119.3
C3—N1—C1—C2	-0.7 (7)	O2—N4—C10—C11	-24.8 (5)
C3—N1—C1—Cl1	178.5 (3)	O1—N4—C10—C11	152.5 (3)
N1—C1—C2—C5	1.1 (7)	C9—C10—C11—C15	-80.5 (4)
Cl1—C1—C2—C5	-178.1 (3)	N4—C10—C11—C15	111.0 (3)
C1—N1—C3—C4	-0.2 (7)	C9—C10—C11—C12	44.6 (4)
N1—C3—C4—C5	0.7 (6)	N4—C10—C11—C12	-123.9 (3)
N1—C3—C4—C6	-175.9 (4)	C10—C11—C12—C13	-56.2 (4)
C1—C2—C5—C4	-0.5 (6)	C15—C11—C12—C13	70.9 (4)
C3—C4—C5—C2	-0.3 (6)	C9—N3—C13—O3	-175.8 (3)

C6—C4—C5—C2	176.2 (3)	C14—N3—C13—O3	17.0 (5)
C9—N2—C6—C4	-34.4 (5)	C9—N3—C13—C12	6.1 (5)
C7—N2—C6—C4	134.8 (3)	C14—N3—C13—C12	-161.1 (3)
C5—C4—C6—N2	124.5 (4)	C11—C12—C13—O3	-143.5 (4)
C3—C4—C6—N2	-59.1 (5)	C11—C12—C13—N3	34.5 (4)
C9—N2—C7—C8	-131.8 (4)	C10—C11—C15—C20	27.1 (5)
C6—N2—C7—C8	58.8 (4)	C12—C11—C15—C20	-94.5 (4)
C6—N2—C9—C10	136.3 (4)	C10—C11—C15—C16	-153.7 (3)
C7—N2—C9—C10	-32.4 (5)	C12—C11—C15—C16	84.6 (4)
C6—N2—C9—N3	-46.4 (4)	C20—C15—C16—F1	-179.0 (3)
C7—N2—C9—N3	144.8 (3)	C11—C15—C16—F1	1.8 (5)
C13—N3—C9—N2	160.7 (3)	C20—C15—C16—C17	0.9 (6)
C14—N3—C9—N2	-32.6 (4)	C11—C15—C16—C17	-178.4 (4)
C13—N3—C9—C10	-21.8 (5)	F1—C16—C17—C18	-179.8 (4)
C14—N3—C9—C10	144.9 (3)	C15—C16—C17—C18	0.4 (6)
N2—C9—C10—N4	-21.6 (6)	C16—C17—C18—C19	-1.2 (6)
N3—C9—C10—N4	161.2 (3)	C17—C18—C19—C20	0.7 (6)
N2—C9—C10—C11	170.6 (3)	C16—C15—C20—C19	-1.4 (5)
N3—C9—C10—C11	-6.6 (5)	C11—C15—C20—C19	177.8 (3)
O2—N4—C10—C9	166.7 (3)	C18—C19—C20—C15	0.7 (5)
O1—N4—C10—C9	-16.0 (5)		

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
C8—H8C···O1 <sup>i</sup>	0.96	2.46	3.383 (5)	161
C6—H6A···O3 <sup>ii</sup>	0.97	2.57	3.506 (5)	161
C3—H3···O1	0.93	2.53	3.320 (5)	143
C2—H2···O3 <sup>iii</sup>	0.93	2.46	3.344 (5)	159

Symmetry codes: (i)  $x+1, y, z$ ; (ii)  $x, y+1, z$ ; (iii)  $-x+1, -y, -z+2$ .